

Electrolyte Disturbances Among Acute Stroke Patients in Manado, Indonesia

by Mieke Kembuan 01

Submission date: 13-Jun-2019 08:34AM (UTC+0700)

Submission ID: 1143105319

File name: isturbances_Among_Acute_Stroke_Patients_in_Manado,_Indonesia.pdf (521.89K)

Word count: 3012

Character count: 15207



1 Electrolyte disturbances among acute stroke patients in Manado, Indonesia

Mieke A.H.N. Kembuan¹, Sekplin A.S. Sekeon²

ABSTRACT

Stroke is known as main public health problem. It is one of the leading causes of chronic disability and death. Both sodium and potassium disturbances have negative influences on the outcome of stroke. There is a scarcity of data about electrolyte disturbances in acute stroke, especially from developing countries. The objectives of this study were to find out the incidence of electrolyte disturbances among acute stroke patients; and their association with severity of acute stroke. This study was a hospital-based cross-sectional study. Samples consisted of 85 patients that meet the inclusion criteria. Data of electrolyte status was obtained from medical record. The result shows that the incidence of electrolyte disturbances among acute stroke was 45.9%. Sodium disturbances was 8.2% and potassium disturbances was 40.0% from all study samples. Mean of NIHSS score on admission was 9.3 (95% CI 8.2-10.2). Mean of sodium was 141.7 (95% CI 140.7-142.7) and potassium was 3.7 (95% CI 3.0-6.0). There was no association between electrolyte disturbances and severity of stroke ($p > 0.05$). Conclusion: The incidence of electrolyte status disorders in acute stroke patients was high but we could not detect any association between electrolyte disturbances and the severity of acute stroke.

Keyword: "electrolyte disturbances", sodium, potassium, severity, stroke, "acute stroke", manado

1 INTRODUCTION

Stroke is a major public health problem. It is globally well-distributed and is ranked at the second top cause of death around the world. Stroke causes a great impact on disability rate. Stroke also has enormous contribution to economic and social burden for patients and their family³. In Indonesia, stroke and other cardiovascular diseases are number one cause of death among major hospitals.^{2,3}

1 In almost all neurological disorders, electrolyte disturbances were prominent. Electrolyte disturbance are commonly found in acute stroke

setting⁴. Hyponatremia, hyponatremia and hypokalemia was the commonest type of disturbance.^{5,6} Recently, research with electrolyte disturbances is not only focusing on the neuro-endocrine mechanism but also on its prevalence, risk factors and association with other medical condition⁵.

Even though there are some data about large number of electrolyte disturbances in acute stroke setting, reports on the association between electrolyte imbalance and severity of acute stroke are still in limited number. There is a lack of data about this association especially from developing

13

GJMEDPH 2014; Vol. 3, issue 1

¹ Head of Department
⁴ Department of Neurology
 Faculty of Medicine
 Sam Ratulangi University/ BLU-RSUP
 Prof. dr. R. D. Kandou
 Jalan Raya Malalayang Manado
 Email:
 dr.miekekembuan@yahoo.com

² Resident at Department of
 Neurology, Faculty of Medicine, Sam
 Ratulangi University, Manado,
 Indonesia
 Telephone: +62-431-3359346;
 Email: sekplin@yahoo.com

Conflict of Interest—none

Funding—none

countries. The objectives of this study were to find out the incidence of electrolyte disturbances in acute stroke setting and their association with severity of acute stroke.

MATERIALS AND METHODS

This was a cross-sectional hospital-based study conducted in Manado General Hospital (formally known as Professor R. D. Kandou General Hospital). All stroke patients were treated in neurological wards. Diagnosis of stroke was based on history of disease, physical examination and confirmed by neuro-imaging study. Electrolytes analyzed in this study were sodium and potassium. Sodium level in a range of 135-153 Meq/L was defined as normal. Potassium level in a range of 3.5-5 Meq/L was defined as normal. Other values were categorized as electrolyte disturbances. Hyponatremia was defined when the level below 135 Meq/L. Hipernatremia was the level of natrium above 153 Meq/L. Hypokalemia was the level of kalium below 3.5 Meq/L, while Hiperkalemia was when the level of kalium above 5 Meq/L. Based on NIHSS (Neurological Institute of Health Science for Stroke) score, we defined mild severity of acute stroke when the NIHSS score less than 4, moderate severity if the score fall within 4 to 15 dan score above 15 was defined as the most severe deficit. All the patients who met the inclusion criteria were

recruited as samples. Inclusion criteria were: 1. Patients are not in diuretic therapy, 2. Patient was not in resuscitation phase. 3. There was no history of kidney or endocrine diseases. Exclusion criteria: We excluded all patients with a complication that might affect electrolyte status during their stroke management.

Data collection for demographic characteristics was obtained from interview. Patients with communication difficulties were represented by family member or spouse. Medical data was attained from medical records. One way Anova was employed to find out the association between each electrolyte disturbances and the severity of stroke. P value < 0.05 level was determined as statistically significant association. Statistical analysis was performed with SPSS version 17. Ethical consideration was obtained from Research Committee of Manado General Hospital.

RESULTS

Among 85 patients who were analyzed, 61.2% was male, 37.6% was in above 60 years age group, mean age was 56.3 (95% CI 53.8-58.8), 50% was reach secondary educational level, mostly were Minahasan ethnicity, majority of them were Christian and largely work as housekeeper or farmer (Table 1).

Table 1. Distribution of demographic characteristics and electrolyte status

Characteristics	n=85	%
Demographic		
Male	52	61.2
51-60 age group	23	27.1
>60 age group	32	37.6
High secondary school	40	50
Minahasan ethnic	62	76
Christian	70	85
Housekeeper	38	46
Farmer	12	15
Ischemic stroke	44	53.7
Sodium level		
Hyponatremia	6	7.1
Normal	78	91.8
Hypernatremia	1	1.2
Potassium level		
Hypokalemia	32	37.6
Normal	51	60.0
Hyperkalemia	2	2.4

Sodium disturbance		
Yes	7	8.2
No	78	91.8
Potassium disturbance		
Yes	34	40.0
No	51	60.0
Electrolyte disturbances		
Yes	39	45.9
No	46	54.1
Only sodium disturbance	7	8.2
Only potassium disturbance	34	40.0
Both	2	2.4

Based on electrolyte status, the current study found that most of acute stroke patients admitted with normal sodium and potassium level (91.8% and 60.0%, respectively). Totally, there were 39 patients with electrolyte disturbances (45.9%). 8.2% of all patients had sodium disturbance, 40.0% with potassium disturbance and 2.4% with both sodium and potassium disturbances. Hyponatremia were found in 6 patients (7.1%) and only 1 case of hypernatremia (1.2%). Hypokalemia was 37.6%

among all patients and hyperkalemia was only detected in 2 patients (2.4%) (Table 1).

The present study found mean GCS score was 12.4 (95% CI 11.7-13.0) for all patients. Mean of NIHSS score on admission was 9.3 (95% CI 8.2-10.2). Mean of sodium was 141.7 (95% CI 140.7-142.7) and potassium was 3.7 (95% CI 3.0-6.0) (Table 2).

Table 2. Distribution of GCS, NIHSS score, sodium and potassium level.

Variable	Mean	SD	Minimal-maximal	95% CI
GCS	12.4	2.9	6-15	11.7-13.0
NIHSS	9.3	4.5	2-20	8.3-10.2
Sodium	141.7	4.8	126-154	140.7-142.7
Potassium	3.7	0.6	3.0-6.0	3.5-3.8

Table 3 revealed the distribution of severity of stroke based on NIHSS score. Most of patients were classified as moderate deficit (80.0%). There were 8

patients admitted with mild severity (9.4%) and 9 patients with severe deficit (10.6%).

Table 3. Distribution of severity of stroke based on NIHSS score

Severity	NIHSS score	n=85	%
Mild	<4	8	9.4
Moderate	4-15	68	80.0
Severe	>15	9	10.6

Based on status of electrolyte disturbances, the present study found means of NIHSS score was 8.31 (95% CI 6.9-9.6) among patients with electrolyte imbalance. In other group, mean score of NIHSS

was 10.1 (95% CI 8.7-11.4). There was no significantly difference¹⁰ between means of NIHSS score on both group ($p > 0.05$) (Table 4).

Table 4. Distribution of means of NIHSS score based on status of electrolyte disturbances

Variable	Mean	SD	95% CI	p-value
Yes	8.31	4.1	6.9-9.6	0.27
No	10.1	4.5	8.7-11.4	

Based on severity of acute stroke, this study found the distribution of means of sodium level for mild deficit neurologic was 139.4 Meq/L (95% CI 136.1-142.5), for moderate deficit neurology it was 142.0 Meq/L (95% CI 140.8-143.2) and it was 141.1 Meq/L (95% CI 139.1-143.1) for patient with severe deficit. However, there was no significantly difference of means of sodium and potassium among these

groups ($p > 0.05$). Distribution of potassium level for mild cases was 3.7 Meq/L (95% CI 3.3-4.2), for moderate cases it was 3.7 Meq/L (95% CI 3.5-3.8) and for severe cases it was 3.5 Meq/L (95% CI 3.3-3.7). Again, we could not detect any significant difference between electrolyte disturbances and severity of acute stroke (Table 5).

Table 5. Distribution of means of electrolyte level based on severity of stroke

Variable	Mean	SD	95% CI	p-value
Sodium				
Mild	139.4	3.8	136.1-142.5	0.3
Moderate	142.0	5.0	140.8-143.2	
Severe	141.1	2.6	139.1-143.1	
Potassium				
Mild	3.7	0.5	3.3-4.2	0.7
Moderate	3.7	0.6	3.5-3.8	
Severe	3.5	0.2	3.3-3.7	

Among patients with electrolyte disturbances, moderately severe stroke was commonly found (84.6%) and only 7.7% was identified as severe cases. The same picture was found among all patients without electrolyte disturbances, 76.1% of which admitted to hospital with moderately severe

stroke. Another 13% of cases were severe. However, there is no evident of significantly association between status of electrolyte imbalance and severity of acute stroke ($p > 0.05$) (Table 6).

Table 6. Distribution of patients based on status of electrolyte imbalance and stroke severity

Electrolyte disturbances	Severity of stroke						Total		p-value
	Mild		Moderate		Severe		n	%	
	n	%	n	%	n	%			
Yes	3	7.7	33	84.6	3	7.7	39	100	>0.05
No	5	10.9	35	76.1	6	13.0	46	100	
Total	8	9.4	68	80.0	9	10.6	85	100	

DISCUSSION

Stroke is dominantly occurred at the middle age group or above a ²⁰ is commonly found in male than female. The current study also revealed that most of the patients are male and above 60 year age group. Most of them reach high secondary educational level. The majorities of patients were from Minahasa ethnicity. The mean of GCS was 12.4 (95% CI 11.7-13.0) and most of the case was ischemic stroke. These characteristics were similar with other study which found that most of the stroke patient was male, in the middle age or above and most of them admitted to hospital with ischemic stroke.^{2,4,7}

We recorded that the incidence of electrolyte disturbances in this study was relatively high. If we compare to other study, it is revealed that the result of this incidence was quiet similar. The result in this study showed that only limited number of patients admitted with sodium disturbances, but large number of patients with potassium disturbances. However, we could not detect any association between electrolyte imbalance and severity of stroke.

It is found from other study that mortality rate of stroke patients with electrolyte imbalance was higher than in patients with normal level. This is especially true for hypernatremia because of its contribution on the development of brain edema. In this study hyponatremia was found in 7.1%. Some researcher reported that acute hyponatremia in acute stroke could affect the outcome of stroke negatively due to poor severity or death.^{4,5,6}

Siddiqui⁴ reported a high number percentage of patients with electrolyte imbalance (53%) of their acute stroke patients. Most of them were in acute hemorrhagic stroke. Among all patients in their study, it was revealed that there were 36% patients with sodium disturbance and 31% patients with potassium disturbance. Among patients with sodium disturbance, most of them were hyponatremia (32%). The majority of potassium disturbance was hypokalemia (30%). According to Siddiqui, minimum setting of acute management probably responsible for a large number of patients with electrolyte disturbances documented in their study. In the present study, we documented 45.9% of patients with electrolyte disturbances. Most of

the disturbances in this stud, potassium disturbance (40.0%) while sodium disturbance was only detected in 8.2% patients. Among sodium disturbances, hyponatremia was found in 7.1% patients and hypernatremia was only 1.2%. Huang⁸ reported an incidence of hyponatremia in their study was 11.6%. Siddiqui recorded an incidence of hypernatremia was 4% and hypokalemia was 30%. In this study we found the incidence of hypokalemia was 37.6% and there were 2.4% of cases with hyperkalemia.

Based on the distribution of severity of stroke according to NIHSS score, the present study documented that most of the patients (80%) was in moderate deficit neurology when admitting to the hospital. There were 10.6% patients with severe NIHSS score and small number of patients with mild deficit (9.4%). The mean of sodium in moderate deficit neurology was 142.0 Meq/L (95% CI 140.8-143.2). The mean of potassium in moderate NIHSS score was 3.7 Meq/L (95% CI 3.5-3.8). According to Huang⁷, there are significant contributions of hyponatraemia in the outcome of stroke. They documented the higher death rate of patients with hyponatremic than normonatremic patients after 3 years of first stroke attack. However, Huang also reported that there was no significantly different between the two group in term of mortality rate while in hospital, functional status when discharge¹⁹ and the rate of stroke recurrence. Also, there was no difference in mortality rate in 1 to 3 months after discharge that study. But, Donati⁹ reported the higher mortality rate of stroke cases with hyponatremia than stroke with normal level of sodium. According to Diaz¹⁰ there are two commonest causes of hyponatremia¹⁸ acute brain disorders which are SIADH (Syndrome of Inappropriate Antidiuretic Hormone) and CSW (Cerebral Salt Wasting).

Among sodium disturbance, hypernatremic patients was 1.2%. In general hospital population, the reported hypernatremia data were in between 0.3%-3.5%⁶. Therefore, the result in current study was relatively ¹⁷ line with previous study. Hypernatremia is one of the predictor of mortality in patients treated in ICU, especially if the sodium level exceeded 160 Meq/L⁶. In the present study, among severe deficit neurologic patients, the mean of sodium was 141.1 Meq/L. According to Aiyagari,

most of the causes of hypernatraemia were induced by osmotic agents. Mortality rate will increase with the increasing of sodium peak level.

In this study, we concluded that the incidence of electrolyte status disorders in acute stroke patients was high but we could not detect any association between electrolyte disturbances and the severity of acute stroke. Both sodium and potassium disturbances in the current study were not significantly associated with the severity of acute stroke. Although hyponatremic or hypernatremic in acute stroke patients were associated with poor outcome of stroke, but it was not found in our study. There are several reasons as to why sodium disturbances were not associated with stroke severity in this study. First, it is because the assessment of NIHSS score was done on first admission. According to Huang⁷ higher mortality rate due to hyponatremia was only significant after 3 years follow up. It means, this study should be conducted not only when the patients admitted to hospital but should be followed by a cohort study. Second, most of patients admitted in relatively moderate severity (based on GCS and NIHSS score).

16 LIMITATION OF THE STUDY

There are several limitations of the current study: First, based on sample size, the study had relatively small samples. Second, there was no long-term follow-up of NIHSS score could be carried out.

REFERENCES

1. WHO. The WHO STEPwise approach to stroke surveillance/non communicable diseases and mental health. WHO press. Geneva. 2005.
2. Misbach. Stroke: Aspek diagnostik, patofisiologi dan manajemen. Perdossi press. Jakarta. 2011.
3. Depkes. Indonesia bangun RS Pusat Otak Nasional. Updated Jul 14, 2012, cited Jan 12, 2013. Available online <http://www.depkes.go.id/index.php/berita/pres-s-release/1705-indonesia-bangun-rumah-sakit-pusat-otak-nasional-national-brain-centre-hospital-.html>.
4. Siddiqui MR, Islam QT, Haque MA, Iqbal MJ, Hossain A, Rahman YU et al. Electrolyte status in different type of acute stroke patients and

their correlation with some common clinical presentation. *J Medicine* 2012; 13: 133-137.

5. Lath R. Hyponatremia in neurological diseases in ICU. *Indian J Critical Care Med* 2005; 9(1): 47-50.
6. Aiyagari V, Deibert E, Diringner MN. Hyponatremia in the neurologic intensive care unit: How high is too high? *Journal of Critical Care* 2006; 21: 163-172.
7. Chu JY, Chu JK, Chung AG. Epidemiology of cerebrovascular disease among Chinese-canadian: A ten years retrospective case-mix study. *Neurology Asia* 2006; 11: 13-18.
8. Huang WY, Weng WC, Peng TI, Chien YY, Wu CL, Lee M et al. Association of hyponatremia in acute stroke stage with three-year mortality in patients with first-ever ischemic stroke. *Cerebrovasc Dis* 2012; 34: 55-62.
9. Donati-Genet PCM, Dubuis JM, Rimensberger PC. Acute symptomatic hyponatremia and cerebral salt wasting after head injury: An important clinical entity. *Journal of Pediatric Surgery*. 2001; 36(1): 1094-97
10. Diaz JL, Granados M, Suarez JI. Management of medical complications in the neuroscience critical care unit. In: Suarez JI. *Critical care neurology and neurosurgery*. New Jersey: Humana press inc, 2004.

Electrolyte Disturbances Among Acute Stroke Patients in Manado, Indonesia

ORIGINALITY REPORT

15%

SIMILARITY INDEX

12%

INTERNET SOURCES

6%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	www.sciencepub.net Internet Source	4%
2	Submitted to University of Scranton Student Paper	2%
3	www.aimdrjournal.com Internet Source	1%
4	academic.oup.com Internet Source	1%
5	"Poster Presentations", International Journal of Gynecology & Obstetrics, 2015 Publication	1%
6	Paweena Susantitaphong, Kamal Sewaralthahab, Ethan M. Balk, Bertrand L. Jaber, Nicolaos E. Madias. "Short- and Long-Term Effects of Alkali Therapy in Chronic Kidney Disease: A Systematic Review", American Journal of Nephrology, 2012 Publication	<1%

7	ccforum.biomedcentral.com Internet Source	<1%
8	www.ijhsonline.com Internet Source	<1%
9	Submitted to Southwestern Oklahoma State University Student Paper	<1%
10	www.frontiersin.org Internet Source	<1%
11	Submitted to Our Lady of Holy Cross Student Paper	<1%
12	article.sciencepublishinggroup.com Internet Source	<1%
13	pdfs.semanticscholar.org Internet Source	<1%
14	journal.unnes.ac.id Internet Source	<1%
15	medforum.pk Internet Source	<1%
16	bmcinfectdis.biomedcentral.com Internet Source	<1%
17	A. Yasmin, A. Aryadi, A. Widya, A. P. Suwirya et al. " Original ResearchHypertension in rural area: The determinants of left ventricular	<1%

geometric alteration among hypertensive patients at Kubu Village, Karangasem, Bali

Effectiveness comparison between lisinopril and amlodipine in Melanesian patients with hypertension

An experimental study of β -D-Glucans extract of polysaccharide peptides as anti-inflammation and anti-oxidant in patients at high-risk of atherosclerosis

Echocardiographic findings among hypertensive patients in rural area: The descriptive study at Kubu Village, Karangasem, Bali

Addition of left ventricular ejection fraction to the GRACE score in prediction of short term prognosis in patients with non ST-elevation myocardial infarction

Effect of colchicine on HsCRP and mean platelet volume in acute myocardial infarction

Correlation between GRACE score with spatial QRS-T angle in non-ST elevation myocardial infarction patients

Prevalence of Impaired Kidney Function in Hospitalized Hypertensive Patients in Agoesdjam General Hospital, Ketapang, West Borneo, Indonesia

Inappropriate initial treatment in acute coronary syndrome patients: A gap in cardiovascular care

Correlation between plasma histamine level and atherosclerosis severity based on carotid intima media thickness (CIMT) measurement in patients with stable coronary artery

disease
Clinical characteristics, management and outcome of patients with ST elevation myocardial infarction in Pekanbaru city: Preliminary result of single center experience
Phase II cardiac rehabilitation program increases functional capacity in post atrial septal defect surgical closure in adult patients
The identification of human telomerase catalytic subunit in cervical cancer patients and its potential usage as strategy to reawaken hibernating myocardium post myocardial infarction
Could EURO heart risk score predict the severity of coronary lesion?
Pericardial effusion in systemic lupus erythematosus
Intracardiac metastasis of hepatocellular carcinoma
Myocardial bridging assessed by computed tomography scan: A cause of chest pain in patients without obstructed coronary arteries
Correlation between resting heart rate and 6-Minute Walk Test distance in systolic heart failure patients
Severe pulmonary hypertension is a predictor of mortality in pregnant unoperated atrial septal defect
High plasma level of soluble ST2 associated with adverse outcomes during acute phase of ST-elevation myocardial infarction
Effect of extracorporeal shockwave myocardial revascularization on spatial QRS – T angle resolution in stable coronary artery

disease patients
Fragmented QRS study, does it have any structural abnormality relation?
Risk of coronary artery lesions severity to erectile dysfunction incidence in stable coronary heart disease patients
Clinical profiles and outcomes of patients with ST-Elevation Acute Coronary Syndrome (STE-ACS) in non-primary PCI capable hospital
Correlation between QTc and QTcD with all caused in-hospital mortality after thrombolytic therapy in Dr. Moewardi Hospital, Surakarta
Functional capacity improvement after newly-modified cardiac rehabilitation programme in post-revascularized coronary artery disease patients
Acute heart failure and diabetes are predictors for mortality in 6 months follow up after acute coronary syndrome
Respiratory training as adjuvant of phase 2 exercise program decreased systolic pulmonary artery pressure in post mitral valve surgery patients with residual pulmonary hypertension
Correlation between Tricuspid Regurgitation Duration with Pulmonary Pressure in Heart Failure Patients
Compatibility of myocardial damage observed by echocardiography to MSCT coronary assessment of arterial stenosis in acute coronary syndrome
Hypolipidemic effect and antioxidant activity of (Tamarind) leaves in hypercholesterolemic-fed rats
Correlation

between pulmonary hypertension and functional tricuspid regurgitation in Cimacan General Hospital

Correlation knowledge and attitude regarding behaviour of controlling blood pressure in elderly hypertension patient in Manggar District

Correlation of lipid profile and high sensitivity C-reactive protein among young people

Increased in-hospital mortality associated with hyponatremia in acute heart failure patients

The association between different criteria of obesity to the severity of coronary artery lesion

Predictor postoperative arrhythmias after pediatric cardiac surgery using aortic cross clamp and cardiopulmonary bypass

Profile of hyperthyroid heart disease patients in National Cardiac Center Harapan Kita : A descriptive study

Effect of cardiac rehabilitation programs on functional capacity of post coronary revascularization patient with decrease ejection fraction using Six Minute Walk-Test

Association of postoperative hyperglycemia with in-hospital mortality in patients undergoing on pump coronary artery bypass grafting

A comparison successful reperfusion with and without fibrinolytic therapy in acute ST-segment elevation myocardial infarct patients

Factors influencing ventricular function in repaired (mid-term evaluation)

Correlation between tricuspid

regurgitation duration with right ventricular dysfunction in atrial septal defect patients
The effect of therapy Spiritual Emotional Freedom Technique (SEFT) changes on blood pressure in elderly hypertension experience in Rejosari Karanganyar Trenggalek 2015
Hyponatremia as a predictor of rehospitalization in patients with acute decompensated heart failure
Effect of sildenafil therapy to functional capacity in congenital heart disease patient with severe pulmonary hypertension in Sardjito Hospital
Eisenmenger syndrome is more prevalence in young female patients with uncorrected atrial septal defect: Result of a single-center registry
Leuko-glycemic index as an in-hospital prognostic marker in patients with ST-segment elevation myocardial infarction in Haji Adam Malik General Hospital
Serum uric acid levels as predictors of incident complications in acute coronary syndrome
Anticoagulant therapy in heart failure patients with sinus rhythm
Epicardial fat thickness is related to the blood pressure response abnormalities in exercise stress testing
Remote ischemic conditioning prior to primary percutaneous coronary intervention: The effect on Six Minutes Walk Test distance and serum NT-ProBNP level
Correlation between urinary protein and vascular

dysfunction before and after Delivery in preeclamptic women
Impact of high hsCRP level on adverse cardiovascular events and mortality in ST-elevation acute myocardial infarction undergoing fibrinolysis
Correlation between baseline soluble ST2 and global longitudinal strain
2D Speckle tracking echocardiography with left ventricle remodeling post acute myocardial infarction
Difference of peripheral P level between univalvular and multivalvular of rheumatic heart disease
Correlation between the severity of hyperthyroidism with increased of systolic blood pressure in hyperthyroidism patients
Correlation of red cell distribution width with the severity of coronary artery disease in patients with stable coronary artery disease
Chest pain presentation on acute coronary syndrome patients in Dustira Hospital
Result of exercise stress testing after cardiac rehabilitation program after coronary artery bypass grafting surgery
Relationship between plasma level of Endothelin-1 with major adverse cardiac event in ST-elevation myocardial infarction undergoing percutaneous coronary intervention
Controlled clinic blood pressure lowers the risk of left ventricular hypertrophy: An internship study among hypertensive Indonesians in rural

areas
Correlation of CHA DS -Vasc-HS scores with coronary artery lesions complexity based on SYNTAX score
The correlation between body mass index and diastolic indices in non-diabetic hypertensive male subjects
Risk of traditional cardiovascular risk factor for the development of erectile dysfunction in stable coronary artery disease patients
Relationship between left ventricle ejection fraction and submaximal exercise capacity in cardiovascular patients
Relationship between type 2 diabetes mellitus with poor heart rate autonomic response in post acute myocardial infarction patients
The ECG features among hypertensive patients in rural area: A descriptive study at Kubu Village, Karangasem, Bali
Chronotropic incompetence in coronary artery bypass graft and heart valve surgery patients: The role of β -blocker
Correlation between fibrinogen and low density lipoprotein (LDL) cholesterol levels with acute myocardial infarction
Cardiac rehabilitation use among patients with coronary artery disease on July–December 2015 in Dr. Hasan Sadikin General Hospital Bandung
Differences of plasma histamine level between acute coronary syndrome and stable coronary artery disease patients
High Endothelin-1 plasma level gives a tendency toward increasing rate of in-hospital major

adverse cardiovascular outcomes in ST-elevation myocardial infarction undergoing fibrinolysis
Correlation between serum p53 levels and vascular age determined by carotid intima media thickness (CIMT) in patients with intermediate cardiovascular risk factor
Clinical significance of precordial ST segment depression on admission electrocardiogram in patients with acute inferior myocardial infarction
Postprocedural high sensitivity C-reactive protein and the risk of recurrent ischemic events after percutaneous coronary intervention in patients with stable angina pectoris
The messenger ribonucleic acid expression of B-type natriuretic peptide, natriuretic peptide receptor type-A and type-C in cardiomyocytes of obese population
The correlation between serum galectin-3 level with wall motion score index in patients with ST-segment elevation myocardial infarction
Risk factors of atrial fibrillation in patients with chronic heart failure
QT interval prolongation after non-ST elevation myocardial infarction in type 2 diabetic compared with nondiabetic patients
The novel effect of polysaccharide peptides of to endothelial dysfunction and dislipidemia in high risk groups of atherosclerosis
Relation of Left Atrial Spontaneous Echocardiographic Contrast to

Neutrophil/Lymphocyte Ratio in Patient With Mitral Stenosis
Analysis of characteristic of atrial fibrillation versus not atrial fibrillation in hospitalized heart failure patients
Pulmonary hypertension features in adult atrial septal defect at RSUP Dr. Sardjito
The association between plasma tryptase with ventricular remodeling after 1 month in patients with acute coronary syndrome
The correlation of plasma histamine and tryptase with ST-elevation acute myocardial infarction in acute coronary syndrome patients
Acute kidney injury after coronary artery bypass graft surgery: Incidence and pre operative risk profile
The effect of diabetes mellitus on in-hospital mortality and major complications after isolated coronary artery bypass grafting surgery
Factor those influence return to work after CABG revascularization on coronary artery disease patients
The correlation of pulmonary vein endothelin-1 level with pulmonary vascular resistance before and after mitral valve surgery with pulmonary hypertension
Mitral valve E-Point septal separation as an independent predictor for in-hospital mortality after acute ST-elevation myocardial infarction
Effects of allopurinol on complications of post coronary artery bypass graft (CABG) surgery in coronary artery

disease's patient with left ventricular dysfunction
Maternal and fetal outcomes of pregnancy with valvular heart disease
QRS duration in ST elevation myocardial infarction patients undergoing a primary percutaneous intervention
Knowledge of resident medical officers on hypertensive emergency: A descriptive study
Hyperuricemia and its association with incident of congestive heart failure in Madurese population
Diagnostic value of Duke Treadmill Score in predicting coronary lesions severity in patients with suspected stable coronary artery diseases
Relationship of plasma glucose levels and troponin-I in patients wi...

Publication

18

www.isaarsci.ir

Internet Source

<1%

19

edoc.pub

Internet Source

<1%

20

Gamil Alrubaiee, Anisah Baharom, Hayati Kadir Shahar, Shaffe Mohd Daud, Huda Omar Basaleem. "Knowledge and practices of nurses regarding nosocomial infection control measures in private hospitals in Sana'a City, Yemen", *Safety in Health*, 2017

Publication

<1%

21

Submitted to Universiti Kebangsaan Malaysia

Student Paper

<1%

22

Jayantee Kalita, Rajesh Kumar Singh, Usha Kant Misra. "Cerebral Salt Wasting Is the Most Common Cause of Hyponatremia in Stroke", Journal of Stroke and Cerebrovascular Diseases, 2017

Publication

<1%

23

www.cureus.com

Internet Source

<1%

24

Kosiborod, Mikhail, Henrik S. Rasmussen, Philip Lavin, Wajeh Y. Qunibi, Bruce Spinowitz, David Packham, Simon D. Roger, Alex Yang, Edgar Lerma, and Bhupinder Singh. "Effect of Sodium Zirconium Cyclosilicate on Potassium Lowering for 28 Days Among Outpatients With Hyperkalemia : The HARMONIZE Randomized Clinical Trial", JAMA, 2014.

Publication

<1%

25

Siddiqui, Mahmudur Rahman, Quazi Tarikul Islam, Md Azharul Haque, Md Javed Iqbal, Ahmed Hossain, Yousuf Ur Rahman, Md Shahriar Mahbub, and Asif Alam Sazzad. "Electrolytes Status in Different Type of Acute Stroke Patients and Their Correlation with Some Common Clinical Presentation", Journal of Medicine, 2012.

<1%

26

Submitted to University of Brighton

Student Paper

<1%

27

"Association of Antihypertensive Monotherapy With Serum Sodium and Potassium Levels in Chinese Patients", American Journal of Hypertension, 01/08/2009

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On