

# Insect On Pig Carcasses As A Model For Predictor Of Death Interval In Forensic Medicine

*by* Erwin Kristanto 1

---

**Submission date:** 14-Aug-2018 04:52PM (UTC+0700)

**Submission ID:** 989882143

**File name:** A\_Model\_For\_Predictor\_Of\_Death\_Interval\_In\_Forensic\_Medicine.pdf (567.6K)

**Word count:** 5306

**Character count:** 29181

## Basic Medical Research

1

## Insects on pig carcasses as a model for predictor of death interval in forensic medicine

Sunny Wangko,<sup>1</sup> Erwin G. Krizanto,<sup>2</sup> Sonny J.R. Kalangi,<sup>1</sup> Johannes Huijbregts,<sup>3</sup> Dantje T. Sembel<sup>4</sup>

<sup>1</sup> Department of Anatomy-Histology, Faculty of Medicine, University of Sam Ratulangi, Manado, Indonesia

<sup>2</sup> Department of Forensic and Medicolegal Science, Faculty of Medicine, University of Sam Ratulangi, Manado, Indonesia

<sup>3</sup> Naturalis Biodiversity Center, Leiden, Netherlands

<sup>4</sup> Department of Pests and Plant Diseases, Faculty of Agriculture, University of Sam Ratulangi, Manado, Indonesia

## ABSTRAK

**Latar belakang:** Entomologi forensik belum dimanfaatkan sebagaimana mestinya di Indonesia. Keberadaan serangga forensik di Indonesia juga belum banyak dilaporkan. Penelitian ini bertujuan untuk mendapatkan jenis-jenis serangga pada bangkai hewan coba yang dapat dipergunakan untuk perkiraan saat kematian.

**Metode:** Empat babi domestik yang dimatikan dengan cara berbeda digunakan sebagai model. Bangkai diamati dua kali sehari (sekitar pukul 09.00 dan 16.00) selama 15 hari untuk mendapatkan tahap-tahap dekomposisi dan koleksi serangga baik imatur maupun dewasa. Serangga imatur dipelihara dan serangga dewasa diidentifikasi di Laboratorium hama dan penyakit tanaman, Universitas Sam Ratulangi, Manado. *Chrysomya megacephala* dan *C. rufifacies* diidentifikasi baik secara morfologik maupun tehnik deoxyribose-nucleic acid (DNA).

**Hasil:** Lima tahap dekomposisi didapatkan dalam penelitian (segar, pembusukan awal, pembusukan aktif, pembusukan lanjut, dan skeletonisasi). Serangga yang ditemukan ialah 11 spesies Diptera and delapan Coleoptera selama 15 hari sukses. *Chrysomya megacephala*, *C. rufifacies* dan *Hermetia illucens* berkolonisasi pada semua bangkai.

**Kesimpulan:** Pada keempat ekor babi serangga yang ditemukan terutama terdiri dari Diptera dan Coleoptera. *Chrysomya megacephala*, *C. rufifacies* dan *Hermetia illucens* tampaknya merupakan kandidat utama untuk perkiraan saat kematian.

## ABSTRACT

1

**Background:** Forensic entomology has not been acknowledged in Indonesia so far. Indonesian carrion insects are very rarely reported. The aim of this study was to obtain the types of insects on pig carcasses that could be used for the estimation of post-mortem interval.

**Methods:** Four domestic pigs sacrificed with different methods were used as a model. The carcasses were observed twice daily (around 9 a.m and 4 p.m) during 15 days to assess the stages of decomposition and to collect insects, both in mature and immature stages. The immature insects were reared and the mature insects were indentified in the Laboratory of Pests and Plant Diseases, University of Sam Ratulangi, Manado. *Chrysomya megacephala* and *C. rufifacies* were identified both morphologically and with deoxyribose-nucleic acid (DNA) techniques.

**Results:** Five stages of decomposition (fresh, bloated, active decay, post-decay, and skeletonization) were observed. A total of 11 Diptera and 8 Coleoptera species were found during a 15-days succession study. *Chrysomya megacephala*, *C. rufifacies* and *Hermetia illucens* colonized in all carcasses.

**Conclusion:** Insects found on four different pig carcasses consisted mainly of widespread Diptera and Coleoptera. *Chrysomya megacephala*, *C. rufifacies* and *Hermetia illucens* seemed to be primary candidates for the estimation of the post-mortem interval.

**Keywords:** *Chrysomya*, forensic entomology, *Hermetia*

2

pISSN: 0853-1773 • eISSN: 2252-8083 • <http://dx.doi.org/10.13181/mji.v24i2.1224> • Med J Indones. 2015;24:70-8

• Received 25 Mar 2015 • Accepted 25 May 2015

Correspondence author: Sunny Wangko, [sunnywangko@yahoo.com](mailto:sunnywangko@yahoo.com)

Copyright © 2015 Authors. This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (<http://creativecommons.org/licenses/by-nc-sa/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are properly cited.

Insects have been used in forensic investigation as early as the thirteenth century in China.<sup>1</sup> The first serious description of insect succession on corpses was given by Ménézier.<sup>2</sup> Early forensic work on insects was done by Nuorteva<sup>3</sup> in Finland and Leclercq<sup>4</sup> in Belgium. After the publication of the first handbook by Smith<sup>1</sup> in 1986, forensic entomology became gradually more fashionable in normal police work. In the last three decades forensic entomology developed rapidly, especially in the areas with a temperate climate. More recently studies on this topic have been carried out in several tropical countries. As the carrion insects in most tropical countries are not very well documented, thus the findings of those insects will be a great challenge. So far, no specific forensic entomology studies have been conducted in Indonesia. Research was done on the island of Sulawesi, which is on the transition of the Oriental and Australian biogeographical regions. Sulawesi has a very peculiar fauna, rich in endemic species. During the Wallace project in 1985<sup>5</sup> carrion flies were collected by Kurahashi, et al<sup>6-9</sup> and carrion beetles by Hanski and Krikken<sup>10</sup>, and Hanski and Niemela.<sup>11</sup> These insects were studied mainly in undisturbed rainforest habitats. Most human corpses are however found in an urban or suburban environment and the carrion fauna of this habitat type was not examined in Sulawesi on a systematic way.

According to the information from the Department of Forensic and Medicolegal Prof. Dr. R. D. Kandou General Hospital Manado, the most common cases of violent deaths in Manado are blunt stroke on the occipital area and cardiac puncture. In order to test whether the cause of death influenced the insect succession, four pigs were sacrificed in different ways. The aim of the present study was to identify the insect succession on pig carcasses as a model in forensic entomology for predictor of death interval.

## METHODS

Field studies were carried out at Winangun and Batu Kota, Manado, North Sulawesi, Indonesia, between January 27<sup>th</sup> and February 15<sup>th</sup>, 2012. Winangun and Batu Kota are located at a latitude 1°26'41"N and longitude 124°50'11"E, respectively, at an altitude of 66 metres above sea level. Four domestic pigs weighing approximately

20 kg each (aged two month old) were used as a model. The 1<sup>st</sup> pig was sacrificed by using 200 g potassium cyanide per oral designated as carcass I; the 2<sup>nd</sup> pig was sacrificed by a blunt stroke on its occipital area, carcass II; the 3<sup>rd</sup> and 4<sup>th</sup> pig were sacrificed by cardiac punctures, carcass III and IV. After death, three pigs were placed in protective cages at a distance of 20 m from each other at Winangun, and the other one (carcass IV) at Batu Kota. The cages consisted of a wooden frame (1.6x1x1 m<sup>3</sup>) were covered with wire meshing (diameter ± 1.7 cm).

The carcasses were observed twice daily (around 9 a.m and 4 p.m) during 15 days to assess the stages of decomposition (fresh, bloated, active decay, post-decay or skeletonization) and to collect insects.

Both mature and immature insects were collected. Insects flying around the carcasses or perched on the carcasses were collected with an insect net and recorded. Immature insects including eggs, larvae in natural cavities and wounds, and larvae from other parts of the carcasses were collected for rearing and identification in the Laboratory of Pests and Plant Diseases, Sam Ratulangi University, Manado. Soil around and underneath the carcasses were examined to observe the presence of insects including pupae. Immature insects were reared on slices of fresh pig liver. Adult insects were sacrificed with ethyl acetate and mounted on entomological pins, while immature specimens were stored in 95% alcohol.

Specific literature for the identification of adult Sulawesi carrion insects is mainly limited to blowflies. Most other species were identified by direct comparison with identified museum material. The identification of immature insects is based on rearing to adults. Voucher specimens were deposited at the Sam Ratulangi University, Manado; and Naturalis Biodiversity Center, Leiden.

*Chrysomya megacephala* and *C. rufifacies* were identified both morphologically and with deoxyribose-nucleic acid (DNA) techniques. Samples for DNA identification were extracted with AxyPrep Multisource Genomic DNA Miniprep Kit. This kit uses column purification technique that extracts both core and mitochondrial DNA.

er extraction, samples with 1270 bp cytochrome oxidase I (COI) gene were amplified with polymerase chain reaction (PCR) using primer for *Chrysomya*, as follows:<sup>12</sup>

- C1-J-1718f (5'-GGAGGATTTGGAAATTGATTAGTTCC)
- TL2-N-3014r (5' -TCCAATGCACTAATCTGCCATATTA)
- TL2-N-3014MODr (5' -TCCATTGCACTAATCTTGCCATATTA)

**Table 1.** Steps of PCR of COI gene

Step	Temperature	Time
Predenaturation	95°C	28 minutes
Denaturation	95°C	30 seconds
Primary annealing	54°C	30 seconds
Extention	72°C	90 seconds
Final extention	72°C	90 seconds

**Table 2.** Sulawesi carrion insects reported in taxonomic literature

Species	Source
<b>Coleoptera</b>	
<i>Hybosoridae</i> <i>Phaeochrous emarginatus</i> Castelnau	Hanski & Krikken, 1991 <sup>10</sup>
<i>Hybosoridae</i> <i>Phaeochrous sulawesi</i> Kuijten	Kuijten, 1978 <sup>13</sup>
<i>Scarabaeidae</i> <i>Onthophagus aper</i> Sharp	Hanski & Krikken, 1991 <sup>10</sup>
<i>Scarabaeidae</i> <i>Onthophagus mentaveiensis</i> Boucomont	Hanski & Krikken, 1991 <sup>10</sup>
<i>Scarabaeidae</i> <i>Onthophagus scrutator</i> Harold	Hanski & Krikken, 1991 <sup>10</sup>
<i>Silphidae</i> <i>Necrophilia renatae</i> (Portevin)	Ruzicka, Schneider, Qubaiová & Nishikawa, 2012 <sup>14</sup>
<i>Silphidae</i> <i>Necrophilia charon</i> Sikes & Madge	Sikes, Madge & Trumbo, 2006 <sup>15</sup>
<i>Silphidae</i> <i>Nicrophorus distinctus</i> (Grouvelle)	Sikes, Madge & Newton, 2002 <sup>16</sup>
<b>Diptera</b>	
<i>Calliphoridae</i> <i>Calliphora hasanuddini</i> Kurahashi & Selomo	Kurahashi & Selomo, 1997 <sup>9</sup>
<i>Calliphoridae</i> <i>Chrysomya greenbergi</i> Wells & Kurahashi	Wells & Kurahashi, 1996 <sup>17</sup>
<i>Calliphoridae</i> <i>Chrysomya nigripes</i> Aubertin	Singh, Kurahashi & Wells, 2011 <sup>18</sup>
<i>Calliphoridae</i> <i>Chrysomya rufifacies</i> (Macquart)	James, 1977 <sup>19</sup>
<i>Calliphoridae</i> <i>Chrysomya yayukae</i> Kurahashi & Magpayo	Kurahashi & Magpayo, 1987 <sup>8</sup>
<i>Calliphoridae</i> <i>Hemipyrellia ligurriens</i> (Wiedemann)	James, 1977 <sup>19</sup>
<i>Calliphoridae</i> <i>Idiella divisa</i> (Walker)	James, 1977 <sup>19</sup>
<i>Calliphoridae</i> <i>Isomya delectans</i> (Walker)	James, 1977 <sup>19</sup>
<i>Calliphoridae</i> <i>Lucilia adisoemartoi</i>	Kurahashi, 1988 <sup>6</sup>
<i>Calliphoridae</i> <i>Lucilia papuensis</i> Macquart	James, 1977 <sup>19</sup>
<i>Calliphoridae</i> <i>Phumosia abdominalis</i> Robineau-Desvoidy	James, 1977 <sup>19</sup>
<i>Calliphoridae</i> <i>Phumosia elegans</i> Kurahashi	Kurahashi, 1989 <sup>7</sup>
<i>Calliphoridae</i> <i>Phumosia indica</i> (Surcouf)	James, 1977 <sup>19</sup>
<i>Calliphoridae</i> <i>Phumosia njonja</i> Kurahashi	Kurahashi 1989 <sup>7</sup>
<i>Calliphoridae</i> <i>Phumosia promittens</i> (Walker)	James, 1977 <sup>19</sup>
<i>Calliphoridae</i> <i>Strongyloneura prolata</i> (Walker)	James, 1977 <sup>19</sup>

Primers were made at PT Genetika Science Indonesia located in Jakarta, Indonesia. PCR was carried out with Biometra® T Personal, with temperatures of each cycle as seen as Table 1.

PCR products were visualized using 1.0% agarose gel, dyed in ethidium bromide 1.5% and transilluminated with ultraviolet (UV) rays, and were then sequenced with COI gene bar coding that is commonly used in flies identification.

## RESULTS

The mean daily temperature at the study sites fluctuated between 23.4°C and 27.2°C. Almost all days had heavy rainfalls.

No comprehensive review of Indonesian carrion insects did exist. Many tropical carrion insects were described a long time ago without mentioning

that they were collected on carrions. More recent taxonomical papers usually contain more ecological information but it is very difficult to trace specific carrion insects in scattered publications. Based on the carrion-taxa found in neighbouring countries, literature was searched for potential carrion insects in Sulawesi. The results of the literature search are summarized in Table 2.

6 Five stages of decomposition were distinguished in all carcasses. The duration of the 27 different stages is illustrated in Table 3. The fresh stage was observed on day one. The bloated stage was observed on days two and three for all carcasses, but facial destruction was first observed on carcass II. The active decay stage was observed on days four and five in carcass II, III and IV meanwhile in carcass I on days four to six. The post-decay stage was observed on days six to eight in carcass II and III, and just on day six in carcass IV. In carcass I, however, this stage lasted until day 10. Skeletonization was observed on day nine for carcasses II and III, on day seven for carcass IV, but on day 11 for carcass I. The different insects in relation to the decomposition stages are summarized in Table 4.

26 **fresh stage**

*C. megacephala* and *C. rufifacies* visited the carcasses at the fresh stage from day one. Eggs

of *Chrysomya* were found on day one in natural orifices (mouth, nose, ear, and anus) and the stab wounds. Other *Diptera* which visited the carcasses on the same day were *Ophyra spinigera* and *Sarcophagidae*. *Formicidae* were found from the first hour of post-mortem in the natural orifices, especially in those with fresh blood.


**Bloated stage**

Blowfly larvae invaded the carcasses' faces voraciously especially in carcasses II, III and IV. *C. rufifacies* and *C. megacephala* were abundant. *Musca domestica*, *Ophyra spinigera*, and *Sarcophagidae* were still found at this stage. *Sarcophagidae* were observed only on intact skin, and not on the natural orifices or wounds. Beetles found at this stage were the predators *Saprinus* and *Philonthus* on day three. *Riptortus linearis* was found on intact abdominal skin on day three. On day two, *Formicidae* left the natural orifices invaded by *Chrysomya* maggots, but were still found in the non-invaded areas.

Act 17 **decay stage**

At the end of this stage, *Diptera* larvae had consumed most part of the body tissues leaving the skin, cartilages, bones, and the last part of muscles and guts. This stage lasted for two days in carcasses II, III and IV, and three days in carcass I.

**Table 3.** Duration of decomposition stages in four different pig carcasses in Manado



**Table 4.** Insects found on four different pig carcasses in Manado in relation <sup>25</sup> to the decomposition stages

24	Species	Fresh	Bloated	Active decay	Post-decay	Skeletonization
<b>Coleoptera</b>						
<i>Cleridae</i>	<i>Necrobia rufipes</i> (De Geer)	-	-	-	A	A
<i>Dermestidae</i>	<i>Dermestes maculatus</i> De Geer	-	-	-	A	A
<i>Histeridae</i>	<i>Pachylister tabellio</i> Marseul	-	-	-	A	-
<i>Histeridae</i>	<i>Saprinus</i> sp.	-	A	A	A	A
<i>Hydrophilidae</i>	<i>Sphaeridium discolor</i> Orchymont	-	-	A	4	4
<i>Silphidae</i>	<i>Diamesus osculans</i> Vigors	-	-	A	A, L	A, L
<i>Staphylinidae</i>	<i>Creophilus flavipennis</i> Hope	-	-	A	A	A
<i>Staphylinidae</i>	<i>Philonthus</i> sp.	-	A	A	A, L	A, L
<b>Dermaptera</b>						
<i>Labiidae</i>		-	-	-	A	-
<b>Diptera</b>						
<i>Calliphoridae</i>	<sup>23</sup> <i>Chrysomya megacephala</i> (Fabricius)	A	A, L	A, L	A, L	A
<i>Calliphoridae</i>	<i>Chrysomya rufifacies</i> (Macquart)	A	A, L	A, L	A, L	A
<i>Calliphoridae</i>	<i>Hemipyrellia ligurriensis</i> (Wiedemann)	-	A	-	-	-
<i>Fanniidae</i>	<i>Fannia</i> sp.	-	-	-	A	-
<i>Muscidae</i>	<i>Atherigona</i> sp.	-	-	A	A	-
<i>Muscidae</i>	<i>Musca domestica</i> L.	-	A	A, L	L	L
<i>Muscidae</i>	<i>Ophyra spinigera</i> Stein	A	A	A	A, L	A
<i>Phoridae</i>	<i>Phoridae</i> sp.	-	-	-	A	-
<i>Piophilidae</i>	<i>Protophiophila</i> sp.	-	-	A	A	A
<i>Sarcophagidae</i>	<i>Sarcophaga</i> s.l. sp.	A	-	-	-	A
<i>Stratiomyidae</i>	<i>Hermetia illucens</i> (L.)	-	-	A	A	-
<b>Heteroptera</b>						
<i>Alydidae</i>	<i>Riptortus linearis</i> (Fabricius)	-	A	A	A	-
<b>Hymenoptera</b>						
<i>Formicidae</i>		A	A	A	A	A

Notes: A = adults; L = larvae

*C. rufifacies* and *C. megacephala* both larvae and adults were abundant. *Musca domestica*, *Ophyra spinigera*, and *Sarcophagidae* still visited the carcasses. Other *Diptera* such as *Protophiophila*, *Phoridae*, and *H. illucens* were found at the end of the active decay stage (day five). Beetles in this stage were *Saprinus*, *Philonthus* and *Sphaeridium discolor*. There were many Formicidae, predating on *Chrysomya* larvae which left the carcasses for pupation. *R. linearis* were still present.

#### Post-decay stage

During this stage, especially *Dermestes* consumed the remaining tissue. Only very few adult *C. rufifacies* and *C. megacephala* were observed. Other *Diptera* (*Protophiophila*, *Phoridae* and *H. illucens*) and *R. linearis* were still found in the early post-decay stage (day six). The beetles of the active decay stage were still present in this

stage as well as a few Formicidae. On day eight, one adult *Labiidae* was found in carcass II.

#### Skeletonization

In this stage, adult *C. rufifacies* and *C. megacephala* of the new generation emerged from their pupae on days 12 and 13. However, their larvae and *Muscidae* larvae were still found under and around the carcasses. *Dermestes maculatus*, *Saprinus*, *Philonthus*, and *Necrobia rufipes* were still present.

## DISCUSSION

#### Decomposition

Generally, five stages of decomposition <sup>15</sup> are recognized. <sup>20-23</sup> Payne<sup>24</sup> reported six stages (fresh, bloating, active decay, advanced decay, dry and

remains) in Clemson, South Carolina. Archer and Elgar<sup>25</sup> reported only two stages (initial decomposition and advanced decomposition) from Victoria, Australia while Wang, et al<sup>26</sup> in South China distinguished four stages.

**6**  
In our study we observed five stages of decomposition in all carcasses; the duration of the decomposition stages in the different carcasses was however not the same. Carcass I took 10 days for skeletonization, carcasses II and III only took eight days, and carcass IV six days. Carcass I had no wounds or external blood in contrast to carcass II (fresh blood that came out of the mouth), III and IV (stab wound); oviposition by blowflies on carcass I was therefore much later. The duration of decomposition (7-10 days) resembled to those reported by Wang, et al<sup>26</sup> in South China and Heo, et al<sup>27</sup> at Tanjung Sepat, peninsular Malaysia meanwhile Apichat, et al<sup>28</sup> in Phitsanulok, Thailand reported one day each for fresh, bloated, and active decay stages, but three days for post-decay stage and a much longer skeletonization period (days 8-31).

**9**  
Generally, *Calliphoridae* are the first insects to lay eggs on a corpse or carcass which makes them the most important tool of the forensic entomologist.<sup>1,29</sup> In our study, *Chrysomya megacephala* and *C. rufifacies* were abundantly found in both adult and immature forms. *C. rufifacies* maggots are predators on *C. megacephala* maggots and their eggs are therefore deposited later than those of *C. megacephala*. These dominant *Chrysomya* species are the same as those from similar experiments in other parts of tropical Asia as reported by Heo, et al<sup>27</sup> in Malaysia, Apichat, et al<sup>28</sup> in Thailand and Wang, et al<sup>26</sup> in South China.

Rearing of larvae and pupae in the laboratory resulted in adult *C. rufifacies* and *C. megacephala*. From both larva and pupa samples of carcasses II and III, the adults emerged one day earlier than those of carcass I. Adult *Chrysomya* at the study site emerged one day later than those in the laboratory. This faster development was caused by the higher temperature in the laboratory (28°-30°C).

Heo, et al<sup>27</sup> reported that the second generation blowflies appeared on day-12 and -13 around the vegetation on **12** study sites, which are similar to our results. Larvae of *C. megacephala* and *C. rufifacies* were still present underneath and around

the carcasses until skeletonization. According to Gunn<sup>30</sup> adult flies also feed on the carcass during the post-decay stage or skeletonization, but they did not initiate oviposition.

*Hermetia illucens* was originally restricted to the New World but it has spread to other regions, both tropical and temperate.<sup>29,31</sup> This species is not often found in corpses, and is reported to colonize carrion in an advanced stage: 20-30 days post-mortem<sup>21,30,32</sup> In our study, adult *Hermetia illucens* were already found on day five. Diclaro and Kauffman<sup>33</sup> reported that *H. illucens* larvae have six instars which needed 14 days, and a pupation period of two weeks. The rearing of soil around the carcasses collected on day 11, resulted in a second generation of *Hermetia illucens* on days 35-39. The growth of the larvae was in accordance with the findings of Tomberlin, et al.<sup>34</sup>

Sarcophagidae larvae are reported from large carrion including forensic cases.<sup>29,35</sup> In our study, adult Sarcophagidae were found in small numbers from day 1 until the end of the study, but no maggots or pupae were observed. As these species apparently did not develop in the carcasses they cannot be used as forensic indicators. The adult *Sarcophagidae* were not identified to species level.

Larvae of *Ophyra* (instar two and three) are predators of larvae *Musca domestica* and other Muscidae larvae.<sup>1,29,36</sup> In our study, adult *Ophyra spinigera* was found from day 1-8, and their larvae on day 6-14 together with other *Muscidae* larvae.

Adults of *Musca domestica* usually visit a corpse after the arrival of blowflies and flesh flies in both the active and post-decay stages. House flies are rarely attracted to fresh corpses, unless there is excreta or exposed gut present.<sup>1,29,30</sup> In our study, *Musca domestica* was found from day 2-5, earlier than that observed by Byrd and Castner<sup>29</sup>, Gunn<sup>30</sup>, and Smith<sup>1</sup>. The first adult fly emerged during rearing in the laboratory on day seven which is similar to that of Gunn<sup>30</sup> who stated that the life cycle of *Musca domestica* takes six to eight days at a relatively high temperature.

In our study, adults of *Phoridae* were found on day-5. Rearing in the laboratory of soil collected at day eight resulted in pupae and adult *Phoridae*

which shows that development of the Phoridae occurred on the carcasses. Unfortunately the identification of *Phoridae* is notoriously difficult which prevents their usage as a forensic indicator at the moment.

In general, adults of *Protopiophila* visit carcasses in an active state of decay and an early dry stage. Larvae, as well as the adults, are necrophagous.<sup>29</sup> In our study, adult *Protopiophila* were found on day 4-6 but their larvae were not found on the carcasses or during rearing of soil samples. Adult *Protopiophila* were not identified to species level.

The beetles found in our experiment did feed on maggots or on parts of the carcass that were not consumed by maggots and therefore, they arrived usually later than the blowflies. The earliest Coleoptera visiting the carcasses were *Staphylinidae* (day-3), followed by *Saprinus* sp. (day-3, -5, -6, -10); these taxa are predators<sup>29</sup> and their presence was in accordance with the abundance of *Diptera* larvae on the carcasses. *Sphaeridium discolor* was found on day-4 and -5. Most *Sphaeridium* species are dung specialists and they were probably attracted by the gut contents of the carcasses. *Dermestes maculatus* (day-6, -10) and *Necrobia rufipes* (day-10) were observed later in the succession. *Dermestes* species are usually found on dry carcasses in the wild. Almost all species are scavengers and feed on various types of dried animal tissue.<sup>29</sup> *Necrobia* sp. have a special preference for fatty tissues: their larvae are sometimes predacious.<sup>37</sup> The *Coleoptera* larvae collected are belong to *Diamesus osculans* and *Staphylinidae*.

*Formicidae* are social insects which are distributed all over the world. In tropical areas they are very abundant and can be found in all stages of decomposition as predators or omnivores feeding on both body tissue and exudate fluids.<sup>1,30</sup> The ants were observed in the first hour after the start of the experiment and during the whole decomposition process in accordance with the statements of Aggarwal<sup>38</sup> and Heo, et al<sup>39</sup>. Ants were usually found in the natural orifices, they will leave these areas after being invaded by the *Chrysomya*.

Adults of *Riptortus linearis* were found on day 2-6. These bugs landed on the intact abdominal skin of carcasses, and not on the natural orifices

or the open wounds, which suggest they are accidental visitors. All *Alydidae*, the family to which *Riptortus* belongs, are apparently plant-feeders<sup>1,40</sup> but Payne<sup>41</sup> reported three species of *Alydidae* feeding on a pig carcass in the USA. The relation between *Coreidae* and carcasses seems doubtful at the moment.

*Labiidae* are widely distributed, especially in hot climates.<sup>42</sup> Several species are predators on other insects by seizing them with their forceps or are omnivorous.<sup>29,40,42</sup> On day eight one adult *Labiidae* was found on carcass II. At this moment we consider this condition as an accidental visitor.

<sup>1</sup> Literature study revealed a total of 11 *Diptera* and eight *Coleoptera* species which were belong to the potential carrion fauna of Sulawesi. Most of these taxa were different from those of the literature study, probably because the literature data were mainly based on the forest environments. The suburban Manado list contains mainly widespread taxa while the species from the literature list mainly endemic species in Sulawesi.

<sup>1</sup> Five stages of decomposition (fresh, bloated, active decay, post-decay and skeletonization) could be distinguished during the 15 days study. *Chrysomya megacephala* and *C. rufifacies* dominated the other insects in all carcasses. *Formicidae* were the first visitors and found during the whole study, but abandoned areas which were invaded by *Chrysomya* larvae. Although the insect succession patterns were generally similar in all carcasses, the duration of decomposition and the moment on which the second generation of *Chrysomya* adults appeared were not the same. Traumatic carcasses showed a shorter decomposition period and an earlier emergence of *Chrysomya* adults. Further experiments are needed to prove that these succession patterns are reliable enough to be used in court.

The morphological identification of *C. megacephala* and *C. rufifacies* was confirmed with DNA-techniques. The carrion fauna of a limited part of Sulawesi is certainly not representative for all regions in Indonesia. Further research in other regions will reveal much more species of the <sup>20</sup> non-community and thus, can lay a fundamental for the development of forensic entomology in Indonesia.



In conclusion, during our succession study on pig carcasses in suburban Manado, 11 Diptera and 8 Coleoptera taxa were collected. Six of the collected taxa were collected as larvae on the field location and two additional taxa were reared to adults from soil samples collected at the field location. The fact that these species developed on the carcasses classify them as potential forensic indicators. From *Chrysomya megacephala*, *C. rufifacies* and *Hermetia illucens*, the development from larvae to adults in relation to the ambient temperature is already known and they seem therefore to be primary candidates for the estimation of the post-mortem interval in suburban Sulawesi.

#### Acknowledgment

We would like to thank the International Affairs Office of University of Sam Ratulangi Manado that facilitated Drs. Johannes Huijbregts from Naturalis Biodiversity Center, Leiden, Netherlands, to come to Manado, Indonesia.

#### Conflict of interest

The authors affirm no conflict of interest in this study.

#### REFERENCES

1. Smith KGV. A manual of forensic entomology. London: The Trustees of the British Museum (Natural History); 1986.
2. Mégnin JP. La faune des cadavres: application de l'entomologie à la médecine légale, Encyclopédie Scientifique des Aide-Mémoire 101. Paris: Masson et Gauthiers-Villars; 1894. France.
3. Nuorteva P. Sarcosaprophagous insects as forensic indicators. In: Tedeschi CG, Eckert WG, Tedeschi LG, editors. Forensic medicine, a study in trauma and environmental hazards. Philadelphia: Saunders; 1977. p. 1072-95.
4. Leclercq M. Entomologie et médecine légale, datation de la mort. Collection de Médecine Légale et de Toxicologie Médicale 108. Paris: Masson; 1978. France.
5. Knight WJ, Holloway JD. Insects and the rain forests of South East Asia (Wallacea). London: Royal Entomological Society of London; 1990.
6. Kurahashi H. A new species of *Lucilia* (Diptera, Calliphoridae) from Sulawesi. Kontyu. 1988;56:144-7.
7. Kurahashi H. The genus *Phumosia* of Sulawesi, Indonesia, with descriptions of two new species (Diptera: Calliphoridae). Jpn J Sanit Zool. 1989;40(3):203-10.
8. Kurahashi H, Magpayo FR. Two new species of the genus *Chrysomya* from Wallacea (Diptera, Calliphoridae). Kontyu. 1987;55:71-9.
9. Kurahashi H, Selomo M. A new species of *Calliphora* from Sulawesi, Indonesia (Diptera, Calliphoridae). Jpn J Syst Ent. 1997;3:123-7.
10. Hanski I, Krikken J. Dung beetles in tropical forests in South-East Asia. In: Hanski I, Cambefort Y, editors. Dung beetle ecology. Princeton: Princeton University Press; 1991. p. 179-97.
11. Hanski I, Niemela J. Elevational distributions of dung and carrion beetles in northern Sulawesi. In: Knight WJ, Holloway JD, editors. Insects and the rain forests of South East Asia (Wallacea). London: Royal Entomological Society of London; 1990. p. 145-52.
12. Harvey ML, Gaudieri S, Villet MH, Dadour IR. A global study of forensically significant calliphorids: implications for identification. Forensic Sci Int. 2008;177(1):66-76.
13. Kuijten PJ. Revision of the Indo-Australian species of the genus *Phaeochrous* Castelnau, 1840 (Coleoptera: Scarabaeidae, Hybosorinae), with notes on the African species. Zool Verh. 1978;165:3-42.
14. Růžička J, Schneider J, Qubaiova J, Nishikawa M. Revision of palaeartic and oriental necrophila Kirby & Spence, part 2: subgenus *Chrysosilpha* Portevin (Coleoptera: Silphidae). Zootaxa. 2012;3261:33-58.
15. Sikes DS, Madge RB, Trumbo ST. Revision of *Nicrophorus* in part: new species and inferred phylogeny of the nepalensis-group based on evidence from morphology and mitochondrial DNA (Coleoptera: Silphidae: Nicrophorinae). Invertebr Syst. 2006;20(3):305-65.
16. Sikes DS, Madge RB, Newton AF. A catalog of the Nicrophorinae (Coleoptera: Silphidae) of the world. Zootaxa. 2002;65:1-304.
17. Wells JD, Kurahashi H. A new species of *Chrysomya* (Diptera: Calliphoridae) from Sulawesi, Indonesia, with a key to the Oriental, Australasian and Oceanian species. Med Entomol Zool. 1996;47(2):131-8.
18. Singh B, Kurahashi H, Wells JD. Molecular phylogeny of the blowfly genus *Chrysomya*. Med Vet Entomol. 2011;25(2):126-34.
19. James MT. Family Calliphoridae. In: Delfinado MD, Hardy DE, editors. A catalog of the Diptera of the oriental region. Honolulu: University Press; 1977. p. 526-56.
20. Goff ML. Early postmortem changes and stages of decomposition. In: Amendt J, Campobasso CP, Goff ML, Grassberger M, editors. Current concepts in forensic entomology. Dordrecht: Springer; 2010. p. 1-24.
21. Goff ML. Forensic entomology. In: Mozayani A, Noziglia C, editors. The forensic laboratory handbook procedures and practice. New York: Humana Press; 2011. p. 448-54.
22. Kreitlow KLT. Insect succession in a natural environment. In: Byrd JH, Castner JL, editors. Forensic entomology: The utility of arthropods in legal investigations. Boca Raton: CRC Press; 2010. p. 251-6.
23. Sharanowski BJ, Walker EG, Anderson GS. Insect succession and decomposition patterns on shaded and sunlit carrion in Saskatchewan in three different seasons. Forensic Sci Int. 2008;179(2-3):219-40.
24. Payne JA. A summer carrion study of the baby pig *Sus scrofa* Linnaeus. Ecology. 1965;46(5):592-602.
25. Archer MS, Elgar MA. Effects of decomposition on carcass attendance in a guild of carrion-breeding flies. Med Vet Entomol. 2003;17(3):263-71.
26. Wang J, Li Z, Chen Y, Chen Q, Yin X. The succession and development of insects on pig carcasses and their significances in estimating PMI in south China. Forensic Sci Int. 2008;179(1):11-8.

27. Heo CC, Mohamad AM, Ahmad FM, Jeffrey J, Kurahashi H, Omar B. Study of insect succession and rate of decomposition on a partially burned pig carcass in an oil palm plantation in Malaysia. *Trop Biomed.* 2008;25(3):202-8.
28. Apichat V, Wilawan P, Udomsak T, Chanasorn P, Saengchai N. A preliminary study on insects associated with pig (*Sus scrofa*) carcasses in Phitsanulok, northern Thailand. *Trop Biomed.* 2007;24(2):1-5.
29. Byrd JH, Castner JL, editors. *Forensic Entomology: The utility of arthropods in legal investigations.* 2nd ed. Boca Raton: CRC Press; 2010.
30. Gunn A. *Essential forensic biology.* 2nd ed. Chichester (UK): Wiley-Blackwell; 2009.
31. Tomberlin JK, Shepard DC. Lekking behavior of the black soldier fly (Diptera: Stratiomyidae). *Fla Entomol.* 2001;84(4):729-30.
32. Lord WD, Goff ML, Adkins TR, Haskell NH. The black soldier fly *Hermetia illucens* (Diptera: Stratiomyidae) as a potential measure of human postmortem interval: observations and case histories. *J Forensic Sci.* 1994;39(1):215-22.
33. Diclaro JW, Kaufman PE. Black soldier fly *Hermetia illucens* Linnaeus (Insecta: Diptera: Stratiomyidae). *EENY.* 2009;461:1-3.
34. Tomberlin JK, Sheppard DG, Joyce JA. Selected life-history traits of black soldier flies (Diptera: Stratiomyidae) reared on three artificial diets *Ann Entomol Soc Am.* 2002;95(3):379-86.
35. Villet MH. African carrion ecosystems and their insect communities in relation to forensic entomology. *Pest Technol.* 2011;5(1):1-15.
36. Vibe-Petersen S. Development, survival and fecundity of the urine fly *Scatella (Teichomyza) fusca* and predation by the black dumpfly, *Hydrotaea aenescens*. *Entomol Exp Appl.* 1998;87(2):157-69.
37. Busvine JR. *Insects and Hygiene. The biology and control of insect pests of medical and domestic importance.* 2nd ed. London: Methuen; 1966.
38. Aggarwal AD. *Estimating the post-mortem interval with the help of entomological evidence [Thesis].* Medical College, Patiala. Baba Farid University of Health Sciences Faridkot, India; 2005.
39. Heo CC, Mohamad AR, Rosli H, Nurul Ashikin A, Chen CD, John J, et al. Ants (Hymenoptera: Formicidae) associated with pig carcasses in Malaysia. *Trop Biomed.* 2009;26(1):106-9.
40. Kalshoven LGE. *Pests of crops in Indonesia.* Jakarta: Ichtar Baru; 1981.
41. Payne JA, Mead FW, King EW. Hemiptera associated with pig carrion. *Ann Entomol Soc Am.* 1968;61:565-7.
42. Gillott C. *Entomology.* 3rd ed. Dordrecht: Springer; 2005.

# Insect On Pig Carcasses As A Model For Predictor Of Death Interval In Forensic Medicine

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

11%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1

[id.portalgaruda.org](http://id.portalgaruda.org)

Internet Source

8%

2

Ulfah Lutfiah, Sri W.A. Jusman, Mochammad Sadikin. "H,K-ATPase and carbonic anhydrase response to chronic systemic rat gastric hypoxia", Medical Journal of Indonesia, 2015

Publication

2%

3

Silvia W. Lestari, Dwi A. Pujianto, Purnomo Soeharso, Evelyn Loanda. "Evaluation of outer dense fiber-1 and -2 protein expression in asthenozoospermic infertile men", Medical Journal of Indonesia, 2015

Publication

1%

4

Zaidi, F.. "A preliminary survey of carrion breeding insects associated with the Eid ul Azha festival in remote Pakistan", Forensic Science International, 20110615

Publication

1%

5

[uad.portalgaruda.org](http://uad.portalgaruda.org)

Internet Source

1%

6

[researchdirect.uws.edu.au](http://researchdirect.uws.edu.au)

Internet Source

<1%

7

A. Yasmin, A. Aryadi, A. Widya, A. P. Suwirya et al. " Original ResearchHypertension in rural area: The determinants of left ventricular geometric alteration among hypertensive patients at Kubu Village, Karangasem, BaliEffectiveness comparison between lisinopril and amlodipine in Melanesian patients with hypertensionAn experimental study of  $\beta$ -D-Glucans extract of polysaccharide peptides as anti-inflammation and anti-oxidant in patients at high-risk of atherosclerosisEchocardiographic findings among hypertensive patients in rural area: The descriptive study at Kubu Village, Karangasem, BaliAddition of left ventricular ejection fraction to the GRACE score in prediction of short term prognosis in patients with non ST-elevation myocardial infarctionEffect of colchicine on HsCRP and mean platelet volume in acute myocardial infarctionCorrelation between GRACE score with spatial QRS-T angle in non-ST elevation myocardial infarction patientsPrevalence of Impaired Kidney

<1%

Function in Hospitalized Hypertensive Patients  
in Agoesdjam General Hospital, Ketapang,  
West Borneo, Indonesia

Inappropriate initial  
treatment in acute coronary syndrome patients:  
A gap in cardiovascular care

Correlation  
between plasma histamine level and  
atherosclerosis severity based on carotid intima  
media thickness (CIMT) measurement in  
patients with stable coronary artery  
disease

Clinical characteristics, management  
and outcome of patients with ST elevation  
myocardial infarction in Pekanbaru city:  
Preliminary result of single center  
experience

Phase II cardiac rehabilitation  
program increases functional capacity in post  
atrial septal defect surgical closure in adult  
patients

The identification of human telomerase  
catalytic subunit in cervical cancer patients and  
its potential usage as strategy to reawaken  
hibernating myocardium post myocardial  
infarction

Could EURO heart risk score predict  
the severity of coronary lesion?

Pericardial  
effusion in systemic lupus  
erythematosus

Intracardiac metastasis of  
hepatocellular carcinoma

Myocardial bridging  
assessed by computed tomography scan: A  
cause of chest pain in patients without  
obstructed coronary arteries

Correlation  
between resting heart rate and 6-Minute Walk

Test distance in systolic heart failure patients  
Severe pulmonary hypertension is a predictor of mortality in pregnant unoperated atrial septal defect  
High plasma level of soluble ST2 associated with adverse outcomes during acute phase of ST-elevation myocardial infarction  
Effect of extracorporeal shockwave myocardial revascularization on spatial QRS – T angle resolution in stable coronary artery disease patients  
Fragmented QRS study, does it have any structural abnormality relation?  
Risk of coronary artery lesions severity to erectile dysfunction incidence in stable coronary heart disease patients  
Clinical profiles and outcomes of patients with ST-Elevation Acute Coronary Syndrome (STE-ACS) in non-primary PCI capable hospital  
Correlation between QTc and QTcD with all caused in-hospital mortality after thrombolytic therapy in Dr. Moewardi Hospital, Surakarta  
Functional capacity improvement after newly-modified cardiac rehabilitation programme in post-revascularized coronary artery disease patients  
Acute heart failure and diabetes are predictors for mortality in 6 months follow up after acute coronary syndrome  
Respiratory training as adjuvant of phase 2 exercise program decreased systolic pulmonary artery pressure in post mitral valve surgery patients with residual pulmonary

hypertension  
Correlation between Tricuspid Regurgitation Duration with Pulmonary Pressure in Heart Failure Patients  
Compatibility of myocardial damage observed by echocardiography to MSCT coronary assessment of arterial stenosis in acute coronary syndrome  
Hypolipidemic effect and antioxidant activity of (Tamarind) leaves in hypercholesterolemic-fed rats  
Correlation between pulmonary hypertension and functional tricuspid regurgitation in Cimacan General Hospital  
Correlation knowledge and attitude regarding behaviour of controlling blood pressure in elderly hypertension patient in Manggar District  
Correlation of lipid profile and high sensitivity C-reactive protein among young people  
Increased in-hospital mortality associated with hyponatremia in acute heart failure patients  
The association between different criteria of obesity to the severity of coronary artery lesion  
Predictor postoperative arrhythmias after pediatric cardiac surgery using aortic cross clamp and cardiopulmonary bypass  
Profile of hyperthyroid heart disease patients in National Cardiac Center Harapan Kita : A descriptive study  
Effect of cardiac rehabilitation programs on functional capacity of post coronary revascularization patient with decrease ejection fraction using Six Minute

Walk-Test Association of postoperative hyperglycemia with in-hospital mortality in patients undergoing on pump coronary artery bypass grafting A comparison successful reperfusion with and without fibrinolytic therapy in acute ST-segment elevation myocardial infarct patients Factors influencing ventricular function in repaired (mid-term evaluation) Correlation between tricuspid regurgitation duration with right ventricular dysfunction in atrial septal defect patients The effect of therapy Spiritual Emotional Freedom Technique (SEFT) changes on blood pressure in elderly hypertension experience in Rejosari Karangas Trenggalek 2015 Hyponatremia as a predictor of rehospitalization in patients with acute decompensated heart failure Effect of sildenafil therapy to functional capacity in congenital heart disease patient with severe pulmonary hypertension in Sardjito Hospital Eisenmenger syndrome is more prevalence in young female patients with uncorrected atrial septal defect: Result of a single-center registry Leuko-glycemic index as an in-hospital prognostic marker in patients with ST-segment elevation myocardial infarction in Haji Adam Malik General Hospital Serum uric acid levels as predictors of incident complications in acute coronary



syndrome  
Anticoagulant therapy in heart failure patients with sinus rhythm  
Epicardial fat thickness is related to the blood pressure response abnormalities in exercise stress testing  
Remote ischemic conditioning prior to primary percutaneous coronary intervention: The effect on Six Minutes Walk Test distance and serum NT-ProBNP level  
Correlation between urinary protein and vascular dysfunction before and after Delivery in preeclamptic women  
Impact of high hsCRP level on adverse cardiovascular events and mortality in ST-elevation acute myocardial infarction undergoing fibrinolysis  
Correlation between baseline soluble ST2 and global longitudinal strain  
2D Speckle tracking echocardiography with left ventricle remodeling post acute myocardial infarction  
Difference of peripheral P level between univalvular and multivalvular of rheumatic heart disease  
Correlation between the severity of hyperthyroidism with increased of systolic blood pressure in hyperthyroidism patients  
Correlation of red cell distribution width with the severity of coronary artery disease in patients with stable coronary artery disease  
Chest pain presentation on acute coronary syndrome patients in Dustira Hospital  
Result of exercise stress testing after

cardiac rehabilitation program after coronary artery bypass grafting surgery  
Relationship between plasma level of Endothelin-1 with major adverse cardiac event in ST-elevation myocardial infarction undergoing percutaneous coronary intervention  
Controlled clinic blood pressure lowers the risk of left ventricular hypertrophy: An internship study among hypertensive Indonesians in rural areas  
Correlation of CHA<sub>2</sub>DS<sub>2</sub>-Vasc-HS scores with coronary artery lesions complexity based on SYNTAX score  
The correlation between body mass index and diastolic indices in non-diabetic hypertensive male subjects  
Risk of traditional cardiovascular risk factor for the development of erectile dysfunction in stable coronary artery disease patients  
Relationship between left ventricle ejection fraction and submaximal exercise capacity in cardiovascular patients  
Relationship between type 2 diabetes mellitus with poor heart rate autonomic response in post acute myocardial infarction patients  
The ECG features among hypertensive patients in rural area: A descriptive study at Kubu Village, Karangasem, Bali  
Chronotropic incompetence in coronary artery bypass graft and heart valve surgery patients: The role of  $\beta$ -blocker  
Correlation between fibrinogen and low density lipoprotein (LDL) cholesterol levels with

acute myocardial infarction  
Cardiac rehabilitation use among patients with coronary artery disease on July–December 2015 in Dr. Hasan Sadikin General Hospital Bandung  
Differences of plasma histamine level between acute coronary syndrome and stable coronary artery disease patients  
High Endothelin-1 plasma level gives a tendency toward increasing rate of in-hospital major adverse cardiovascular outcomes in ST-elevation myocardial infarction undergoing fibrinolysis  
Correlation between serum p53 levels and vascular age determined by carotid intima media thickness (CIMT) in patients with intermediate cardiovascular risk factor  
Clinical significance of precordial ST segment depression on admission electrocardiogram in patients with acute inferior myocardial infarction  
Postprocedural high sensitivity C-reactive protein and the risk of recurrent ischemic events after percutaneous coronary intervention in patients with stable angina pectoris  
The messenger ribonucleic acid expression of B-type natriuretic peptide, natriuretic peptide receptor type-A and type-C in cardiomyocytes of obese population  
The correlation between serum galectin-3 level with wall motion score index in patients with ST-segment elevation myocardial infarction  
Risk

factors of atrial fibrillation in patients with chronic heart failure  
QT interval prolongation after non-ST elevation myocardial infarction in type 2 diabetic compared with nondiabetic patients  
The novel effect of polysaccharide peptides of to endothelial dysfunction and dislipidemia in high risk groups of atherosclerosis  
Relation of Left Atrial Spontaneous Echocardiographic Contrast to Neutrophil/Lymphocyte Ratio in Patient With Mitral Stenosis  
Analysis of characteristic of atrial fibrillation versus not atrial fibrillation in hospitalized heart failure patients  
Pulmonary hypertension features in adult atrial septal defect at RSUP Dr. Sardjito  
The association between plasma tryptase with ventricular remodeling after 1 month in patients with acute coronary syndrome  
The correlation of plasma histamine and tryptase with ST-elevation acute myocardial infarction in acute coronary syndrome patients  
Acute kidney injury after coronary artery bypass graft surgery: Incidence and pre operative risk profile  
The effect of diabetes mellitus on in-hospital mortality and major complications after isolated coronary artery bypass grafting surgery  
Factor those influence return to work after CABG revascularization on coronary artery disease patients  
The correlation of pulmonary vein

endothelin-1 level with pulmonary vascular resistance before and after mitral valve surgery with pulmonary hypertension Mitral valve E-Point septal separation as an independent predictor for in-hospital mortality after acute ST-elevation myocardial infarction Effects of allopurinol on complications of post coronary artery bypass graft (CABG) surgery in coronary artery disease's patient with left ventricular dysfunction Maternal and fetal outcomes of pregnancy with valvular heart disease QRS duration in ST elevation myocardial infarction patients undergoing a primary percutaneous intervention Knowledge of resident medical officers on hypertensive emergency: A descriptive study Hyperuricemia and its association with incident of congestive heart failure in Madurese population Diagnostic value of Duke Treadmill Score in predicting coronary lesions severity in patients with suspected stable coronary artery diseases Relationship of plasma glucose levels and troponin-I in patients wi...

Publication

8

Toni Sutono. "Efficacy of *Garcinia mangostana* L. (mangosteen rind extract) to reduce acne severity", *Medical Journal of Indonesia*, 2013

Publication

<1%

---

9	<a href="http://www.aafs.org">www.aafs.org</a> Internet Source	<1%
10	<a href="http://pure.bond.edu.au">pure.bond.edu.au</a> Internet Source	<1%
11	<a href="http://de.scribd.com">de.scribd.com</a> Internet Source	<1%
12	Olsen, A.R.. "Regulatory Action Criteria for Filth and Other Extraneous Materials", Regulatory Toxicology and Pharmacology, 199812 Publication	<1%
13	<a href="http://ejournal.undip.ac.id">ejournal.undip.ac.id</a> Internet Source	<1%
14	<a href="http://doaj.org">doaj.org</a> Internet Source	<1%
15	Comstock, Jenna L., Jean-Paul Desaulniers, Hélène N. LeBlanc, and Shari L. Forbes. "New decomposition stages to describe scenarios involving the partial and complete exclusion of insects", Canadian Society of Forensic Science Journal, 2014. Publication	<1%
16	Essam A. Shaalan, Zeinab A. El-Moaty, Salaheldin Abdelsalam, Gail S. Anderson. "A Preliminary Study of Insect Succession in Al-Ahsaa Oasis, in the Eastern Region of the	<1%

---

Kingdom of Saudi Arabia", Journal of Forensic Sciences, 2017

Publication

17

M. Lee Goff. "Early post-mortem changes and stages of decomposition in exposed cadavers", Experimental and Applied Acarology, 06/25/2009

Publication

<1%

18

Fer Coenders, Cees Terlouw, Sanne Dijkstra, Jules Pieters. "The Effects of the Design and Development of a Chemistry Curriculum Reform on Teachers' Professional Growth: A Case Study", Journal of Science Teacher Education, 2017

Publication

<1%

19

[www.termedia.pl](http://www.termedia.pl)

Internet Source

<1%

20

"South America", Contemporary Topics in Entomology, 2015.

Publication

<1%

21

Goodbrod, J. R., and M. L. Goff. "Effects of Larval Population Density on Rates of Development and Interactions between Two Species of Chrysomya (Diptera: Calliphoridae) in Laboratory Culture", Journal of Medical Entomology, 1990.

Publication

<1%

22

E. ANTON. "Beetles and flies collected on pig carrion in an experimental setting in Thuringia and their forensic implications", *Medical and Veterinary Entomology*, 12/2011

Publication

<1%

23

api.equinoxpub.com

Internet Source

<1%

24

Grassberger, M., and C. Frank. "Initial Study of Arthropod Succession on Pig Carrion in a Central European Urban Habitat", *Journal of Medical Entomology*, 2004.

Publication

<1%

25

documents.mx

Internet Source

<1%

26

Kittikhun Moophayak, Kabkaew L. Sukontason, Watcharapong Ruankham, Jeffery K. Tomberlin, Nophawan Bunchu. "Variation in the Time of Colonization of Broiler Carcasses by Carrion Flies in Nakhonsawan Province, Thailand", *Journal of Medical Entomology*, 2017

Publication

<1%

27

Terry L. Eberhardt, Douglas A. Elliot. "A preliminary investigation of insect colonisation and succession on remains in New Zealand", *Forensic Science International*, 2008

Publication

<1%



---

28

[www.geotechuk.com](http://www.geotechuk.com)

Internet Source

<1%

---

29

B. SINGH. "Molecular phylogeny of the blowfly genus *Chrysomya*", *Medical and Veterinary Entomology*, 06/2011

Publication

<1%

---

30

[www.scielo.br](http://www.scielo.br)

Internet Source

<1%

---

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On