

Manadoperoxides, a new class of potent antitrypanosomal agents of marine origin

by Henny Dien 9

Submission date: 13-Mar-2019 03:45PM (UTC+0700)

Submission ID: 1092548796

File name: new_class_of_potent_antitrypanosomal_agents_of_marine_origin.pdf (454.53K)

Word count: 10656

Character count: 51012

28
Manadoperoxides, a new class of potent antitrypanosomal agents of marine origin†Giuseppina Chianese,^a Ernesto Fattorusso,^a Fernando Scala,^a Roberta Teta,^a Barbara Calcinaï,^b Giorgio Bavestrello,^c Henny A. Dien,^d Marcel Kaiser,^{e,f} Deniz Tasdemir^g and Orazio Tagliatalata-Scafati^{*a}

Received 12th June 2012, Accepted 6th July 2012

DOI: 10.1039/c2ob26124c

1
Chemical investigation of the marine sponge *Plakortis* cf. *lita* afforded a library of endoperoxylketal polyketides, manadoperoxides B–K (3–5 and 7–13) and peroxyplakoric esters B₃ (6) and C (14). Eight of these metabolites are new compounds and some contain an unprecedented chlorine-bearing THF-type ring in the side chain. The library of endoperoxide derivatives was evaluated for *in vitro* activity against *Trypanosoma brucei rhodesiense* and *Leishmania donovani*. Some compounds, such as manadoperoxide B, exhibited ultrapotent trypanocidal activity (IC₅₀ = 3 ng mL⁻¹) without cytotoxicity. Detailed examination of the antitrypanosomal activity data and comparison with those available in the literature for related dioxane derivatives enabled us to draw a series of structure–activity relationships. Interestingly, it appears that minor structural changes, such as a shift of the methyl group around the dioxane ring, can dramatically affect the antitrypanosomal activity. This information can be valuable to guide the design of optimized antitrypanosomal agents based on the dioxane scaffold.

1. Introduction

57
African trypanosomiasis (also called sleeping sickness) is a 47
disease caused by the single-celled protozoan parasites *Trypanosoma brucei gambiense* (in Western and Central Africa) and *T. 34
rhodesiense* (in Eastern and Southern Africa) and transmitted to humans by bites of tsetse flies (*Glossina* genus), which have acquired their infection from human beings or from animals harbouring the hu 15
pathogenic parasites.¹ Once injected into the humans, trypanosomes multiply in subcutaneous tissues, blood and lymph and, in a second stage, they 2
cross the blood-brain barrier and attack the central nervous system. The process can take years with *T. b. gambiense*, while *T. b. rhodesiense* is responsible for acute infections. When the

2
parasites invade the central nervous system, 2
they lead to neurological symptoms, including behavioural changes, coma, and ultimately, if untreated, death. Disturbance of the sleep cycle, which gives the disease its name, is a characteristic feature of this second stage of the disease.

Recent figures ha 2
estimated that sleeping sickness threatens at least 60 million people in 36 countries with the number of infections ranging between 50 000 and 70 000 per year, thus qualifying 15
as the world's third most devastating tropical disease.² The rural populations living in regions where transmission occurs and which depend on agriculture, fishing, animal husbandry or hunting are the most exposed to the bite of the tsetse fly and therefore to the disease. Socio-political instability and the consequent population movements, combined with a shortage of funds have largely contributed to determining the recent spread of trypanosomiasis, which, together with other tropical diseases, results in billions of dollars of lost productivity, contributing to determine the poverty and the social isolation of th 2
regions.

Treatment of African trypanosomiasis is still today based on very 2
few molecules, almost all introduced about one century ago. Pentamidine and suram 2
are the preferred options for the treatment of the first stage, while to treat the second, cerebral stage, eflornithine and the trivalent arsenic derivative melarsoprol are practically the only therapeutic options.³ Melarsoprol has many disadvantages since it can induce a reactive encephalopathy (encephalopathic syndrome) which can be fatal, while eflornithine is very expensive and ineffective against *T. b. rhodesiense*. Moreover, an increase of cross-resistance to

27
^aDipartimento di Chimica delle Sostanze Naturali, Università di Napoli "Federico II", Via D. Montesano, 49, 80131 Napoli, Italy; E-mail: scatagli@unina.it; Fax: +(0039) 081-678552; Tel: +(0039) 0 26 78509

^bDipartimento di Scienze del Mare, Università Politecnica delle Marche, Via Brecce Bianche, 60131 Ancona, Italy

^cDipartimento per lo Studio del Territorio e delle sue Risorse, Università 44
nuova, Corso Europa 26, 16132 Genova, Italy

^dFaculty of Fishery and Marine Science, Sam Ratulangi University, Manado, Ind 36
ia

^eDepartment of Medical Parasitology and Infection Biology, Swiss Tropical and Public Health Institute, CH-4002 Basel, Switzerland
^fUnivers 61
f Basel, Petersplatz 1, CH-4003 Basel, Switzerland

^gSchool of Chemistry, National University of Ireland, Galway, University 70
K Galway, Ireland

†Electronic supplementary information (ESI) available: 1D and 2D NMR spectra for compounds 7–14. See DOI: 10.1039/c2ob26124c




Fig. 1 Chemical structures of known compounds: plakortin (1), manadoperoxides A–D (2–5) and peroxyplakoric ester B₃ (6).

these drugs has been observed in recent times, especially in central Africa and, thus, the few available drugs are rapidly losing their effectiveness.⁴ In recent years, only one compound, pafuramidine (a synthetic analogue of pentamidine), has been clinically evaluated against African trypanosomiasis, but further development was discontinued in 2008 due to liver toxicity.⁵ Recently, the nitroimidazole derivative fexinidazole, bearing structural similarities with metronidazole and nifurtimox, has entered Phase II/III clinical trials⁶ and is the most advanced drug candidate against trypanosomiasis. However, there is still an urgent need to find new, effective and, above all, affordable alternatives to the existing options for treatment of sleeping sickness.

We have been actively engaged in the field of tropical disease treatment and discovered that some classes of natural products are endowed with promising activity against *Plasmodium falciparum* and *Leishmania* parasites.^{7–9} Among them, plakortin (1, Fig. 1), a simple 1,2-dioxane polyketide obtained from the Caran sponge *Plakortis simplex*⁷ was demonstrated to possess significant *in vitro* antimalarial activity against CQ-resistant *P. falciparum* strains without cellular toxicity.⁷ By means of a multidisciplinary approach (including computational and experimental studies) on natural analogues^{10,11} and semisynthetic derivatives,¹² we also obtained valuable information about the SARs needed for the antimalarial activity of these simple 1,2-dioxanes. These results clearly indicated the crucial role of the endoperoxide functionality, suggested the importance of the alkyl side chain, and revealed conformation-dependent features critical for antimalarial activity.¹³ We recently developed two different synthetic approaches for the preparation of simplified plakortin analogues and some of the prepared compounds succeeded in reaching the *in vitro* antimalarial potency of plakortin.^{14,15}

In this context, we have recently reported the results of the chemical investigation of the Indonesian sponge *Plakortis* cf. *simplex*,¹⁶ which resulted in the isolation of four endoperoxyketal derivatives named manadoperoxide A–D (2–5, Fig. 1),¹⁷ These compounds were assayed *in vitro* against CQ-R and CQ-S *P. falciparum* strains and showed relatively poor antimalarial

potency (5–10 μM), more than ten times lower than that exhibited by plakortin (1).

As part of our ongoing screening of natural sources to find antiprotozoal lead compounds, we have undertaken the chemical analysis of the sponge *Plakortis* cf. *lita* de Laubenfels, a species widely distributed in the Indo-West Pacific,¹⁸ which was collected along the coasts of the Bunaken Marine Park of Manado (North Sulawesi, Indonesia). Careful inspection of the organic extract of this organism afforded the known manadoperoxides B–D (3–5) and peroxyplakoric ester B₃ (6) (Fig. 1), along with eight new peroxyketal polyketides, which we named manadoperoxides E–K (7–13) and peroxyplakoric ester C (14) (Fig. 2). Herein we provide details about their isolation, stereostructural determination and SARs, as well as *in vitro* trypanocidal and leishmanicidal effects. The present study provides the first data on the antitrypanosomal activity of endoperoxyketal derivatives.

2. Results

2.1 Chemistry

The sponge *Plakortis* cf. *lita* de Laubenfels (order Homosclerophorida, family Plakinidae) was collected in January 2010 from along the coasts of Bunaken Island (Manado, Indonesia). After homogenization, the organism was exhaustively extracted sequentially with MeOH and CHCl₃. The combined extracts were subjected to chromatography over reversed-phase silica column eluting with a solvent gradient of decreasing polarity in water to chloroform. Selected fractions were combined and further fractionated by gravity column chromatography (CC) on silica gel, followed by repeated open CC and CC (*n*-hexane–EtOAc mixtures) to afford the new polyketide manadoperoxides E–K (7–13) and peroxyplakoric ester C (14) in the pure state, along with the known manadoperoxides B–D (3–5)¹⁷ and peroxyplakoric ester B₃ (6).¹⁹ The structures of the known compounds 3–6 were deduced by comparing their spectroscopic data with those previously reported,^{17,19} while the structures of the new compounds 7–14 have been deduced as follows.




Fig. 2 Chemical structures of the new compounds manadoperoxides E–K (7–13) and peroxyplakoric ester C (14). Asterisks indicate that the configuration of the moiety is relative.

Compounds 7–14 have been subjected to detailed spectroscopic (ESI-MS) and spectroscopic (mainly 1D and 2D NMR) analysis, which allowed the definition of their stereostructures (Fig. 2) and the complete assignment of NMR resonances (Tables 1 and 2 or Experimental section). In particular, the NMR analysis revealed that both the ^{13}C NMR resonances from C-1 to 69, including 4-Me and 6-OMe (moiety A), and the ^1H NMR resonances and coupling constants from H₂-2 to H₂-5 were practically identical within the entire series of compounds 3–5 and 7–13. This spectroscopic evidence clearly indicated that the new compounds 7–13 must share with the co-occurring manadoperoxides B–D (3–5) the structure of the six-membered dioxanated ring, including the relative configuration at the stereogenic carbons present in this moiety (2D NMR ROESY cross-peaks fully supported this stereochemical assignment). Since we have previously assigned the absolute configuration at all the stereogenic carbons of manadoperoxides A–D (2–5) through chemical derivatization and application of the Mosher's method,¹⁷ it is biosynthetically reasonable to assume that the configurations at C-3, C-4 and C-6 reported for compounds 7–13 in Fig. 2 are also the absolute ones. Thus, the new compounds 7–13 differ from manadoperoxides A–D exclusively by the structure of the long ("western") side chain (moiety B) and we will here describe in detail only the determination of this part of their structures.

Manadoperoxide E (7) was isolated as a colorless amorphous solid with molecular formula C₁₉H₃₄O₇, which, taking into account the structure of the dioxane ring, implied that moiety B should have a C₁₀H₁₉O₂ composition and include a single unsaturation degree. The ^1H and ^{13}C NMR spectra of 7 (Experimental section), analyzed with the help of COSY and HSQC data, revealed the presence of two spin systems which closely paralleled those present in the structure of manadoperoxide D (5). In particular, the first spin system, from H₂-7 to the sp² methine

H-11 (δ_{H} 5.45, δ_{C} 127.1) with an oxymethine at H-10 (δ_{H} 4.40, δ_{C} 70.1), is identical to that present in the structure of 5, while the second spin system included a second oxymethine (H-13) attached to $^{12}\text{CH}_2\text{CH}_3$ moiety. The HMBC cross-peaks of 12-Me with C-11, C-12, and C-13 allowed the connection of the above moieties and defined the gross structure of the side chain (moiety B) for manadoperoxide E, which is a homologue of manadoperoxide D. The ROESY cross-peak of 12-Me with H-10 indicated that also in this case the side chain double bond possessed *E* geometry. Following the same procedure readily applied for manadoperoxide D, we determined the absolute configuration at the two stereogenic carbons C-10 and C-13 of 7 through application of the Riguera method for 1,*n*-diols.²⁰ Thus, two aliquots of 7 were allowed to react with *R*- and *S*-MTPA chloride obtaining the diesters 7a and 7b, respectively. According to the Riguera model,²⁰ in 1,4 MTPA diesters, the combined shielding of the two auxiliary reagents induces positive $\Delta\delta$ (*S*-*R*) values for the carbinol protons and for all the protons between them. Thus, the positive $\Delta\delta$ (*S*-*R*) values measured for protons going from H-9 to H-14 (Fig. 3) indicated a 10*S*,13*S* configuration.

The molecular formula of manadoperoxide F (8), determined by HR-ESIMS, was identical with that of compound 7, suggesting the same C₁₀H₁₉O₂ composition for the side chain. However, analysis of 1D NMR spectra (Tables 1 and 2), together with 2D COSY and HSQC NMR, showed that the moiety B of 8 significantly differed from that of 7 and included a methoxy group (δ_{H} 3.17, δ_{C} 49.3), an sp³ oxymethine (δ_{H} 3.60, δ_{C} 73.6) and two mutually coupled sp² methines (δ_{H} 5.67, δ_{C} 126.8; δ_{H} 5.49, δ_{C} 132.2). The COSY spectrum arranged all the proton multiplets of the side chain of 8 within two spin systems: the first spans from H₂-7 to the double bond,^{10,11} while the second spin system includes only the oxymethine (H-13) coupled with a

68

Table 1 ^1H NMR (500 MHz) data of compounds **8–10** and **12–13** in CDCl_3

Pos.	32 δ_{H} , mult. (<i>J</i> in Hz)	9 δ_{H} , mult. (<i>J</i> in Hz)	10 δ_{H} , mult. (<i>J</i> in Hz)	12 δ_{H} , mult. (<i>J</i> in Hz)	13 δ_{H} , mult. (<i>J</i> in Hz)
43					
2a	2.91, dd (15.6, 9.5)	2.92, dd (15.5, 9.4)	2.92, dd (15.5, 9.5)	2.92, dd (15.7, 9.5)	2.94, dd (15.6, 9.4)
2b	2.52, dd (15.6, 4.2)	2.43, dd (15.5, 4.5)	2.43, dd (15.5, 3.8)	2.42, dd (15.7, 3.5)	2.44, dd (15.6, 3.8)
3	4.37, ddd (9.5, 4.2, 3.0)	4.46, ddd (9.4, 4.5, 3.0)	4.46, ddd (9.5, 3.8, 3.0)	4.44, ddd (9.5, 3.5, 3.0)	4.44, ddd (9.4, 3.8, 3.0)
4	2.50, m	2.57, m	2.56, m	2.57, m	2.57, m
5a	1.68, dd (13.3, 4.3)	1.68 ^a	1.71 ^a	1.70 ^a	1.70 ^a
5b	1.39 ^a	1.30 ^a	1.32 ^a	1.32 ^a	1.31 ^a
7a	1.66 ^a	1.63 ^a	1.58 ^a	1.69 ^a	1.59 ^a
7b	1.62 ^a	1.60 ^a	1.55 ^a	1.32 ^a	1.55 ^a
8a	1.46, m	1.53, m	1.30 ^a	1.26 ^a	1.44 ^a
8b	1.46, m	1.47, m			1.32 ^a
9a	2.14, q (7.1)	1.76, m	1.32 ^a	1.53 ^a	1.72 ^a
9b		1.59 ^a		1.30 ^a	1.51 ^a
10	5.67, dt (15.9, 7.1)	3.08, m	1.24 ^a		3.92, m
11	5.49, d (15.9)	3.20, d (1.7)	1.68 ^a	4.55, bs	3.50, d (7.4)
12			2.50, m		
13a	3.60, q (6.4)	2.08, s	1.66 ^a	4.30, t (7.3)	3.74, t (7.3)
13b			1.41 ^a		
14a	1.08, d (6.4)		1.33 ^a	2.19, dq (14.2, 7.1)	1.50 ^a
14b				1.97, m	1.33 ^a
15			0.88, t (7.0)	1.00, t (7.1)	1.02, t (7.2)
4-Me	0.87, d (7.0)	0.87, d (7.1)	0.85, d (7.0)	0.84, d (7.0)	0.85, d (7.0)
12-Me	1.20, s	3.73, s	9.79, d (2.5)	5.10, bs	1.25, s
1-OMe	3.71, s	3.27, s	3.73, s	5.29, bs	3.73, s
6-OMe	3.23, s		3.28, s	3.73, s	3.28, s
12-OMe	3.17, s			3.28, s	3.33, s

^a Overlapped with other signals.**Table 2** ^{13}C NMR data (125 MHz) of compounds **8–10** and **12–13** in CDCl_3

Pos.	8 δ_{C} , mult.	9 δ_{H} , mult.	10 δ_{H} , mult.	12 δ_{H} , mult.	13 δ_{H} , mult.
1	173.0, s	172.4, s	171.9, s	171.9, s	172.3, s
2	31.0, t	31.2, t	31.1, t	30.9, t	30.8, t
3	80.1, d	79.9, d	80.1, d	79.6, d	80.3, d
4	27.2, d	27.4, d	26.9, d	26.9, d	27.1, d
5	34.2, t	34.6, t	33.9, t	34.0, t	34.6, t
6	103.7, s	102.8, s	103.0, s	102.8, s	103.1, s
7	32.4, t	32.0, t	32.1, s	33.5, t	33.2, t
8	23.4, t	22.5, t	21.8, t	23.5, t	22.8, t
9	32.6, t	36.5, t	30.1, t	30.0, t	30.8, t
10	126.8, d	57.6, d	28.7, t	116.2, s	79.0, d
11	132.2, d	59.5, d	32.2, t	59.7, d	68.2, d
12	80.7, s	205.8, s	43.5, s	141.6, s	82.4, s
13	73.6, d	24.6, q	32.0, t	83.7, d	84.3, d
14	16.6, q		22.8, t	26.1, t	25.5, t
15			14.2, q	10.4, q	17.0, q
4-Me	16.0, q	16.3, q	16.9, q	17.0, q	17.3, q
12-Me	14.3, q	52.0, q	201.1, d	105.6, t	14.0, q
1-OMe	52.3, q	48.5, q	51.7, q	52.0, q	51.9, q
6-OMe	47.7, q		48.5, q	48.5, q	48.4, q
12-OMe	49.3, q				53.1, q

methyl doublet (Me-14). The HMBC cross-peaks of the methyl singlet at δ_{H} 1.20 (12-Me) with C-11 and the oxygenated C-12 (δ_{C} 80.7) and C-13 (δ_{C} 73.6) and of the methoxy protons with C-12, determined the connection for the above moieties, thus defining the planar structure of the side chain of manadoperoxide F as shown in Fig. 2. The value $J_{\text{H-10,H-11}} = 15.9$ Hz indicated the *E* geometry of Δ ; while, due to the very small amounts of manadoperoxide F isolated (~1.0 mg), needed for

pharmacological evaluation, we could not attempt any determination of the configuration at the two stereogenic carbons of the side chain.

The molecular formula of manadoperoxide G (**9**) ($\text{C}_{16}\text{H}_{26}\text{O}_7$) was indicative of a $\text{C}_7\text{H}_{11}\text{O}_2$ side chain, implying two unsaturation degrees. ^1H and ^{13}C NMR data of **9** (Tables 1 and 2), analyzed with the help of the 2D HSQC spectrum, sorted out the seven carbon atoms of this moiety as three methylenes, two methines resonating at δ_{H} 3.08 and 3.20, methyl singlet (δ_{H} 2.08) and a ketone carbonyl (δ_{C} 205.8). The 2D NMR COSY spectrum grouped all the proton multiplets of the side chain in a single spin system terminating with the oxymethines H-10 and H-11. The relatively high field ^{13}C NMR resonances of C-10 and C-11 (δ_{C} 57.6 and 59.5, respectively) were strongly suggestive of the presence of an epoxide group, which also accounted for the remaining unsaturation degree.

The 2D HMBC cross-peaks of the methyl singlet at δ_{H} 2.08 with the ketone carbonyl (C-12) and with the epoxide carbon C-11 allowed the connection between the two moieties and the definition of the planar structure of manadoperoxide G (**9**), which can be viewed as an epoxidated manadoperoxide C (**4**). The ROESY correlation between H-11 (δ_{H} 3.20) and H₂-9 indicated the *trans* orientation of the epoxide protons. It should be noted that this relative configuration at C-10 and C-11 has not been connected with that of the remaining stereogenic carbons of the molecule.

Manadoperoxide H (**10**), $\text{C}_{19}\text{H}_{34}\text{O}_6$ by HR-ESIMS, should possess a $\text{C}_{10}\text{H}_{19}\text{O}$ composition for moiety B. The presence of an aldehyde carbonyl was suggested by the NMR resonances at δ_{H} 9.79 and δ_{C} 201.1 (associated by means of the 2D HSQC spectrum), and this group accounted for the single oxygen atom




Fig. 3 Application of the modified Riguera method to manadoperoxide E (7). $\Delta\delta_{(S-R)}$ values are in ppm.




Fig. 4 COSY and key HMBC correlations for the side chains of manadoperoxides I–K.

and the single unsaturation degree implied for the side chain. Accordingly, the 2D COSY spectrum revealed the presence of a single spin system spanning from H₂-7 to Me-15 (δ_{H} 0.88, t) and including a methine resonating at δ_{H} 2.50 (H-12) coupled to the doublet at δ_{H} 9.79. Thus, the aldehyde group should be attached C-12, as further confirmed by its HMBC cross-peaks with C-11, C-12 and C-13. The configuration at the stereogenic C-12 has been left undetermined.

Manadoperoxide I (11) was found to have the molecular formula C₁₈H₃₀O₇ by HR-ESIMS, thus implying a C₉H₁₅O₂ side chain with two unsaturation degrees. Analysis of ¹H and ¹³C NMR spectra (C₆D₆, see Experimental) of 11, guided by the 2D HSQC spectrum, revealed the presence of an sp² methine singlet (δ_{H} 5.24, δ_{C} 122.7), an sp³ oxymethine (δ_{H} 4.19, δ_{C} 77.6) and an hemiketal carbon resonating at δ_{C} 102.1. In the COSY spectrum, two spin systems (highlighted in bold in Fig. 4) were visible: the first includes a sequence of three methylenes, while the second encompasses only the oxymethine proton coupled to a methyl group. The HMBC cross-peaks of H₂-9 with the hemiketal carbon C-10 and with the sp² methine C-11 and those of the allylic methyl singlet 12-Me with C-11, C-12 and with the oxymethine C-13 provided the information needed to link the above moieties. Moreover, the HMBC cross-peak H-13/C-10 indicated the presence of the oxygen bridge between C-13 and C-10, thus accounting for the remaining unsaturation degree and completing the gross structure of manadoperoxide I (11). A series of signals attributable to a minor compound were detectable both in ¹H and ¹³C NMR spectra of 11 and all our initial attempts to further purify compound 11 failed. However, once the structure of manadoperoxide I was disclosed, these signals were easily rationalized with the presence of an equilibrating mixture (ca. 4 : 1 ratio) of the two epimers at C-10. Selected ¹H NMR resonances of the minor epimer have been reported in the Experimental section.

Manadoperoxide J (12), C₁₉H₃₁ClO₇ by HR-ESIMS, was demonstrated to share with manadoperoxide I (11) the presence of an oxygenated five-membered ring and an hemiketal functionality at C-10, in the moiety B. Accordingly, also in this case we observed the presence of an equilibrating mixture (ca. 4 : 1 ratio) of the two epimers at C-10. A detailed analysis of ¹H and ¹³C NMR data of the major epimer of 12 (Tables 1 and 2, respectively), aided by 2D NMR spectra, revealed that manadoperoxide J (12) actually differs from manadoperoxide I (11) by possessing: (i) an ethyl group attached at the oxygenated C-13 (see the COSY cross-peaks shown in Fig. 4) in place of the methyl group; (ii) an sp² methylene (δ_{H} 5.29 and 5.10, δ_{C} 105.6) instead of the allylic methyl group (iii) a chlorine-bearing sp³ methine (δ_{H} 4.55, δ_{C} 59.7) instead of the sp² methine. The 2D HMBC cross-peaks shown by H₂-9 (C-10 and C-11) and by the hexomethylene protons (with C-11, C-12 and C-13), together with the cross-peak H-13/C-10, proved to be crucial to join these moieties as indicated in Fig. 4. The ROESY cross-peak H-11/H-13 revealed the *cis* relative orientation of these protons, but it was not possible to connect this relative configuration with that of the stereogenic carbons of the dioxane ring.

Manadoperoxide K (13), C₂₀H₃₅ClO₇ by HR-ESIMS, should possess a C₁₁H₂₀ClO₂ side chain with a single unsaturation degree. The ¹H NMR spectrum of 13 (Table 1) revealed, in addition to the signals of moiety A, the resonances of a third methoxy group (δ_{H} 3.33) and of three methines (δ_{H} 3.92, 3.50 and 3.74) in the midfield region of the spectrum. All the proton signals were associated to those of the directly linked carbon atoms by means of a 2D HSQC experiment, and then identification of the 2D COSY experiment disclosed the existence of two spin systems, shown in bold in Fig. 4. The first spin system spans from H₂-7 to the chlorine-bearing methine H-11 (δ_{H} 3.50, δ_{C} 68.2) and includes the oxymethine at δ_{H} 3.92 (δ_{C} 79.0). The second spin system spans from the oxymethine H-13 (δ_{H} 3.74)

to methyl triplet Me-15. The remaining unprotonated carbon atom (δ_C 82.4) present in the ^{13}C NMR spectrum of **13** was attributed to a methoxy-bearing carbon (C-12) located between the two above described moieties on the basis of the $^{2,3}J_{\text{C,H}}$ HMBC cross-peaks of the methyl singlet at δ_H 1.24 with C-12, C-11 and C-13 and of the cross-peak of the methoxy group (12-OMe) with C-12. The presence of an ether bridge between C-10 and C-13 to define a THF-type ring was inferred by the HMBC cross-peak H-13/C-10 and accounted for the single unsaturation degree present in the side chain of **13**. This THF-ring includes four stereogenic carbons, whose relative configuration was deduced by the ROESY cross-peaks 12-Me/H₂-9, 12-Me/H-11, H-11/H-13, and H₂-9/H-13. As for compounds **9** and **12**, this relative configuration has not been connected with that of the stereogenic carbons of the dioxane ring.

Peroxyplakoric ester C (**14**), $\text{C}_{17}\text{H}_{28}\text{O}_6$ by HR-ESIMS, proved to belong to the structural series of peroxyplakoric esters,¹⁹ which differ from manadoperoxides by the substituents attached at positions 2 and 4 of the dioxane ring (see Fig. 1). A detailed analysis of its ^1H and ^{13}C NMR spectra (see Experimental section), accomplished by inspection of 2D COSY and HSQC NMR spectra, revealed that compound **14** shared with peroxyplakoric ester B₃ (**6**)¹⁹ all the carbon and proton resonances, along with the proton–proton coupling constants, for the positions C-1 to C-6 (including also 2-Me and 6-OMe). On the other hand, ^1H and ^{13}C NMR resonances of the side chain of **14** were practically superimposable to those of manadoperoxide C (**4**), including an α,β -unsaturated ketone group (δ_C 198.7). The 2D HMBC spectra provided key information to support these assignments and to join the above partial moieties, thus fully confirming the structure of peroxyplakoric ester C (**14**).

Thus, analysis of the organic extract of *P. cfr. lita* afforded twelve endoperoxyketal-containing compounds, eight of which are being described for the first time. These compounds, with the exception of **6** and **14**, share the same stereostructure of the dioxane ring while they differ by the length and/or the functionalization of the “western” side chain. Recently, Garson and co-workers have reported the isolation of a series of dioxane derivatives from the sponge *Plakinastrella clathrata* differing by modifications in the side chain,²¹ evidencing the existence of an extraordinary enzymatic activity in Plakinidae sponges, especially dedicated to the functionalization of side chains. However, some of the metabolites isolated in the course of this study are unique in many extents: three of them show the formation of an unprecedented five-membered oxygenated ring which, in two cases, bears also a chlorine atom. To the best of our knowledge this has never been reported for marine endoperoxides.

2.2 Antiprotozoal activity

The library of endoperoxide derivatives obtained from the chemical investigation of *P. cfr. lita* have been evaluated *in vitro* for their antiprotozoal activity against *Trypanosoma brucei rhodesiense* and *Leishmania donovani*. Results are reported in Table 3.

Some of the tested compounds showed an ultrapotent activity against *T. b. rhodesiense* at low ng mL⁻¹ range. In particular,

Table 3 *In vitro* antiprotozoal activity (IC₅₀) and calculated log*P* values of compounds **3–6**, **8–11**, and **13–14**^a

Compounds	<i>T. b. rhodesiense</i>	<i>L. donovani</i>	log <i>P</i> ^b
Manadoperoxide B (3)	0.003	0.589	4.84
Manadoperoxide C (4)	0.678	3.24	2.52
Manadoperoxide D (5)	36.7	19.2	2.33
Peroxyplakoric ester B ₃ (6)	3.61	12.0	4.53
Manadoperoxide F (8)	0.792	5.73	2.82
Manadoperoxide G (9)	1.84	3.22	1.75
Manadoperoxide H (10)	0.375	2.44	4.25
Manadoperoxide I (11)	0.062	0.633	3.00
Manadoperoxide K (13)	0.087	1.89	3.63
Peroxyplakoric ester C (14)	30.9	43.4	2.55
Reference compound	0.003 ^c	0.174 ^d	—


^a IC₅₀ values are in $\mu\text{g mL}^{-1}$ and mean values from at least two replicates (the variation is max. 20%). ^b Calculated by Virtual Computational Chemistry Laboratory software. ^c Melarsoprol. ^d Miltefosine.

manadoperoxide B (**3**) (IC₅₀ = 3 ng mL⁻¹) proved to be as active as the reference compound and, to our knowledge, it can be viewed as the most potent trypanocidal marine natural product reported to date. Also remarkably active agents are manadoperoxides I (**11**) and K (**13**) (IC₅₀ = 60–90 ng mL⁻¹) and manadoperoxides H (**10**), C (**4**) and F (**8**) (IC₅₀ = 300–800 ng mL⁻¹), although their potencies are, respectively, about 30 and 200 times lower than that of manadoperoxide B. The remaining manadoperoxides and the two peroxyplakoric esters proved to be much less active, with IC₅₀s in the low–medium $\mu\text{g mL}^{-1}$ range. The tested compounds displayed moderate to good activity against *L. donovani*. The most active compounds of the series were manadoperoxide B (IC₅₀ = 0.589 $\mu\text{g mL}^{-1}$) and manadoperoxide I (IC₅₀ = 0.633 $\mu\text{g mL}^{-1}$) with comparable activities to the reference compound miltefosine (IC₅₀ = 0.174 $\mu\text{g mL}^{-1}$). The remaining compounds showed activity in the $\mu\text{g mL}^{-1}$ range. All the test compounds proved to be non-toxic (IC₅₀ > 10 $\mu\text{g mL}^{-1}$) towards the human cell line HMEC-1 (human microvascular endothelial cell line, data not shown). The calculated partition coefficient (log*P*) values of tested metabolites vary between 1.75 to 4.84 (Table 3), hence they comply with Lipinski's rule of five.²² This also indicates that manadoperoxides will be able to penetrate through lipid cellular membranes and potentially through the blood brain barrier (BBB), which is critical for the treatment of cerebral stage of *Trypanosoma* infections.

3. Discussion

As part of our ongoing screening of natural sources to find new antiprotozoal lead compounds,^{7–15} we have now discovered that the endoperoxyketal derivatives manadoperoxides constitute a new class of potent antitrypanosomal agents. The structural variability present within the library of endoperoxides tested in this study clearly defines some structure–activity relationships.

The comparison of the activities of manadoperoxides with those of the closely related peroxyplakoric esters provides valuable information. Indeed, in addition to a negligible difference in the side chain, manadoperoxide B (**3**) and peroxyplakoric ester



<i>Plasmodium falciparum</i>	IC ₅₀ = 2.30 μg/mL	IC ₅₀ = 0.040 μg/mL
<i>Trypanosoma brucei rhodesiense</i>	IC ₅₀ = 0.003 μg/mL	IC ₅₀ = 3.61 μg/mL

Fig. 5 Comparison of the structures and the antiprotozoal activities of manadoperoxide B (3) and peroxyplakoric ester B₃ (6).

B₃ (6) differ ⁴⁰ by the position of a methyl group on the ring moiety. In 3 a methyl group is present on the dioxane ring (at C-4), while in 6 it is attached on the short ester-containing side chain (at C-2) (Fig. 5). Interestingly, this minor structural difference is able to induce dramatic effects both in the antimalarial and in the antitrypanosomal activities, but in opposite directions. Indeed, while 6 has been demonstrated to be about 60-fold more active as an antimalarial agent,¹⁷ manadoperoxide B (3) is more than 1000-fold more potent as an antitrypanosomal agent. This trend is fully supported by the antitrypanosomal activity of manadoperoxide C (4, IC₅₀ = 0.678 μg mL⁻¹) (Fig. 1) and peroxyplakoric ester C (14, IC₅₀ = 30.9 μg mL⁻¹) (Fig. 2). These two compounds share the same structure of the “western” side chain and, thus, they differ exclusively by the shift of a methyl group from C-4 (in 4) to C-2 (in 14). As observed for the pair 3/6, this structural change induces a marked reduction of the antitrypanosomal activity in 14.

These results seem to exclude a non-specific antiprotozoal activity for these peroxyketal derivatives, strongly suggesting their molecular interaction with a defined target(s). In this regard, our studies on plakortin derivatives provided indications about the mechanism of the antitrypanosomal action of simple 1,2-dioxanes¹³ which, upon reaction with Fe(II) (most likely the heme iron), should generate an oxygen radical, simultaneously transferred to a side chain carbon. The so formed C-centered radical should then represent the toxic species responsible for the antimalarial activity. In the case of manadoperoxides and peroxyplakoric esters we already demonstrated that the different pattern of substitution around the dioxane ring can induce different conformational preferences of this ring and, consequently, different accessibilities for the interaction with heme iron.¹⁷ Intriguingly, the situation for the antitrypanosomal activity seems to be completely different. In this case, the different pattern of substitution around the dioxane ring and the consequent different conformational behaviour appear to play in favour of the antitrypanosomal activity of manadoperoxides. It is also interesting to notice that the above data suggest, indirectly, that the target(s) of these endoperoxyketal compounds might be different in the two different protozoan parasites.

The complete homogeneity of the stereostructure of the dioxane ring (moiety A) in the series of manadoperoxides B–K can allow a clear evaluation of the impact on the trypanocidal activity of modifications on the alkyl side chain (moiety B). The most active compound of the series, manadoperoxide B (3), shows a methyl branched C₉ side chain including a conjugated

diene system, while the least active compound is manadoperoxide D (5), where the presence of a diol group provokes an almost complete loss of activity. The negative effects of an increase in the side chain polarity is fully supported by the trend of activities measured for the other manadoperoxides. Manadoperoxides I (11) and K (13), bearing a five-membered ring in the side chain, retain very good antitrypanosomal activity, while the carbonyl-containing manadoperoxides H (10, saturated C₉ alkyl chain with an aldehyde group in place of the methyl group) and C (4, conjugated ketone group in side chain) evidence an about 100-fold lower activity compared to manadoperoxide B. An even lower activity was exhibited by manadoperoxides F (8, hydroxy group in side chain) and G (9, epoxy-ketone group in side chain).

An almost linear relationship between the antitrypanosomal activity and the lipophilicity of the moiety B is clearly indicated by the log*P* values calculated for the members of the manadoperoxide series (Table 3). This relationship suggests that this moiety is probably not directly involved in the interaction with the target(s), but is responsible for the pharmacokinetic behaviour of these compounds, thus modulating their activities.

Manadoperoxides represent the first endoperoxide derivatives bearing a ketal functional group on the dioxane ring to be evaluated against *Trypanosoma*. A comparison with data available in the literature for the antitrypanosomal activity of related 1,2-dioxanes seems to indicate a positive effect for the alkoxy group at C-6. Indeed, all the natural²³ and synthetic²⁴ analogues showing an ethyl group or an hydrogen atom in place of the methoxy group at C-6 proved to be less active than manadoperoxides. However, an unambiguous evaluation of this relationship is prevented by the differences in the side chain structures.

Endoperoxide-containing compounds have been reported to target several protozoan parasites and to be potential lead compounds against different tropical diseases. However, while some members of this class show a broad spectrum activity against *Plasmodium*, *Trypanosoma*, *Leishmania*, etc., other endoperoxide-bearing derivatives show a specific action against selected protozoa. Probably the most remarkable example of this second type of antiprotozoal agents is given by artemisinin, a well known antimalarial compound (activity against *P. falciparum* in the low nM range),²⁵ displaying a very modest activity (about 25 μg mL⁻¹) against *T. b. rhodesiense*.²⁵ As we have shown before, peroxyplakoric ester B₃ (6) and its analogue 14 follow the same behaviour as artemisinin. On the other hand, manadoperoxides show an opposite behaviour, possessing moderate




Fig. 6 Schematic view of the structure–activity relationships for anti-trypansomal 1,2-dioxanes.

activity against *Plasmodium falciparum* (3–10 μM), which is ca. 1000 times lower than the trypanocidal activity described herein (Fig. 5).

Since the cell organ⁴⁹ and the biochemical composition of trypanosomes differ in numerous aspects from those of mammalian cells, several parasite-specific processes may provide selective targets for drug development and, among these, purine salvage, fatty acid elongation, cytochrome-independent oxidase and trypanothione reductase have been suggested.^{26,27} This latter flavoenzyme is of key importance to maintain the redox balance within the parasite acting as regulator of the levels of trypanothione, a molecule composed by glutathione and spermidine and involved in the reduction of hydroperoxides and ribonucleotides.²⁷ Although a shortage of material prevented us from investigating the mechanism of action of manadoperoxides, an interference with the finely regulated redox balance within the parasite could be anticipated.

The analysis of our library of natural compounds and comparison with data⁵⁵ available in the literature have demonstrated unambiguously that the endoperoxy group does not confer *per se* activity against *Trypanosoma* (and other protozoan parasites responsible for tropical diseases), while the contribution of the carbon skeleton is of pivotal importance. We have summarized in Fig. 6 the most important structural requirements for the class of simple 1,2-dioxanes as they have emerged from the present study.

18 Experimental section

4.1 General experimental procedures

Low and high resolution ESI-MS spectra were performed on a LTQ OrbitrapXL (Thermo Scientific) mass spectrometer. ¹H (500 MHz) and ¹³C (125 MHz) NMR spectra were measured on Varian INOVA spectrometers. Chem³⁷ shifts were referenced to the residual solv⁴ t signal (CDCl₃: δ_{H} 7.26, δ_{C} 77.0; C₆D₆: δ_{H} 7.16, δ_{C} 128.1). Homonuclear ¹H connectivities were determined by the COSY experiment. Through-space ¹H connectivities were evidenced using a ROESY experiment with a mixing time of 500 ms. One-bond heteronuclear ¹H–¹³C connectivities were determined by the HSQC experiment; two- and three-bond ¹H–¹³C connectivities by gradient-HMBC experiments optimized for a ^{2,3}J of 8 Hz. Medium pressure liquid chromatography was performed¹⁴ on a Büchi apparatus using a silica gel (230–400 mesh) column. HPLC were achieved on a Knauer apparatus equipped with a refractive index detector and LUNA

(Phenomenex) SI60 or C18 (250 × 4 mm) columns⁴² the logP values of manadoperoxides were estimated using VCCLAB, Virtual Computational Chemistry Laboratory, online server (<http://www.vcclab.org>).³²

4.2 Animal material, extraction, isolation

A specimen of *Plakortis lita*⁹ de Laubenfels (order Homosclerophorida, family Plakinidae) was collected in January 2010 from along the coasts of the Bunaken island in the Bunaken Marine Park¹⁴ Manado. A voucher sample (Man/10/02) has been deposited at the Dipartimento di Chimica delle Sostanze Naturali, Università di Napoli Federico II. The sponge is massive, roundish; the colour *in vivo* and in the preserved state is light brown externally and internally. The sponge is very soft and easy to cut, but it contracts after preservation. Surface smooth in the living specimen, but after collection, especially in the dried state, the surface became irregular and wrinkled. Ostia are not detectable on the living specimens. Oscules with a very low rim, about 1.5 mm in diameter. The ectosomal skeleton is an irregular reticulation of rounded or elliptical meshes. In some area a disorganized arrangement of tangential spicules is present. Choanosome with a confused skeleton of scattered spicules. Spicules are straight or bent, thin diods (98 × 2.5 μm on average); rare triods (actines 33 μm on average); irregular strongyloid microrhabds (2.5–25 μm) and smooth ovoid spheres (about 2 μm in diameter). The studied material is very similar to *P. lita* de Laubenfels in the spicular complement, which consists of diods, triods and in irregular strongyloid microrhabds, comparable in size and similar in shape. A skeletal organisation of the ectosome with rounded meshes, typical of *P. lita* de Laubenfels, is also present. This species differs from *P. lita* de Laubenfels by the presence of tangential spicules irregularly arranged, never reported in *P. lita*, and by the colour which in this last species may be black, dark brown, or grey-brown. Spheres, detected in the present material, were not described in *P. lita*. The present sample has been compared also to *Plakortis hooperi* Muricy¹⁸ from Papua New Guinea. *P. hooperi* possesses a spicule complement, comparable to that of the present material, of diods, triods, microrhabds and spheres; it is characterized by a brown colour, similar to that of *P. cfr. lita* and by a confused ectosomal skeleton, without trace of reticulation. *P. hooperi* differs from *P. cfr. lita* in lacking an ectosomal reticulation of rounded meshes, in the shape and size of the diods, relatively thick and without a well-marked centre, in the shape and size of microrhabds (shorter and irregular in *P. hooperi*), and of the spheres (more spherical and larger). Pending deeper morphological studies, the present sample is provisionally assigned to *P. cfr. lita* on the basis of the numerous shared morphological characters.

After homogenization, the organism was exhaustively extracted, in sequence, with methanol and chloroform. The combined organic extracts (8.16 g) were subjected to chromatography over C18 silica column (200–400 mesh) eluting with a solvent gradient of decreasing polarity from water to chloroform. Fractions eluted with H₂O–MeOH 2 : 8¹⁰ and H₂O–MeOH 1 : 9 were combined (1.35 g) and further fractionated by gravity column chromatography⁴ on silica gel using a *n*-hexane–EtOAc gradient. Fractions eluted with *n*-hexane–EtOAc 9 : 1 were

subjected to repeated column and HPLC chromatographies (*n*-hexane–EtOAc 95:5) affording manadoperoxides B (3, 31 mg), J (12, 0.9 mg) and I (11, 1.3 mg) in the pure state. Fractions eluted with *n*-hexane–EtOAc 8:2 were re-chromatographed by HPLC (*n*-hexane–EtOAc 8:2, flow 0.8 mL min⁻¹) to give manadoperoxide K (13, 2.3 mg), F (8, 1.1 mg) and G (9, 1.3 mg). Fractions eluted with *n*-hexane–EtOAc 7:3 were re-chromatographed by HPLC (*n*-hexane–EtOAc 75:25) affording pure manadoperoxide C (4, 1.1 mg), H (10, 1.3 mg) and another fraction, further purified by RP-HPLC (MeOH–H₂O 85:15, flow 0.8 mL min⁻¹) to yield peroxyplakoric ester C (14, 1.3 mg). Fractions eluted with *n*-hexane–EtOAc (1:9) were re-chromatographed by RP-HPLC (MeOH–H₂O 6:4, flow 0.8 mL min⁻¹) affording manadoperoxide D (5, 2.1 mg) and E (7, 2.5 mg).

4.3 Manadoperoxide E (7)

Colorless amorphous solid; $[\alpha]_D^{25} -13.0$ (*c* 0.2, CHCl₃); ¹H NMR (CDCl₃, 500 MHz): δ_H 5.45 (1H, d, *J* = 8.7 Hz, H-11), 4.43 (60 ddd, *J* = 9.5, 4.3, 3.0 Hz, H-3), 4.40 (1H, m, H-10), 4.26 (1H, 9 H-13), 3.72 (3H, s, 1-OMe), 3.26 (3H, s, 6-OMe), 2.97 (1H, dd, *J* = 29.5, 9.5 Hz, H-2a), 2.57 (1H, m, H-4), 2.44 (1H, dd, *J* = 15.5, 4.3 Hz, 23), 1.71 (3H, bs, H-17), 1.69 (1H, overlapped, H-5a), 1.64 (1H, overlapped, H-9a), 1.62 (1H, overlapped, H-7a), 1.61 (2H, overlapped, H₂-14), 1.55 (1H, overlapped, 11 H-9b), 1.42 (1H, overlapped, H-8a), 1.35 (1H, overlapped, H-8b), 1.34 (1H, overlapped, H-7b), 1.00 (3H, t, *J* = 7.1 Hz, Me-15), 0.86 (7, d, *J* = 7.1 Hz, 4-Me); ¹³C NMR (CDCl₃, 125 7 Hz) δ_C 172.5 (s, C-1), 142.0 (s, C-12), 127.1 (d, C-11), 103.0 (s, C-6), 79.7 (d, C-3), 73.4 (33, C-13), 70.1 (d, C-10), 52.2 (q, 1-OMe), 86 7 (q, 6-OMe), 38.4 (t, C-9), 34.5 (t, C-5), 31.7 (t, C-2), 11 4 (t, C-7), 27.9 (t, C-14), 27.2 (d, C-4), 22.0 (t, C-8), 17.0 (q, C-7), 10.1 (q, C-15). (+) ESI-MS *m/z* 375 [M + H]⁺, 397 [M + Na]⁺. HR-ESIMS (positive ions): found *m/z* 397.2200, C₁₉H₃₄NaO₇ requires *m/z* 397.2202.

4.4 Reaction of manadoperoxide E (7) with *R*- and *S*-MTPA chloride

Compound 7 (1.0 mg, 2.8 μmol) was treated with *R*-MTPA chloride (30 μL) in 400 μL of dry pyridine with a catalytic amount of DMAP overnight at rt. Then, the solvent was removed and the product was purified by HPLC (*n*-hexane–EtOAc, 97:3) to obtain the *S*-MTPA diester 7a (1.2 mg, 60% yield). When compound 7 (1.0 mg, 2.8 μmol) was treated with *S*-MTPA chloride, following the same procedure, 1.5 mg (69% yield) of *R*-MTPA diester 7b was obtained. ***S*-MTPA diester 7a:** amorphous solid; ESI-MS (positive ions) *m/z* 807 [M + H]⁺; ¹H NMR (CDCl₃): selected values δ 5.61 (1H, m, H-10), 5.45 (1H, d, *J* = 25 Hz, H-11), 5.43 (1H, m, H-13), 1.77 (3H, s, 12-Me), 1.60 (1H, m, H-9a), 1.55 (2H, m, H-7), 1.40 (1H, m, H-9b), 1.34 (2H, overlapped, H₂-14). ***R*-MTPA diester 7b:** amorphous solid; ESI-MS (positive ions) *m/z* 29 07 [M + H]⁺; ¹H NMR (CDCl₃): selected values δ 5.59 (1H, m, H-10), 5.43 (1H, d, *J* = 8.7 Hz, H-11), 5.38 (1H, m, H-13), 4.37 (1H, m, H-3), 1.75 (25 s, 12-Me), 1.59 (1H, m, H-9a), 1.55 (2H, m, H-7), 1.39 (1H, m, H-9b), 1.62 (2H, overlapped, H₂-14).

4.5 Manadoperoxide F (8)

Colorless amorphous solid; $[\alpha]_D^{25} -12.1$ (*c* 0.1, CHCl₃); ¹H NMR (CDCl₃, 500 MHz): Table 1; ¹³C NMR (CDCl₃, 125 MHz): Table 2; (+) ESI-MS *m/z* 375 [M + H]⁺, 397 [M + Na]⁺. HR-ESIMS (positive ions): found *m/z* 397.2197, C₁₉H₃₄NaO₇ requires *m/z* 397.2202.

4.6 Manadoperoxide G (9)

Colorless amorphous solid; $[\alpha]_D^{25} -13.5$ (*c* 0.2, CHCl₃); ¹H NMR (CDCl₃, 500 MHz): Table 1; ¹³C NMR (CDCl₃, 125 MHz): Table 2; (+) ESI-MS *m/z* 331 [M + H]⁺, 353 [M + Na]⁺. HR-ESIMS (positive ions): found *m/z* 353.1510; C₁₆H₂₆NaO₇ requires *m/z* 353.1506.

4.7 Manadoperoxide H (10)

Colorless amorphous solid; $[\alpha]_D^{25} -18.1$ (*c* 0.1, CHCl₃); ¹H NMR (CDCl₃, 500 MHz): Table 1; ¹³C NMR (CDCl₃, 125 MHz): Table 2; (+) ESI-MS *m/z* 359 [M + H]⁺, 381 [M + Na]⁺. HR-ESIMS (positive ions): found *m/z* 381.2259; C₁₉H₃₄NaO₆ requires *m/z* 381.2253.

4.8 Manadoperoxide I (11)

Colorless amorphous solid; $[\alpha]_D^{25} -19.3$ (*c* 0.1, CHCl₃); ¹H NMR (C₆D₆, 500 MHz): δ_H 5.24 (1H, bs, H-11), 4.50 (1H, m, H-3), 4.19 (1H, q, 11 6.9 Hz, H-13), 3.34 (3H, s, 1-OMe), 3.20 (21, s, 6-OMe), 2.88 (1H, dd, *J* = 15.5, 9.3 Hz, H-2a), 2.50 (1H, 53 H-4), 2.14 (1H, dd, *J* = 15.5, 4.4 Hz, H-2b), 1.68 (1H, m, H-7a), 1.29 (1H, m, H-9a), 1.62 (1H, m, H-9b), 1.59 (1H, m, H-7b), 1.44 (1H, overlapped, H-5a), 1.38 (1H, m, H-8a), 1.36 (3H, bs, 12-Me), 1.33 (1H, m, H-8b), 1.17 (74, d, *J* = 6.9 Hz, Me-14), 0.88 (1H, overlapped, H-5b), 0.38 (3H, d, *J* = 7.0 Hz, 4-Me); ¹H NMR data for the minor epimer (only resonances differing from those of the major epimer are reported): δ_H 4.15 (59 q, *J* = 6.9 Hz, H-13), 1.19 (3H, d, *J* = 6.9 Hz, H-14), 1.69 (1H, m, H-9a), 1.58 (1H, m, H-9b); ¹³C NMR (C₆D₆, 125 7 Hz) δ_C 171.8 (s, C-1), 135.8 (s, C-12), 122.7 (d, C-11), 102.9 (s, C-6), 102.1 (s, C-10), 79.7 (d, C-3), 77.6 (d, C-13), 52.2 (q, 1-OMe), 48.8 (q, 6-OMe), 40.8 (t, C-9), 34.3 (t, C-5), 33.8 (C-7), 31.7 (t, C-2), 27.1 (d, C-4), 22.5 (t, C-8); 18.5 (q, C-14), 17.2 (q, 12-Me), 15.8 (q, 4-Me); (+) ESI-MS: *m/z* 359 [M + H]⁺, 381 [M + Na]⁺. HR-ESIMS (positive ions): found *m/z* 381.1892; C₁₈H₃₀NaO₇ requires *m/z* 381.1889.

4.9 Manadoperoxide J (12)

Colorless amorphous solid; $[\alpha]_D^{25} -11.9$ (*c* 0.1, CHCl₃); ¹H NMR (CDCl₃, 500 MHz): Table 1; ¹H NMR data for the minor epimer (only resonances differing from those of the major epimer are reported): δ_H 4.55 (1H, bs, H-11), 4.38 (1H, m, H-13), 1.78 (1H, overlapped, H-14a), 1.56 (1H, overlapped, H-14b), 1.00 (3H, t, *J* = 7.0 Hz, H-15); ¹³C NMR (CDCl₃, 125 MHz): Table 2; (+) ESI-MS *m/z* 97 and 409 (about 3:1 ratio) [M + H]⁺, 429, 431 (about 3:1 ratio) [M + Na]⁺.

HR-ESIMS found m/z 429.1663; $C_{19}H_{31}^{35}ClNaO_7$ requires m/z 429.1656.

4.10 Manadoperoxide K (13)

Colorless amorphous solid; $[\alpha]_D^{25}$ -11.5 (c 0.3, $CHCl_3$); 1H NMR ($CDCl_3$, 500 MHz): Table 1; ^{13}C NMR ($CDCl_3$, 125 MHz): Table 2; (+) ESI-MS: m/z 449 and 425 (about 3 : 1 ratio) $[M + H]^+$, 445 and 447 (about 3 : 1 ratio) $[M + Na]^+$. HR-ESIMS (positive ions): found m/z 445.1972; $C_{20}H_{35}^{35}ClNaO_7$ requires m/z 445.1969.

4.11 Peroxyplakoric ester C (14)

Colorless amorphous solid; $[\alpha]_D^{25}$ -3.0 (c 0.3, $CHCl_3$); 1H NMR ($CDCl_3$, 500 MHz): δ_H 6.77 (1H, dt, $J = 16.5, 6.5$ Hz, H-10), 6.18 (1H, d, $J = 16.5$ Hz, H-11), 4.19 (1H, m, H-3), 3.75 (3H, s, 1-OMe), 3.27 (3H, s, 6-OMe), 3.07 (1H, dq, $J = 9.5, 7.0$ Hz, H-2), 2.25 (2H, m, H₂-9), 2.24 (3H, s, Me-13), 2.10 (1H, m, H-4a), 1.75 (1H, m, H-5a), 1.69 (1H, m, H-7a), 1.65 (1H, m, H-5b), 1.53 (2H, t, H₂-8), 1.40 (1H, m, H-4b), 1.30 (1H, m, H-7b), 1.12 (3H, d, $J = 7.0$ Hz, 2-Me); ^{13}C NMR ($CDCl_3$, 125 MHz): δ_C 198.7 (s, C-12), 172.5 (s, C-1), 147.0 (d, C-10), 132.1 (d, C-11), 131.5 (s, C-6), 81.0 (d, C-3), 52.0 (q, 1-OMe), 48.8 (6-OMe), 41.6 (2-Me), 32.9 (t, C-7), 32.6 (t, C-9), 27.3 (q, C-13), 26.5 (t, C-5), 21.8 (t, C-8), 20.5 (t, C-4), 17.0 (q, 2-Me). (+) ESI-MS m/z 351 $[M + Na]^+$. HR-ESIMS (positive ions): found m/z 351.1790, $C_{17}H_{28}NaO_6$ requires m/z 351.1784.

4.12 Activity against *Trypanosoma brucei rhodesiense*

Minimum Essential Medium (50 μ L) supplemented with 25 mM HEPES, 1 g L^{-1} additional glucose, 1% MEM non-essential amino acids (100 \times), 0.2 mM 2-mercaptoethanol, 1 mM Na-pyruvate and 15% heat inactivated horse serum was added to each well of a 96-well microtitre plate. Serial drug dilutions of eleven 3-fold dilution steps covering a range from 100 to 0.002 μ g mL^{-1} were prepared. Then 10^4 bloodstream forms of STIB 900 strain of *T. b. rhodesiense* in 50 μ L was added to each well and the plate incubated at 37 $^{\circ}C$ under a 5% CO_2 atmosphere for 72 h. 10 μ L of a resazurin solution (12.5 mg resazurin dissolved in 100 mL double-distilled water) was then added to each well and incubation continued for a further 2–4 h. Then the plates were read in a Spectramax Gemini XS microplate fluorometer (Molecular Devices Cooperation, Sunnyvale, CA, USA) using an excitation wavelength of 536 nm and an emission wavelength of 588 nm. The IC_{50} values were calculated by linear regression (Huber 1993) from the sigmoidal dose inhibition curves using SoftmaxPro software (Molecular Devices Cooperation, Sunnyvale, CA, USA). Melarsoprol was the standard drug.

4.13 Activity against *Leishmania donovani*

Amastigotes of *L. donovani* (strain MHOM/ET/67/L82) grown in axenic culture at 37 $^{\circ}C$ in SM medium at pH 5.5, supplemented with 10% heat-inactivated fetal bovine serum under an atmosphere of 5% CO_2 in air. One hundred μ L of culture medium with 10^5 amastigotes from axenic culture with or

without a serial drug dilution were seeded in 96-well microtitre plates. Serial drug dilutions covering a range from 90 to 0.001 μ g mL^{-1} were prepared. After 72 h of incubation the plates were inspected under an inverted microscope to assure growth of the controls and sterile conditions. 10 μ L of a resazurin solution (12.5 mg resazurin dissolved in 100 mL double-distilled water) was then added to each well and the plates incubated for another 2 h. Then the plates were read in a Spectramax Gemini XS microplate fluorometer using an excitation wavelength of 536 nm and an emission wavelength of 588 nm. Data were analyzed using the software Softmax Pro. Decrease of fluorescence (*i.e.* inhibition) was expressed as percentage of the fluorescence of control cultures and plotted against the drug concentrations. The IC_{50} values were calculated from the sigmoidal inhibition curves. Miltefosine was used as a reference drug.

4.14 Cell cytotoxicity assay

HMEC-1 cells immortalized by SV 40 large T antigen was maintained in MCDB 131 medium supplemented with 10% fetal calf serum, 10 ng mL^{-1} of epidermal growth factor, 1 μ g mL^{-1} of hydrocortisone, 2 mM glutamine, 100 U mL^{-1} of penicillin, 100 mL^{-1} of streptomycin, and 20 mM Hepes buffer. Cells were treated with serial dilutions of test compounds and cell proliferation evaluated using the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay.

Conclusions

In the last decade, the urgent need for viable alternatives to the existing antiquated and unsatisfactory therapeutic options has stimulated a remarkable increase in the efforts towards finding new antitrypanosomal agents, often as a result of a series of public–private partnerships. A number of different classes of active compounds have been proposed and, among them, purine nitriles, thiosemicarbazones, quinolones, and chalcone–benzoxaborole appear to display better trypanocidal potential and better selectivity. Somewhat surprisingly, in spite of the incredible chemical diversity associated with the secondary metabolites elaborated by terrestrial plants and marine invertebrates, the contribution of natural products chemistry to the fight against trypanosomiasis has been until now modest. As matter of fact, the above reported classes of recently developed trypanocidal compounds include only synthetically produced molecules.

Data presented in this paper have disclosed that manadoperoxides constitute one of the most potent classes of natural products acting against sleeping sickness isolated to date and strongly suggest that endoperoxyketal-containing molecules are of promise for further development as lead structures against African trypanosomiasis. Moreover, the structure–activity relationships deduced have revealed that minor structural changes around the dioxane ring can dramatically affect the pharmacological activity, thus suggesting the interaction of this moiety with a specific target(s).

Since trypanosomiasis affects the poorest countries, the cost of the treatment is one of the main concerns in the design of new chemotherapeutics and, although effective, too expensive drugs

should be discarded. In this regard, the natural product state of manadoperoxides should not be considered as a drawback on the way to their development as antitrypanosomal drugs; indeed their relatively simple structures should guarantee the feasibility of a total synthesis. They are small, lipophilic and appear to have good pharmacokinetic properties, so they offer significant drug-likeness to become bioavailable lead compounds. A number of syntheses of the six-membered endoperoxyketal scaffold have been already published in the literature, very recently also by our group.¹⁵ These should allow the preparation of large libraries of compounds and the development of this new class of antitrypanosomal agents.

Acknowledgements

This work was supported by MIUR (PRIN2008, 20084MMXNM: Leads ad Attività Antimalarica di Origine Naturale, Ottimizzazione e Valutazione Biologica). MS and NMR spectra were registered by Authors at CSIAS, Centro Interdipartimentale di Analisi Strumentale, Faculty of Pharmacy, University of Naples.

References

- C. Burri, S. Nkunku, A. Merolle, T. Smith, J. Blum and R. Brun, *Lancet*, 2000, **355**, 1419–1425.
- E. M. Fèvre, B. V. Wissmann, S. C. Welburn and P. Lutumba, *PLoS Negl. Trop. Dis.*, 2008, **2**, e333.
- N. M. El-Sayed, P. J. Myler and D. C. Bartholomeu, *Science*, 2005, **309**, 409–415.
- M. P. Barrett, I. M. Vincent, R. J. S. Burchmore, A. J. N. Kazibwe and E. Matovu, *Fut. Microbiol.*, 2011, **6**, 1037–1047.
- P. G. Kennedy, *Ann. Neurol.*, 2008, **64**, 116–126.
- <http://www.dndi.org/portfolio/fexinidazole.html>
- E. Fattorusso, S. Parapini, C. Campagnuolo, N. Basilico, O. Tagliatalata-Scafati and D. Taramelli, *J. Antimicrob. Chemother.*, 2002, **50**, 883–888.
- G. Chianese, S. R. Yerbanga, L. Lucantoni, A. Habluetzel, N. Basilico, D. Taramelli, E. Fattorusso and O. Tagliatalata-Scafati, *J. Nat. Prod.*, 2010, **73**, 1448–1452.
- F. Scala, E. Fattorusso, M. Menna, O. Tagliatalata-Scafati, M. Tierney, M. Kaiser and D. Tasdemir, *Mar. Drugs*, 2010, **8**, 2162–2174.
- C. Campagnuolo, E. Fattorusso, A. Romano, O. Tagliatalata-Scafati, N. Basilico, S. Parapini and D. Taramelli, *Eur. J. Org. Chem.*, 2005, 5077–5083.
- E. Fattorusso, O. Tagliatalata-Scafati, A. Ianaro and M. Di Rosa, *Tetrahedron*, 2000, **56**, 7959–7967.
- C. Fattorusso, G. Campiani, B. Catalanotti, M. Persico, N. Basilico, S. Parapini, D. Taramelli, C. Campagnuolo, E. Fattorusso, A. Romano and O. Tagliatalata-Scafati, *J. Med. Chem.*, 2006, **49**, 7088–7094.
- O. Tagliatalata-Scafati, E. Fattorusso, A. Romano, F. Scala, V. Barone, P. Cimino, E. Stendardo, B. Catalanotti, M. Persico and C. Fattorusso, *Org. Biomol. Chem.*, 2010, **8**, 846–856.
- C. Fattorusso, M. Persico, N. Basilico, D. Taramelli, E. Fattorusso, F. Scala and O. Tagliatalata, *Bioorg. Med. Chem.*, 2011, **19**, 312–320.
- M. Persico, A. Quintavalla, F. Rondinelli, C. Trombini, M. Lombardo, C. Fattorusso, V. Azzarito, D. Taramelli, S. Parapini, Y. Corbett, G. Chianese, E. Fattorusso and O. Tagliatalata-Scafati, *J. Med. Chem.*, 2011, **54**, 8526–8540.
- In our paper cited in ref. 17 this sponge was tentatively attributed to *Plakortis* cfr. *simplex* on the basis of its morphological relationships with the *Plakortis simplex* species complex. However, on the basis of a comparison with the detailed morphological data later published (ref. 18), we consider conspecific the specimens analyzed both in this study and in our previous work (ref. 17) and attribute them to the species *Plakortis* cfr. *lita*.
- C. Fattorusso, M. Persico, B. Calcinai, C. Cerrano, S. Parapini, D. Taramelli, E. Novellino, A. Romano, F. Scala, E. Fattorusso and O. Tagliatalata-Scafati, *J. Nat. Prod.*, 2010, **73**, 1138–1145.
- M. Muricy, *J. Mar. Biol. Assoc. UK*, 2011, **91**, 303–319.
- M. Kobayashi, K. Kondo and I. Kitagawa, *Chem. Pharm. Bull.*, 1993, **41**, 1324–1326.
- F. Freire, J. M. Seco, E. Quinoa and R. Riguera, *J. Org. Chem.*, 2005, **70**, 3778–3790.
- K. W. L. Yong, L. K. Lambert, P. Y. Hayes, J. J. De Voss and M. J. Garson, *J. Nat. Prod.*, 2012, **75**, 351–360.
- C. A. Lipinski, F. Lombardo, B. W. Dominy and P. J. Feeney, *Adv. Drug Delivery Rev.*, 2001, **46**, 3–26.
- M. H. Kossuga, A. M. Nascimento, J. Q. Reimao, A. G. Tempone, N. N. Taniwaki, K. Veloso, A. G. Ferreira, B. C. Cavalcanti, C. Pessoa, M. O. Moraes, A. M. S. Mayer, E. Haidu and R. G. S. Berlink, *J. Nat. Prod.*, 2008, **71**, 334–339.
- H. Holla, M. Labaied, N. Phom, I. D. Jenkins, K. Stuart and R. J. Quinn, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 4793–4797.
- M. Kaiser, S. Wittlin, A. Nehrbass-Stuedli, Y. Dong, X. Wang, A. Hemphill, H. Matile, R. Brun and J. L. Vennerstrom, *Antimicrob. Agents Chemother.*, 2007, **51**, 2991–2993.
- A. Cavalli and M. L. Bolognesi, *J. Med. Chem.*, 2009, **52**, 7339–7359.
- R. L. Krauth-Siegel, H. Bauer and R. H. Schirmer, *Angew. Chem., Int. Ed.*, 2005, **44**, 690–715.
- Z. Qiao, Q. Wang, F. Zhang, Z. Wang, T. Bowling, B. Nare, R. T. Jacobs, J. Zhang, D. Ding, Y. Liu and H. Zhou, *J. Med. Chem.*, 2012, **55**, 3553–3557.
- J. P. Mallari, A. A. Shelat, T. O'Brien, C. R. Caffrey, A. Kosinski, M. Connelly, M. Harbut, D. Greenbaum, J. H. McKerrow and R. K. Guy, *J. Med. Chem.*, 2008, **51**, 545–552.
- J. P. Mallari, A. A. Shelat, A. Kosinski, C. R. Caffrey, M. Connelly, F. Zhu, J. H. McKerrow and R. K. Guy, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 2883–2885.
- G. Hiltensperger, N. G. Jones, S. Niedermeier, A. Stich, M. Kaiser, J. Jung, S. Puhl, A. Damme, H. Braunschweig, L. Meinel, M. Engstler and U. Holzgrabe, *J. Med. Chem.*, 2012, **55**, 2538–2548.
- I. V. Tetko, J. Gasteiger, R. Todeschini, A. Mauri, D. Livingstone, P. Ertl, V. A. Palyulin, E. V. Radchenko, N. S. Zefirov, A. S. Makarenko, V. Y. Tanchuk and V. V. Prokopenko, *J. Comput. Aided Mol. Des.*, 2005, **19**, 453–463.
- W. Huber and J. C. Koella, *Acta Trop.*, 1993, **55**, 257–261.

Manadoperoxides, a new class of potent antitrypanosomal agents of marine origin

ORIGINALITY REPORT

24%

SIMILARITY INDEX

15%

INTERNET SOURCES

21%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

1

eprints.uni-kiel.de

Internet Source

2%

2

"Handbook of Marine Natural Products",
Springer Nature America, Inc, 2012

Publication

1%

3

katsuokurabayashi.com

Internet Source

1%

4

Caterina Fattorusso, Giuseppe Campiani,
Bruno Catalanotti, Marco Persico et al.
"Endoperoxide Derivatives from Marine
Organisms: 1,2-Dioxanes of the Plakortin
Family as Novel Antimalarial Agents", Journal
of Medicinal Chemistry, 2006

Publication

1%

5

Submitted to The Hong Kong Polytechnic
University

Student Paper

1%

6

Giuseppina Chianese, Johanna Silber, Paolo
Luciano, Christian Merten et al. " Antiprotozoal

1%

Linear Furanosesterterpenoids from the Marine Sponge ", Journal of Natural Products, 2017

Publication

-
- | | | |
|----|---|-----|
| 7 | Submitted to Mahidol University
Student Paper | 1% |
| 8 | Caterina Fattorusso, Marco Persico, Nicoletta Basilico, Donatella Taramelli et al.
"Antimalarials based on the dioxane scaffold of plakortin. A concise synthesis and SAR studies", Bioorganic & Medicinal Chemistry, 2011
Publication | 1% |
| 9 | Angawi, Rihab F., Giorgio Bavestrello, Barbara Calcinaï, Henny Adeleida Dien, Giovanna Donnarumma, Maria Antonietta Tufano, Iole Paoletti, Elena Grimaldi, Giuseppina Chianese, Ernesto Fattorusso, and Orazio Tagliatela-Scafati. "Aurantioside J: a New Tetramic Acid Glycoside from Theonella swinhoei. Insights into the Antifungal Potential of Aurantiosides", Marine Drugs, 2011.
Publication | 1% |
| 10 | Fattorusso, Ernesto, Paolo Luciano, Adriana Romano, Orazio Tagliatela-Scafati, Giovanni Appendino, Marianna Borriello, and Caterina Fattorusso. "Stereostructure Assignment of Medium-Sized Rings through an | <1% |

NMR–Computational Combined Approach.
Application to the New Germacrane
Ketopelenolides C and D", Journal of Natural
Products, 2008.

Publication

11

L. Yang. "Biotransformation of α - and 6β -
santonin by fungus and plant cell cultures",
Journal of Asian Natural Products Research,
6/1/2006

Publication

12

ir.uiowa.edu

Internet Source

13

Pagani, A.. "Cannabioxepane, a novel
tetracyclic cannabinoid from hemp, *Cannabis
sativa* L.", Tetrahedron, 20110513

Publication

14

Giuseppina Chianese, Ernesto Fattorusso,
Orazio Taglialatela-Scafati, Giorgio Bavestrello
et al. "Desulfohaplosamate, a new phosphate-
containing steroid from *Dasychalina* sp., is a
selective cannabinoid CB2 receptor ligand",
Steroids, 2011

Publication

15

projectdisaster.com

Internet Source

16

Submitted to University of Queensland

Student Paper

<1%

<1%

<1%

<1%

<1%

<1%

17

Submitted to University College London

Student Paper

<1%

18

Persico, Marco, Silvia Parapini, Giuseppina Chianese, Caterina Fattorusso, Marco Lombardo, Luca Petrizza, Arianna Quintavalla, Francesca Rondinelli, Nicoletta Basilico, Donatella Taramelli, Claudio Trombini, Ernesto Fattorusso, and Orazio Taglialatela-Scafati. "Further Optimization of Plakortin Pharmacophore: Structurally Simple 4-Oxymethyl-1,2-Dioxanes with Promising Antimalarial Activity", *European Journal of Medicinal Chemistry*, 2013.

Publication

<1%

19

Alì Snene, Carmina Sirignano, Daniela Rigano, Carmen Formisano et al. "Antiproliferative metabolites from the Northern African endemic plant *Daucus virgatus* (Apiaceae)", *Phytochemistry*, 2017

Publication

<1%

20

Putra, Masteria Yunovilsa, Angela Ianaro, Elisabetta Panza, Giorgio Bavestrello, Carlo Cerrano, Ernesto Fattorusso, and Orazio Taglialatela-Scafati. "Sinulasulfoxide and sinulasulfone, sulfur-containing alkaloids from the Indonesian soft coral *Sinularia* sp.", *Tetrahedron Letters*, 2012.

<1%

21

Submitted to Higher Education Commission
Pakistan

Student Paper

<1%

22

Submitted to Universiti Malaysia Sabah

Student Paper

<1%

23

Fattorusso, E.. "Coelodiol and coeloic acid, ent-
isocopalane diterpenes from the Indonesian
sponge Coelocarteria cfr. singaporensis",
Tetrahedron Letters, 20060327

Publication

<1%

24

Submitted to Universita degli Studi di Torino

Student Paper

<1%

25

Williamson, R.. "Taveuniamides: new
chlorinated toxins from a mixed assemblage of
marine cyanobacteria", Tetrahedron, 20040809

Publication

<1%

26

www.cambridge.org

Internet Source

<1%

27

Taglialatela-Scafati, Orazio, Ernesto
Fattorusso, Adriana Romano, Fernando Scala,
Vincenzo Barone, Paola Cimino, Emiliano
Stendardo, Bruno Catalanotti, Marco Persico,
and Caterina Fattorusso. "Insight into the
mechanism of action of plakortins, simple 1,2-
dioxane antimalarials", Organic & Biomolecular

<1%

Chemistry, 2010.

Publication

28

"Contents list", Organic & Biomolecular Chemistry, 2012.

Publication

<1%

29

Abad, A.. "A unified synthetic approach to trachylobane-, beyerane-, atisane- and kaurane-type diterpenes", Tetrahedron, 20060403

Publication

<1%

30

Submitted to University of Scranton

Student Paper

<1%

31

europemc.org

Internet Source

<1%

32

www.rsc.org

Internet Source

<1%

33

Fontana, A.. "Structure and absolute stereochemistry of stolonoxide A, a novel cyclic peroxide from the marine tunicate *Stonica socialis*", Tetrahedron Letters, 20000115

Publication

<1%

34

Submitted to Misr International University

Student Paper

<1%

35

Submitted to Lander University

Student Paper

<1%

36

academic.oup.com

Internet Source

<1%

37

Concetta Imperatore, Marco Persico, Maria Senese, Anna Aiello et al. "Exploring the antimalarial potential of the methoxy-thiazinoquinone scaffold: Identification of a new lead candidate", *Bioorganic Chemistry*, 2019

Publication

<1%

38

K. C. Nicolaou, Daniel Schlawe, David W. Kim, Deborah A. Longbottom et al. "Total Synthesis of Halipeptins: Isolation of Halipeptin D and Synthesis of Oxazoline Halipeptin Analogues", *Chemistry - A European Journal*, 2005

Publication

<1%

39

Patrizia Ciminiello, Carmela Dell'Aversano, Ernesto Fattorusso, Silvana Magno, Maurizio Pansini. "Chemistry of Verongida Sponges. 10. Secondary Metabolite Composition of the Caribbean Sponge", *Journal of Natural Products*, 2000

Publication

<1%

40

Kamaljit Singh, Tavleen Kaur. "Pyrimidine-based antimalarials: design strategies and antiplasmodial effects", *MedChemComm*, 2016

Publication

<1%

Putra, M.Y.. "Polyhydroxylated sterols from the

41 Indonesian soft coral *Sinularia* sp. and their effect on farnesoid X-activated receptor", *Steroids*, 201204 $<1\%$
Publication

42 tutcris.tut.fi $<1\%$
Internet Source

43 gene-quantification.net $<1\%$
Internet Source

44 beilstein-journals.org $<1\%$
Internet Source

45 Nguyen Huu Tung, Chau Van Minh, Phan Van Kiem, Hoang Thanh Huong et al. "Chemical components from the Vietnamese soft coral *Lobophytum* sp.", *Archives of Pharmacal Research*, 2010 $<1\%$
Publication

46 www.jove.com $<1\%$
Internet Source

47 who.int $<1\%$
Internet Source

48 res.mdpi.com $<1\%$
Internet Source

49 R. Luise Krauth-Siegel, Holger Bauer, R. Heiner Schirmer. "Dithiol Proteins as Guardians of the Intracellular Redox Milieu in Parasites: $<1\%$

Old and New Drug Targets in Trypanosomes and Malaria-Causing Plasmodia", *Angewandte Chemie International Edition*, 2005

Publication

50

Appleton, D.R.. "Novel tryptophan-derived dipeptides and bioactive metabolites from the sea hare *Aplysia dactylomela*", *Tetrahedron*, 20011217

<1%

Publication

51

repository.up.ac.za

Internet Source

<1%

52

real.mtak.hu

Internet Source

<1%

53

Submitted to Chungnam National University

Student Paper

<1%

54

Valentina Sepe, Raffaella Ummarino, Maria Valeria D'Auria, Maria Giovanna Chini et al. "Conicasterol E, a Small Heterodimer Partner Sparing Farnesoid X Receptor Modulator Endowed with a Pregnane X Receptor Agonistic Activity, from the Marine Sponge ", *Journal of Medicinal Chemistry*, 2011

<1%

Publication

55

Liu, Dong-Ze, and Ji-Kai Liu. "Peroxy natural products", *Natural Products and Bioprospecting*, 2013.

<1%

Publication

56	theses.gla.ac.uk Internet Source	<1%
57	www.vanguardngr.com Internet Source	<1%
58	msbmb.org Internet Source	<1%
59	Clark, J.S.. "Synthesis of a lactone natural product found in Greek tobacco", Tetrahedron Letters, 20070402 Publication	<1%
60	Submitted to Nottingham Trent University Student Paper	<1%
61	ecc.isc.gov.ir Internet Source	<1%
62	Marcos, I.S.. "Synthetic studies towards picrasane quassinoids", Tetrahedron, 20070305 Publication	<1%
63	Carmen Festa, Simona De Marino, Maria Valeria D'Auria, Eric Deharo et al. "Gracilioethers E–J, new oxygenated polyketides from the marine sponge Plakinastrella mamillaris", Tetrahedron, 2012 Publication	<1%
64	Gemma, S.. "Synthetic studies toward 1,2-dioxanes as precursors of potential	<1%

endoperoxide-containing antimalarials",
Tetrahedron Letters, 20091014

Publication

65

www.freepatentsonline.com

Internet Source

<1%

66

www.oatext.com

Internet Source

<1%

67

pubs.rsc.org

Internet Source

<1%

68

www.mdpi.com

Internet Source

<1%

69

Sinisi, Annamaria, Barbara Calcinai, Carlo Cerrano, Henny A. Dien, Angela Zampella, Claudio D'Amore, Barbara Renga, Stefano Fiorucci, and Orazio Taglialatela-Scafati. "Isoswinholide B and swinholide K, potently cytotoxic dimeric macrolides from *Theonella swinhoei*", *Bioorganic & Medicinal Chemistry*, 2013.

Publication

<1%

70

Banwell, Martin G., Mark J. Coster, Natasha L. Hungerford, Mary J. Garson, Stephen Su, Andrew C. Kotze, and Murray H. G. Munro. "3,4- β -Linked bis(piperidines) related to the haliclonyclamine class of marine alkaloids: synthesis using crossed-aldol chemistry and

<1%

preliminary biological evaluations", Organic & Biomolecular Chemistry, 2012.

Publication

71

Peter E. Cockram, Terry K. Smith. "Active Natural Product Scaffolds against Trypanosomatid Parasites: A Review", Journal of Natural Products, 2018

Publication

<1%

72

publish.csiro.au

Internet Source

<1%

73

researchspace.auckland.ac.nz

Internet Source

<1%

74

patentscope.wipo.int

Internet Source

<1%

75

Submitted to Macau University of Science and Technology

Student Paper

<1%

76

Fattorusso, E.. "Loboanthamine, a new zoanthamine-type alkaloid from the Indonesian soft coral Lobophytum sp.", Tetrahedron Letters, 20080331

Publication

<1%

77

Liang, Lin-Fu, Li-Xin Gao, Jia Li, Orazio Tagliatela-Scafati, and Yue-Wei Guo. "Cembrane diterpenoids from the soft coral Sarcophyton trocheliophorum Marenzeller as a

<1%

new class of PTP1B inhibitors", *Bioorganic & Medicinal Chemistry*, 2013.

Publication

78

Ernesto Fattorusso, Adriana Romano, Orazio Taglialatela-Scafati, M. Janib Achmad, Giorgio Bavestrello, Carlo Cerrano. "Xenimanadins A–D, a family of xenicane diterpenoids from the Indonesian soft coral *Xenia* sp.", *Tetrahedron*, 2008

<1%

Publication

79

de Champdore, M.. "In-water reactivity of nucleosides and nucleotides: one-step preparation and biological evaluation of novel ferrocenyl-derivatives", *Tetrahedron*, 20040726

<1%

Publication

80

Adio, A.M.. "Sesquiterpenoid constituents of the liverwort *Marsupella aquatica*", *Tetrahedron: Asymmetry*, 20070611

<1%

Publication

81

Fattorusso, E.. "Xenimanadins A-D, a family of xenicane diterpenoids from the Indonesian soft coral *Xenia* sp.", *Tetrahedron*, 20080331

<1%

Publication

82

Gribble, Gordon. "Biological Activity of Recently Discovered Halogenated Marine Natural Products", *Marine Drugs*, 2015.

<1%

Publication

83

Jung Oh, Buyng Hwang, Ok-Hwa Kang, Dong-Yeul Kwon, Jung-Rae Rho. "New Constituents from the Korean Sponge *Plakortis simplex*", *Marine Drugs*, 2013

Publication

<1%

84

siba-ese.unile.it

Internet Source

<1%

85

D. P. Sonawane, M. Persico, Y. Corbett, G. Chianese et al. "New antimalarial 3-methoxy-1,2-dioxanes: optimization of cellular pharmacokinetics and pharmacodynamics properties by incorporation of amino and N-heterocyclic moieties at C4", *RSC Advances*, 2015

Publication

<1%

86

R.-T. Li. "Chemical constituents from *Schisandra plena*", *Journal of Asian Natural Products Research*, 12/1/2005

Publication

<1%

87

Festa, Carmen, Simona De Marino, Maria Valeria D'Auria, Orazio Tagliatela-Scafati, Eric Deharo, Sylvain Petek, and Angela Zampella. "New antimalarial polyketide endoperoxides from the marine sponge *Plakinastrella mamillaris* collected at Fiji Islands", *Tetrahedron*, 2013.

Publication

<1%

88

Okasaka, M.. "Terpenoids from *Juniperus polycarpus* var. *seravschanica*",
Phytochemistry, 200612

Publication

<1%

89

Christophe Daepfen, Marcel Kaiser, Markus Neuburger, Karl Gademann. "Preparation of Antimalarial Endoperoxides by a Formal [2 + 2 + 2] Cycloaddition", Organic Letters, 2015

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On