

Serum nitric oxide and pediatric sepsis outcomes

by Jeanette Manoppo 6

Submission date: 26-Aug-2019 11:53AM (UTC+0700)

Submission ID: 1163476629

File name: 6._Serum_Nitric_Oxide_and_Pediatric_Sepsis_Outcomes.pdf (109.07K)

Word count: 3843

Character count: 21561

Serum nitric oxide and pediatric sepsis outcomes

Ronald Chandra, Jose M. Mandei, Jeanette I. Ch. Manoppo,
Rocky Wilar, Ari L. Runtunuwu, Phey Liana

Abstract

Background Sepsis is the complex pathophysiologic responses of the host against septic infection. Sepsis can cause severe conditions such as septic shock and multiple organ failure. Although we have a better understanding of the molecular basis of sepsis as well as aggressive therapy, the mortality rate remains high, between 20-80%. Nitric oxide (NO) is one of the mediators associated with cardiovascular failure, apoptosis and organ dysfunction in sepsis.

Objective To evaluate for a possible correlation between NO levels and outcomes in pediatric sepsis.

Methods A prospective cohort study was conducted at the pediatric intensive care unit (PICU) of Prof. Dr. R.D. Kandou General Hospital in Manado, from June to November 2012. Forty children aged one month to five year old, fulfilled the *International Pediatrics Sepsis Consensus Conference 2005* criteria were recruited. Nitrite oxide metabolites (nitrite and nitrate) levels were measured using a calorimetric assay kit (Cayman®, Catalog No.780001) from venous blood specimens collected at admission. All patients received antibiotics empirically within an hour of the diagnosis. Outcomes of patients recorded were survivor or died, and length of stay in PICU.

Results Mann-Whitney U test revealed a significant difference between median serum NO levels in survivors and those who died (18.60 vs. 36.50 $\mu\text{M/L}$, respectively; $P=0.016$).

Conclusion Serum NO concentration is higher in those who died than in survivors of pediatric sepsis. Specific NO inhibition may be beneficial in decreasing morbidity and mortality in this condition. [Paediatr Indones. 2014;54:213-8].

Keywords: nitric oxide, sepsis, mortality, pediatric

Sepsis is one of leading causes of admission to pediatric intensive care units (PICUs) in the world with significant morbidity and mortality in patients and an elevated cost for society.^{1,2} In United States, sepsis and septic shock are major causes of morbidity and mortality, with an estimated 42,000 cases per year and an associated mortality of 10,3% with more than 4,300 cases.²⁻⁴ However, in developing countries, the mortality rate of sepsis in children from PICU is higher than 50%.⁵ The incidence of sepsis was reported to be highest in children less than 1 year of age (5.16/1,000), and decreased to 0.2/1,000 in children 10-14 years of ages.²

The sepsis syndrome is characterized by alterations in vascular tone and, in the most severe forms of septic shock, the peripheral vasculature is markedly refractory to the vasoconstrictive effects of alpha-1-adrenergic receptor agonist agents.⁶ The prevailing theory is that sepsis represents an uncontrolled-inflammatory response with loss of the normal homeostatic balance between systemic inflammation and the anti-inflammatory re-

³⁴ From the Department of Child Health, Sam Ratulangi University Medical School, Manado¹ and ² Clinical Pathology of Sriwijaya University Medical School, Palembang², Indonesia.

Reprint requests to: Ronald Chandra, Departement of Child Health, Sam Ratulangi University Medical School, Jalan Raya Tanawangko no.56, Malalayang, Manado, Indonesia. Tel +62-431-821562, Fax +61-431-859091. E-mail: ronald_chandr@yahoo.co.id.

sponse.^{7,8} Cytokines and mediators that are released may cause collapse of circulation, pan-endothelial vasodilation,¹⁵ trauma, and increased of vascular permeability,⁹ leading to septic shock and multiple organ failure.⁷ Recently, it has been proposed that impaired cellular oxygen utilization, termed cytopathic hypoxia, rather than inadequate oxygen delivery, may play an important role in the development of multiple organ dysfunction syndrome (MODS).^{7,10} It may lead to suggest that, in early sepsis is associated with microcirculatory disturbances, in late course of disease the problem seems to be related to mitochondrial dysfunction. Nitric oxide (NO) is known to be mediator in microcirculation and cytopathic hypoxia.^{7,11,12}

Nitric oxide, a potent endogenous vasodilator produced by various mammalian cells, has been postulated to play a role in the hemodynamic derangements of sepsis syndrome.^{6,13,14} The inducible isoform of NO is responsible for increased NO output after induction by lipopolysaccharide and various pro-inflammatory cytokines [i.e., tumor necrosis factor-alpha (TNF- α), interleukin-1 beta (IL-1 β), and interferon gamma (IFN- γ)],¹⁴⁻¹⁸ which are of pathogenic importance during the sepsis syndrome.^{6,19} Although NO is known to have a role in sepsis, its action has not been elucidated by recent studies. Some studies reported a positive correlation between NO metabolites and sepsis,²⁰ septic shock,^{6,21} organ failure,²¹ severity of inflammation,⁶ and mortality.²² However, others reported no correlation between NO metabolites and sepsis,²³ severity of sepsis,²⁰ or mortality.^{20,24} We hypothesized NO concentration be higher in sepsis those who died than survivors. This study was undertaken to evaluate a possible correlation between NO levels and outcomes in pediatric sepsis.

Methods

A prospective cohort study was conducted at the PICU, Prof. Dr. R.D. Kandou General Hospital in Manado, from June to November 2012. The study protocol was approved by the Ethics Committee of the hospital. All patients' parents provided written informed consent following a full explanation of the study and reading of the information sheet. Infants and children, aged one month to five years, who fulfilled the *International Pediatrics Sepsis Conference Consensus criteria* (2005)³

were consecutively recruited. We excluded infants and children with diagnoses of severe dehydration, severe malnutrition, obesity, immunodeficiency disorders, trauma, scald burns, acute pancreatitis, and patients with a history of liver and kidney diseases in the 6 months prior to the study.

At time of admission, subjects' blood specimens were drawn in order to evaluate complete blood counts, urea and creatinine levels, blood cultures, and NO metabolites (nitrite and nitrate) levels. Clinical information was also recorded (age, gender, nutritional status and diagnosis) at that time. All subjects were treated with early goal-directed therapy. Combination antibiotics comprising of the 3rd generation cephalosporines, such as ceftriaxone or cefotaxime, and gentamicin, were given empirically within an hour of the diagnosis. The antibiotic regimen was appropriately changed after blood culture and sensitivity test results became available. During treatment, all subjects were followed-up daily until patients either recovered enough to be moved to the pediatric ward or died. Patients were considered to have recovered from sepsis if they were free from a systemic inflammation response for two days. Outcomes of patients were recorded (survivor or died and length of stay in PICU). None of the children were treated with inhaled NO or nitrovasodilators.

Venous blood specimens were collected at admission and immediately centrifuged at 1,500 rpm for 15 minutes. Serum samples were then removed and stored at -30 degrees Celsius until they were used for NO metabolite assays. Nitrate and nitrite measurements were made using the nitrate/nitrite colorimetric assay kit (Cayman®, catalog No. 780001). Two hundred μL of assay buffer and up to 80 μL of sample dilutions (diluted sample 1:1 with distilled water) were added to the wells. Ten μL of enzyme cofactor mixture, followed by 10 μL nitrate reductase mixture were added to each well. Plates were then covered and incubated for 1 hour at room temperature. We then added 50 μL of Griess Reagent R1 followed immediately by 50 μL of Griess Reagent R2 to the wells. Plates were incubated for 10 minutes at room temperatures to develop the color. Absorbance was read at 550 nm using a plate reader. Serum NO metabolites were plotted on the nitrate standard curve. The detection limit for this assay was 2.5 μM . Nitric oxide metabolites are stable and most eliminated by

the kidney. As such, indexing of NO metabolites to serum creatinine concentrations is considered to be representative of endogenous NO.

Statistical analysis consisted of descriptive and comparative analyses. Normality of data was analyzed by Kolmogorov-Smirnov test. Descriptive analyses were used in order to analyze the characteristics, clinical and laboratory findings, diagnoses as well as antibiotic use, and were reported in a distribution table. Parametric data was expressed as mean, standard deviation (SD), while non-parametric data was expressed as median, and minimum-maximum values.

We used the Mann-Whitney rank sum test to compared the ratio of serum NO metabolites to serum creatinine concentrations in pediatric sepsis between survivors and those who died. Data were considered to be statistically significant for $P < 0.05$. All analyses were performed using SPSS version 19.0.

Results

The final PICU dataset was comprised of 40 children with a median age of 8.5 (range 2 - 70) months, of whom 22 were boys. Thirty-three children were categorized in the under 24 months' old grouping. Subjects' nutritional status were as follows: 28 well-nourished, 11 undernourished, and 1 overweight. Two children had culture-positive sepsis, *Streptococcus pneumoniae* and *Enterobacter aerogenes*, respectively. The sources of sepsis included bronchopneumonia (33/40), meningococcal meningitis (11/40) and acute diarrhea (11/40). Thirty-one children were treated with a combination of cefotaxime and gentamycin. During treatment, 31 children survived and were allowed to move to pediatrics ward, while 9 children died. The median length of PICU stay was 5 (range 1-15) days (Table 1).

Table 1. Baseline characteristics of subjects

Characteristics	Overall n=40	Survived n=31	Died n=9
Gender			
Male, n	22	15	7
Median age (range), months	8.5 (2 - 70)**	10.11 (7.2)*	8 (2 - 26)**
1 to < 24 months, n	35	25	9
24 to < 72 months, n	5	5	0
Nutritional status			
Overweight, n	1	1	0
Well-nourished, n	28	22	6
Undernourished, n	11	8	3
Sources of sepsis, n			
Bronchopneumonia, n	33	26	7
Meningococcal meningitis, n	11	7	4
Acute diarrhea, n	11	7	4
Median creatinine (range), mg/dL	0.65 (0.2 - 2.5)**	0.7 (0.3 - 2.5)**	0.61 (0.28)*
Blood cultures, n			
<i>S. pneumoniae</i> , n	1	1	0
<i>E. aerogenes</i> , n	1	1	0
Antibiotics			
Cefotaxime+ gentamycin, n	31	23	8
Ceftriaxone+ gentamycin, n	9	8	1
Median length of PICU stay (range), days	5 (1 - 15)**	5 (2 - 15)**	1 (1 - 12)**

Note: * parametric data: mean (SD);** non-parametric data: median (range)

Table 2. Ratio of serum NO metabolites to serum creatinine concentration in survivors and those who died with pediatric sepsis.

Variable	Survived	Died	P value
Median ratio of serum NO to serum creatinine (range), $\mu\text{M/L}$	18.60 (4.20-59.30)	36.50 (12.90-93.00)	0.016

Mann-Whitney U test

The Mann-Whitney rank sum test analysis revealed that the ratio of serum NO metabolites to serum creatinine concentrations in those who died was higher than in survivors within pediatric sepsis (Table 2).

Discussion

Nitric oxide metabolites concentrations are influenced by their synthesis and elimination rates. As NO metabolites are excreted predominantly via the kidneys, differences in serum NO concentrations may occur as a result of renal dysfunction.^{6,25,26} Renal dysfunction in sicker children may be a consequence of increased disease severity.⁶ We attempted to correct any confounding effects of renal dysfunction in our patients by indexing serum concentrations of NO metabolites to serum creatinine concentrations to be a better proxy for the level of endogenous NO.

Nitric oxide has an important role in microvascular regulation.²⁷ Generated in physiologic amounts, NO has clearly been shown to have a beneficial role in organ perfusion by acting as a regulator of vascular tone and blood flow distribution. In basal amounts, NO is also considered to be important in immune-mediated disease by protecting various tissue against injury by endotoxin-induced inflammation. However, inducible nitric oxide synthetase is generated in response to inflammatory stimuli, and is able to produce vastly increased amounts of NO. This NO relaxes vascular tone, causes vascular hyporeactivity against vasopressor agents, depresses myocardium, and can lead to multiple organ dysfunction and death.²⁵⁻²⁹ Nitric oxide can also react with free oxygen and hydroxy radicals to form toxic peroxynitrite radicals which can induce damage to endothelial cells.¹⁵

In our study, we found that the ratio of serum NO metabolites to serum creatinine concentrations in those who died was significantly higher than in survivors. This finding was consistent with the role of endogenous NO overproduction in sepsis pathogenesis. Our findings were also in agreement with other study which reported that NO concentrations in pediatric sepsis were higher in non-survivors than in survivors and could be used as predictor of mortality.³⁰ In addition, a study showed that NO concentrations in non-survivors were higher than in survivors. The difference was not statistically

significant on the first day of examination, but it was significant on the 2nd and 3rd days of examination.²¹ A study on adults reported that serial NO concentrations were higher in sepsis non-survivors than in survivors (P=0.005) and was related to the severity of disease.²⁷ However, other studies reported no significant differences in NO concentration between survivors and non-survivors.^{25,29} All these other studies had different subject characteristics from our study, though most studies did not index the NO metabolites to serum creatinine concentrations in order to correct for renal dysfunction.

Interestingly, all those who died in this study had septic shock during the course of disease (data not shown). Mortality due to cardiovascular failure caused by septic shock may occur in the initial stages of sepsis. Nitric oxide is also known as a potent vasodilator and can decrease cardiac contractility.²¹ Our findings are consistent with others reports, such as Galley *et al*.³¹ who reported that NO concentration was elevated in non-survivors due to shock septic and decreased over the study period only in patient who survived, and other studies^{6,28} which reported that the ratio of serum NO metabolites to serum creatinine concentration was higher in the hypotensive sepsis group than in the non-hypotensive group (P<0.05). Nitric oxide concentrations also increased in sepsis patients who needed vasopressor agents (dopamine, dobutamine, or nor-epinephrine). They concluded that NO was associated with cardiovascular failure.^{6,21,28,32} An animal study reported that NO and peroxynitrite were involved in cytopathic hypoxia. Microvascular dysfunction and cytopathic hypoxia are reported to be good predictors for mortality.³³

Baines *et al*. were also in agreement with our findings in that they found a relationship between NO and mortality risk in children with meningococemia by calculating PRISM scores (Spearman's rho=0.72; P=0.0003).²⁶ Mitaka *et al*. reported that NO concentrations were related to APACHE II and SOFA scores in adults,²⁵ while Groeneveld *et al*. showed relationships between NO concentrations and TNF- α (Spearman's rho=0.69; P< 0.0001) or IL-6 (Spearman's rho=0.44; P<0.0001) in sepsis. Nitric oxide is reportedly increased when excess of cytokine pro-inflammation is produced during the first phase of sepsis but tends to decrease when cytokine anti-inflammation is produced during the secondary phase.³⁴ Doughty *et al*. reported that

children with elevated IL-6 concentration had higher NO concentrations than those with undetected IL-6 concentration in sepsis, but both were still higher than controls ($P < 0.005$).³⁵ Interleukin-6 concentration has been reported to be associated with mortality in pediatric sepsis.³⁶

Exogenous NO influences the endogenous NO levels.³⁷ We did not calculate the effect of oral intake before admission to PICU since our subjects were categorized as critically ill and did not receive exogenous NO during treatment (dietary nitrite, inhaled NO or nitrovasodilators). Our findings were consistent with another study which reported that oral intake contributed little NO metabolites to critically ill patients admitted to the hospital.³⁷

In addition to nitrite and nitrate as NO metabolites, NO also reacts with reactive oxygen species (ROS) to form reactive nitroso species (RNS), such as nitrosotyrosine (NT) and nitrosothiol.³⁸ A study reported that some septic shock patients had NO concentrations within normal limits, but higher NT concentrations.³⁸ Wong *et al.* also reported that variable NO concentrations in septic children occurred because the biological effects of NO are short lived and are likely to occur at local tissue and intracellular levels.⁶ All these possibilities may explain why some of our septic patients who died had lower NO concentrations. Because of the NO concentration variability in our patients, we should not use NO concentration as a single predictor of mortality in pediatric septic patients.

Limitations of our study are the single blood examination of NO, lack of measurements of other NO bioreactives, such as NT and nitrosothiol, and no calculation of the synthesis effect of NO. Further study is needed with serial examinations of NO metabolites (nitrate, nitrite, NT, and nitrosothiol) in order to more accurately evaluate the role of NO as a predictor of mortality. In conclusion, serum NO level is higher in those who died than in survivors pediatric sepsis. Specific NO inhibition may be beneficial in decreasing morbidity and mortality in this condition.

References

1. Jaramillo-Bustamante JC, Marin-Agudelo A, Fernandez-Laverde M, Bareno-Silva J. Epidemiology of sepsis in pediatric intensive care units: first Colombian multicenter study. *Pediatr Crit Care Med.* 2012;13:501-8
2. Thomas NJ, Tamburro RF, Hall MW, Rajasekaran S, Venglarcik JS. Bacterial sepsis and mechanisms of microbial pathogenesis. In: Nichols DG, Ackerman AD, Argent AC, editors. *Rogers' Textbook of Pediatric Intensive Care.* 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2008. p. 1300-27.
3. Goldstein B, Giroir B, Randolph A, International Consensus Conference on Pediatric Sepsis. International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med.* 2005;6:2-8.
4. Pavare J, Grope I, Gardovska D. Prevalence of systematic inflammatory response syndrome (SIRS) in hospitalized children: a point prevalence study. *BMC Pediatr.* 2009;9:25.
5. Khan MR, Maheshwari PK, Masood K, Qamar FN, Haque A. Epidemiology and outcome of sepsis in a tertiary care PICU of Pakistan. *Indian J Pediatr.* 2012;79:1454-8.
6. Wong HR, Carcillo JA, Burckart G, Shah N, Janosky JE. Increased serum nitrite and nitrate concentrations in children with the sepsis syndrome. *Crit Care Med.* 1995;23:835-42.
7. von-Dessauer B, Bongain J, Molina V, Quilodrán J, Castillo R, Rodrigo R. Oxidative stress as a novel target in pediatric sepsis management. *J Crit Care.* 2011;26:103e1-7.
8. Weighardt H, Holzmann B. Role of Toll-like receptor responses for sepsis pathogenesis. *Immunobiology.* 2007;212:715-22.
9. Brill RJ, Goldstein B. Pediatric sepsis definitions: Past, present and future. *Pediatr Crit Care Med.* 2005;6:6-8.
10. Peters K, Unger RE, Brunner J, Kirkpatrick CJ. Molecular basis of endothelial dysfunction in sepsis. *Cardiovasc Res.* 2003;60:49-57.
11. Azevedo LCP. Mitochondrial dysfunction during sepsis. *Endocr Metab Immune Disord Drug targets.* 2010;10:214-23.
12. Victor VM, Esplugues JV, Hernandez-Mijares A, Rocha M. Oxidative stress and mitochondrial dysfunction in sepsis: a potential therapy with mitochondria-targeted antioxidants. *Infect Disord Drug Targets.* 2009;9:376-89.
13. Cena J, Chow AK, Schulz R. Nitric oxide, peroxynitrite and matrix metalloproteinases: insight into the pathogenesis of sepsis. In: Tota B, Trimmer B, editors. *Nitric Oxide.* 1st ed. Oxford: Elsevier; 2007. p. 367-95.
14. Victor VM, Rocha M, De la Fuente M. Immune cells: free radicals and antioxidants in sepsis. *Int Immunopharmacol.* 2004;4:327-47.
15. Symeonides S, Balk RA. Nitric oxide in the pathogenesis of sepsis. *Infect Dis Clin North Am.* 1999;13:449-63.
16. Gross SS, Wolin MS. Nitric oxide: pathophysiological

- mechanisms. *Ann Rev Physiol.* 1995;57:737-69.
17. Wearden ME. Nitric oxide synthase inhibitors for septic shock. *Sem Pediatr Infect Dis.* 2001;12:42-5.
 18. Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev.* 2007;87:315-424.
 19. Suchner U, Heyland DK, Peter K. Immune-modulatory actions of arginine in the critically ill. *Br J Nutr.* 2002;87:121-32.
 20. Duke T, South M, Stewart A. Activation of L-arginine nitric oxide pathway in severe sepsis. *Arch Dis Child.* 1997;76:203-9.
 21. Doughty L, Carcillo JA, Kaplan S, Janosky J. Plasma nitrite and nitrate concentrations and multiple organ failure in pediatric sepsis. *Crit Care Med.* 1998;26:157-62.
 22. Barthlen W, Stadler J, Lehn NL, Miethke T, Bartels H, Siewert JR. Serum levels of end products of nitric oxide synthesis correlate positively with tumor necrosis factor alpha and negatively with body temperature in patients with postoperative abdominal sepsis. *Shock.* 1994;2:398-401.
 23. Terregino CA, Lopez BL, Karras DJ, Killian AJ, Arnold GK. Endogenous mediators in emergency department patients with presumed sepsis: are levels associated with progression to severe sepsis and death? *Ann Emerg Med.* 2000;35:26-34.
 24. Adamik B, Kubler-Keilb J, Golebiowska B, Gamian A, Kubler A. Effect of sepsis and cardiac surgery with cardiopulmonary bypass on plasma level of nitric oxide metabolites, neopterin, and procalcitonin: correlation with mortality and postoperative complications. *Intensive Care Med.* 2000;26:1259-67.
 25. Mitaka C, Hirata Y, Yokoyama K, Wakimoto H, Hirokawa M, Nosaka T, et al. Relationships of circulating nitrite/nitrate levels to severity and multiple organ dysfunction syndromes in systemic inflammatory response syndrome. *Shock.* 2003;19:305-9.
 26. Baines PB, Stanford S, Bishop-Bailey D, Sills JA, Thomson APJ, Mitchell JA, Fear SC, Hart CA, Petros AJ. Nitrite oxide production in meningococcal disease is directly related to disease severity. *Crit Care Med.* 1999;27:1187-90.
 27. Ho JT, Chapman MJ, O'Connor S, Lam S, Edwards J, Ludbrook G, Lewis JG, et al. Characteristics of plasma NOx levels in severe sepsis: high interindividual variability and correlation with illness severity, but lack of correlation with cortisol levels. *Clin Endocrinol.* 2010;73:413-20.
 28. Wong HR, Carcillo JA, Burckart G, Kaplan SS. Nitric oxide production in critically ill patients. *Arch Dis Child.* 1996;74:482-9.
 29. Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. *Analysis of nitrate, nitrite, and [¹⁵N] nitrate in biological fluids.* *Anal Biochem.* 1989;126:131-8.
 30. El-Gawad TAA, El-Sahrigy SAF, Abdel-Rahman AMO, Ghaffar EA, El-Rasheed EA. Plasma levels of nitric oxide and carbon monoxide in critically ill children with septic syndrome. *J Med Sci.* 2007;7:769-75.
 31. Galley HF, Webster NR. Plasma nitrite is increased in patients who die from sepsis. *Intensive Care Med.* 1996;22:S327.
 32. Fortin CF, McDonald PP, Fulop T, Lesur O. Sepsis, leukocytes, and nitric oxide (NO): an intricate affair. *Shock.* 2010;33:344-52.
 33. Collaborative group for the study of sepsis in PICUs in Shanghai area. Hospital epidemiology, management and outcome of pediatric sepsis and severe sepsis in 4 PICUs in Shanghai. *Zhonghua Er Ke Za Zhi.* 2012;50:172-7.
 34. Groeneveld PH, Kwappenberg KM, Langermans JA, Nibbering PH, Curtis L. Relation between pro- and anti-inflammatory cytokines and the production of nitric oxide (NO) in severe sepsis. *Cytokine.* 1997;9:138-42.
 35. Doughty LA, Kaplan SS, Carcillo JA. Inflammatory cytokines and nitric oxide responses in pediatric sepsis and organ failure. *Crit Care Med.* 1996;24:1137-43.
 36. Survillan JS, Kilpatrick L, Costarino AT Jr, Lee SC, Harris MC. Correlation of plasma cytokine elevations with mortality rate in children with sepsis. *J Pediatr.* 1994;124:105-111.
 37. Ellis G, Adatia I, Yazdanpanah M, Makela SK. Nitrite and nitrate analyses: a clinical biochemistry perspective. *Clin Biochem.* 1998;31:195-220.
 38. Strand OA, Leone A, Giercksky KE, Kirkeboen KA. Nitric oxide indices in human septic shock. *Crit Care Med.* 2000;8:2779-85.

Serum nitric oxide and pediatric sepsis outcomes

ORIGINALITY REPORT

21%

SIMILARITY INDEX

9%

INTERNET SOURCES

17%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

1

Fatemeh Rezaei, Yury Gorbanev, Michael Chys, Anton Nikiforov et al. "Investigation of plasma-induced chemistry in organic solutions for enhanced electrospun PLA nanofibers", Plasma Processes and Polymers, 2018

Publication

2%

2

Submitted to Sriwijaya University

Student Paper

2%

3

Submitted to Higher Education Commission Pakistan

Student Paper

1%

4

Lesley Doughty. "Plasma nitrite and nitrate concentrations and multiple organ failure in pediatric sepsis", Critical Care Medicine, 01/1998

Publication

1%

5

Jaramillo-Bustamante, Juan C., Alejandro MarÃn-Agudelo, Mauricio FernÃndez-Laverde, and JosÃ© BareÃ±o-Silva. "Epidemiology of sepsis in pediatric intensive care units : First

1%

Colombian Multicenter Study*", Pediatric Critical Care Medicine, 2012.

Publication

6

Bettina von Dessauer, Jazmina Bongain, Víctor Molina, Julio Quilodrán, Rodrigo Castillo, Ramón Rodrigo. "Oxidative stress as a novel target in pediatric sepsis management", Journal of Critical Care, 2011

Publication

1%

7

Paul B. Baines. "Nitric oxide production in meningococcal disease is directly related to disease severity", Critical Care Medicine, 06/1999

Publication

1%

8

"Yearbook of Intensive Care and Emergency Medicine 1997", Springer Science and Business Media LLC, 1997

Publication

1%

9

Abraham IJ Gajardo, Bettina von Dessauer, Víctor Molina, Sergio Vera, Matías Libuy, Ramón Rodrigo. "Plasma Antioxidant Potential at Admission is Associated with Length of ICU Stay in Child with Sepsis: A Pilot Study", Fetal and Pediatric Pathology, 2018

Publication

1%

10

www.emjreviews.com

Internet Source

1%

11	von Dessauer, B.. "Oxidative stress as a novel target in pediatric sepsis management", Journal of Critical Care, 201102 Publication	1%
12	preview-ccforum.biomedcentral.com Internet Source	1%
13	journals.sagepub.com Internet Source	1%
14	jamanetwork.com Internet Source	1%
15	link.springer.com Internet Source	1%
16	dmsjournal.biomedcentral.com Internet Source	1%
17	Submitted to Universiti Sains Malaysia Student Paper	<1%
18	www.stemcellethics.net Internet Source	<1%
19	L. ARMSTRONG. "Differential expression of Toll-like receptor (TLR)-2 and TLR-4 on monocytes in human sepsis", Clinical & Experimental Immunology, 5/2004 Publication	<1%
20	PAŞA, Semir, ALTINTAŞ, Abdullah, TÜMER,	<1%

Cemil, DEMİRCİN, Mustafa, ÇİL, Timuçin, BAYAN, Kadim, DANIŞ, Ramazan, TÜZÜN, Yekta, ATAYAN, Yahya, AYYILDIZ, Orhan and BAHÇECİ, Mithat. "Prothrombin time, activated thromboplastin time, fibrinogen and D-dimer levels and von-Willebrand activity of patients with sheehan's syndrome and the effect of hormone replacement therapy on these factors", Akademi Doktorlar Yayınevi, 2010.

Publication

21

Wu, Huang-Pin, Kong Chung, Chun-Yao Lin, Bor-Yiing Jiang, Duen-Yau Chuang, and Yu-Chih Liu. "Associations of T helper 1, 2, 17 and regulatory T lymphocytes with mortality in severe sepsis", Inflammation Research, 2013.

Publication

<1%

22

www.mdpi.com

Internet Source

<1%

23

Gaspar, Heloisa Amaral, Werther Brunow Carvalho, and Artur Figueredo Delgado. "New Approach to Insert Peripherally Inserted Central Catheter in Critically Ill Pediatric Patients :", Pediatric Critical Care Medicine, 2013.

Publication

<1%

24

eprints.ucm.es

Internet Source

<1%

Young A Kim, Eun-Ju Ha, Won Kyoung Jhang,

25	Seong Jong Park. "Early blood lactate area as a prognostic marker in pediatric septic shock", <i>Intensive Care Medicine</i> , 2013 Publication	<1%
26	anesthesiology.pubs.asahq.org Internet Source	<1%
27	ccforum.biomedcentral.com Internet Source	<1%
28	"Biomarkers in Cardiovascular Disease", Springer Science and Business Media LLC, 2016 Publication	<1%
29	fedetd.mis.nsysu.edu.tw Internet Source	<1%
30	www.arca.fiocruz.br Internet Source	<1%
31	Hasan Özkan, Nevin Uzuner, Hale Ören, Nur Çabuk, Hüray İşlekel. "Urinary Nitrite Excretion after Prophylactic Intravenous Immunoglobulin in Premature Infants", <i>Neonatology</i> , 2000 Publication	<1%
32	Submitted to University of Sydney Student Paper	<1%
33	Terregino, C.A.. "Endogenous mediators in emergency department patients with presumed	<1%

sepsis: Are levels associated with progression to severe sepsis and death?", Annals of Emergency Medicine, 200001

Publication

34

Submitted to Universitas Airlangga

Student Paper

<1%

35

Zahra Bahadoran, Parvin Mirmiran, Zhaleh Tahmasebinejad, Fereidoun Azizi, Asghar Ghasemi. "Serum nitric oxide metabolites and hard clinical endpoints: a population-based prospective study", Scandinavian Cardiovascular Journal, 2019

Publication

<1%

36

K.N. Santhosh, D. Pavana, N.B. Thippeswamy. "Impact of scorpion venom as an acute stressor on the neuroendocrine-immunological network", Toxicon, 2016

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On