

THE MANGROVE FLORA IN THE SOUTH AND NORTH COASTAL AREAS OF THE NORTHERN PART OF SULAWESI ISLAND, INDONESIA: AN IMPLICATION FOR SPECIES CONSERVATION

Submission date: 18-Nov-2019 08:48AM (UTC+0700)
by Rignolda Djamaluddin 20

Submission ID: 1215765228

File name: ARTIKEL_MANGROVE_FLORA_SULAWESI_BAGIAN_UTARA.pdf (257.81K)

Word count: 3604

Character count: 21014

1 THE MANGROVE FLORA IN THE SOUTH AND NORTH COASTAL AREAS OF THE NORTHERN PART OF SULAWESI ISLAND, INDONESIA: AN IMPLICATION FOR SPECIES CONSERVATION

1 Rignolda Djamaluddin
Faculty of Fishery and Marine Science, Sam Ratulangi University
rignolda@unsrat.ac.id; rignolda@gmail.com

ABSTRACT

Biogeographically, the mangrove flora in the northern part of Sulawesi Island may include two regions, Indo-Malesia and Asia, Australasia and the Western Pacific, supporting the high diversity of mangrove species. Collections of mangrove specimens have been integrated in various mangrove surveys since 1995, using transect quadrat and spot check methods. As the result, at least 31 species of mangrove were collected from this study area. Three species, *Avicennia alba* Blume, *Campostemon philippinense* (Vidal) Becc, *Sonneratia ovata* Baker seemed to be a typical species in the north coast, explaining its distribution limit in Indo-Malesia region. Three other species of *Aegiceras floridum* Roemer & Schultes, *Heritiera globulus*, and *Osbornia octodonta* F. Muell occurred only in the south coast, confirming its disjunctive distribution. Among all collected species in the two coastal areas, stands of several species including *A. alba* Blume, *Ceriops zippiliana* Blume, *Bruguiera cylindrica*, (Linnaeus) Blume, *S. ovata* Backer, appeared to be uncommon species. This study confirmed that the coastal areas are floristically rich, and there is a concern of species conservation at certain locations.

Keyword: Australasia, conservation, Indo-Malesia, mangrove, Sulawesi

ABSTRAK

Berdasarkan aspek biogeografi, tumbuhan mangrove di bagian utara Pulau Sulawesi termasuk dalam dua wilayah, yakni: Indo-Malesia dan Asia, Australasia dan Pacific Bagian Barat, dimana kondisi di wilayah ini mendukung diversitas spesies mangrove yang tinggi. Pengumpulan spesimen mangrove telah diintegrasikan dalam berbagai survei mangrove sejak tahun 1995 menggunakan metode kuadrat transek dan pengecekan poin. Hasilnya, terkoleksi paling sedikit 31 spesies mangrove di wilayah tersebut. Tiga spesies yakni *Avicennia alba* Blume, *Campostemon philippinense* (Vidal) Becc, *Sonneratia ovata* Baker merupakan spesies tipikal pantai Utara dan kehadirannya juga menjelaskan batas distribusi spesies tersebut di wilayah Indo-Malesia. Tiga spesies lainnya yakni *Aegiceras floridum* Roemer & Schultes, *Heritiera globulus*, and *Osbornia octodonta* F. Muell terdapat hanya di pantai Selatan. Dari semua spesies yang telah terkoleksi beberapa spesies yakni *A. alba* Blume, *Ceriops zippiliana* Blume, *Bruguiera cylindrica*, (Linnaeus) Blume, *S. ovata* Backer, semua merupakan spesies yang tidak umum. Studi ini mengkonfirmasi bahwa spesies mangrove di wilayah pantai bagian Utara Pulau Sulawesi kaya akan spesies mangrove, juga hasil studi mengindikasikan pentingnya konservasi spesies pada beberapa lokasi tertentu.

INTRODUCTION

The origin of the term of mangrove remains ambiguity. It may be related to the old Malay word of 'mangi' or 'manggi-manggi', or 'mangle' in *Rhizophora mangle* Linnaeus (Claridge & Burnett, 1993), or according to Vannucci (1998) 'mangue' - the national language of Senegal. In practice, this term is used for the species of higher plants which have been uniquely successful

in inhabiting intertidal habitats of tropical and sub-tropical regions (Clough, 1979; Duke, 1992; Maxwell, 2015). Mangrove is halophytic, tolerance to salty environment (Spalding *et al.*, 1997), includes a variety of functional forms/ growth habits, such as trees, shrubs, a palm and a ground fern (Duke *et al.*, 1998). According to Srikanth *et al.* (2015), the survival and success of mangrove vegetation in colonizing high tress

habitats is due to its successful adaptation in morphology, anatomy, physiology, and molecule.

In several reports (e.g. Gieasen *et al.* 2006; Spaldi *et al.* 2010; Richards & Friess, 2015; Hamilton & Casey, 2016) the world distribution of mangrove are widely explained. The largest areas of mangrove is in Southeast Asia, encompassing of 5.1 million ha, representing 33.5% of the world's total (Spalding *et al.*, 2010). Indonesia has the largest areas of mangrove in Southeast Asia with almost 60% of Southeast Asia's total (Gieasen *et al.*, 2006). Estimation given by Richard & Friess (2015) that the total mangrove area over this Country was approximately 2,788,683 ha in 2000, and the rate of mangrove loss between 2000 and 2012 was calculated at level of 1.72%.

In the context of biogeographic regions, mangroves in Indonesia are part of Indo-Malesia and Asia, Australasia and the Western Pacific (Duke *et al.*, 1992). There are 50 mangrove species in Indo-Malesia and 47 species in Australasia, including several putative hybrids (Duke *et al.* 1998). In both biogeographic regions, the total number of species is estimated at 57 species, since there is an overlap of 39 species and several species definitely occur either in Indo-Malesia or Australasia (Djamaluddin, 2004). Gieasen *et al.* (2006) claims that Indonesia is the more species diversity of the Southeast Asia countries with 48 species of mangrove. With particular concern to mangroves of the northern part of Sulawesi Island, some 34 true mangrove species may be found (Tomlinson, 1986).

Two factors, oceanic circulation and climate condition, according to Thom (1982), may influence the distribution of mangrove. The marine environment of the north coast is under influence by dominant seawater mass approaching from the northern Pacific Ocean, flowing through Makasar Strait (Van Bennekom, 1988). Bingham & Lukas (1994) explains that Mindanao Current from the coastal areas in the southeast of the Philippines archipelago strengthen the flow of seawater mass from the northern Pacific. Another significant seawater mass is approaching from the

western Pacific flowing through the Maluku Sea. So that, in the view of seawater circulation, the two coastal areas in the northern part of Sulawesi Island seems to be affected by these two different flows of seawater mass. These may influence the distribution patterns of mangrove propagules and seeds in the northern coastal areas of Sulawesi Island.

Two wind systems, north-westerly and south-easterly winds, are reported to have strong influence on rainfall in northern part of Sulawesi Island (Davie *et al.*, 1996; Djamaluddin, 2004). During September and April the north-westerly wind brings moisture after blowing over the South China Sea. This wind arrives in the North Sulawesi in November through the Sulawesi Sea. From the wintery Australian land the south-easterly wind blows towards eastern Sulawesi. This wind causes a short dry season in Manado from August to October. According to data released by Manado Meteorology and Geo-physic Office, the total annual rainfall ranges from 2,501 to 3,000 mm. Within the period from 1973 to 2016, the annual temperature of North Sulawesi varies between 25.5° C and 27.0° C (Djamaluddin, 2018).

The high diversity of mangrove forests in the northern coast in particular the area of Bunaken National Park seems to be unique and unusual in Southeast Asia (Davie *et al.* 1996; Djamaluddin, 2004). Indication of difference in species composition between the north and south coastal areas has been indication in report of Tomini Bay mangrove survey (Djamaluddin, 2015). The interplay between factors of geo-physic, geomorphic and biology is believed to support the mangrove distribution and diversity in the intertidal environments of both coastal areas.

A long-term investigation on mangrove species within the northern coast of Sulawesi Island provides ample time and opportunity to collect and to record any detailed information of mangrove flora in this specific region of Indo-Malesia and Asia, Australasia and the Western Pacific. Geographical distribution limits and patterns as well as population status of certain species may also be clarified from this investigation. Moreover, results from this study may be

important in the context of species conservation.

METHOD

Almost all mangrove forests in the north coast and south coasts have been visited since 1995, and some 235 locations were observed deeply for community structure including collection of mangrove specimens. Two methods of transect quadrat and spot check were used during the observation. At any locations where a new specimen was found, geographical position was marked. Field determination of mangrove specimens was made *in situ* for well-known mangrove

species, and other specimens were collected for further identification for uncommon or expected new mangrove species. Identification of mangrove specimens were based on morphological characteristics by using a number of references of systematic reviews such as Stennis (1955-58), Ding Hou (1958), Percival & Womersley (1975), Blasco (1984), Fernando & Panco (1980), Tomlinson (1986), Mabeber *et al.* (1995), Noor *et al.* (2006). All the specimens used for determination of the flora were photographed and documented. In Figure 1 mangrove areas in the north and south coast covered by the survey are presented.



Figure 1. Surveyed coastal areas in the north and south coasts of the Northern Sulawesi Island.

RESULTS AND DISCUSSION

Mangrove flora

In total, there were at least 31 species of 13 families were identified to occur in the north and south coast of the northern part of Sulawesi Island. Considering the list of Tomlinson (1986) for the biogeographic region between 120° and 135° E, three

species of *Aegialitis annulata* R. Brown, *Bruguiera exaristata* Ding Hou, and *Bruguiera hainesii* C.G. Rodgers were not found. It is also clear that several species are typical north or south coasts in term of their range of distribution. All collections specimen for both coastal areas are presented in Table 1.

Table 1. All mangrove species found in the north and south coastal areas in the northern part of Sulawesi Island.

Family	Species	Common Name
Acanthaceae:	1. <i>Acanthus ilicifolius</i> Linnaeus ^{1,2}	Holly mangrove
	2. <i>Avicennia alba</i> Blume ¹	Api-api putih
	3. <i>Avicennia marina</i> (Forssk.) Vierh ^{1,2}	Grey/white mangrove
Arecaceae:	4. <i>Nypa fruticans</i> (Thunb.) Wurm. ^{1,2}	Mangrove palm
Bombacaceae:	5. <i>Camptostemon philippinense</i> (Vidal) Becc. ¹	Gapas gapas
Combretaceae:	6. <i>Lumnitzera littorea</i> (Jack) Voigt. ^{1,2}	Red-flowered black mangrove
	7. <i>Lumnitzera racemosa</i> Willd ^{1,2}	White-flowered black mangrove
Euphorbiaceae:	8. <i>Excoecaria agallocha</i> Linnaeus ^{1,2}	Milky mangrove, Blind-your-eye
Meliaceae:	9. <i>Xylocarpus granatum</i> König ^{1,2}	Cannonball mangrove
	10. <i>Xylocarpus molucensis</i> Pierre ^{1,2}	Cedar mangrove
Primulaceae:	11. <i>Aegiceras corniculatum</i> (Linnaeus) Blanco ^{1,2}	River mangrove, Black mangrove
	12. <i>Aegiceras floridum</i> Roemer & Schultes ²	-
Pteridaceae:	13. <i>Acrosticum aureum</i> Linnaeus ^{1,2}	Golden mangrove fern
	14. <i>Acrosticum speciosum</i> Wildenow ^{1,2}	Showy mangrove fern
Myrtaceae	15. <i>Osbornia octodonta</i> F. Muell ²	Myrtle mangrove
Rhizophoraceae:	16. <i>Bruguiera cylindrica</i> (Linnaeus) Blum ^{1,2}	Large-leafed orange mangrove
	17. <i>Bruguiera gymnorhiza</i> (Linnaeus) Lamk. ^{1,2}	Large-leafed orange mangrove
	18. <i>Bruguiera parviflora</i> Weight & Arnold ex Griffith ^{1,2}	Small-leafed orange mangrove
	19. <i>Bruguiera sexangula</i> (Lour.) Poir. ^{1,2}	Upriver orange mangrove
	20. <i>Ceriops zippeliana</i> Blume ²	Tengat merah
	21. <i>Ceriopa tagal</i> (Perr.) C.B. Robinson ^{1,2}	Rib-fruited yellow mangrove
	22. <i>Rhizophora apiculata</i> Blum ^{1,2}	Corky stilt mangrove
	23. <i>Rhizophora mucronata</i> Lamk. ^{1,2}	Upstream stilt mangrove
	24. <i>Rhizophora stylosa</i> Griffith ^{1,2}	Long-styled stilt mangrove
	Rubiaceae:	25. <i>Scyphiphora hydrophyllacea</i> Gaertn. f. ^{1,2}
Lythraceae:	26. <i>Pemphis acidula</i> Forst & Forst ^{1,2}	Bonsai mangrove
	27. <i>Sonneratia alba</i> J. Smith ^{1,2}	White-flowered apple mangrove
	28. <i>Sonneratia caseolaris</i> (Linnaeus) Engler ^{1,2}	Red-flowered apple mangrove
Sterculiaceae:	29. <i>Sonneratia ovata</i> Backer ¹	Ovate-leafed apple mangrove
	30. <i>Heritiera littoralis</i> Dryand ^{1,2}	Looking-glass mangrove
	31. <i>Heritiera globulus</i> ²	-

Notes: ¹⁾ species found in the north coast, ²⁾ species found in the south coast

There are several changes in species listed in Tabel 1 comparing to the previous species lists for the north coast (Davie *et al.*, 1996; Djamaluddin, 2004) and the south coast (Djamaluddin, 2015). Part of these

changes in particular for the mangrove flora in Bunaken National Park has been explained in Djamaluddin (2018). Species of *Avicennia marina* var. *rumphiana* (Hall. F.) Bakhuizen was excluded from the previous

list and corrected as *Avicennia marina* (Forssk.) Vierh. Specimen that was identified as *Avicennia officinalis* Linnaeus (in the list of Davie *et al.*, 1996) was verified as *Avicennia alba* Blume. Detail check of specimen previously identified as *Kandelia candel* (Linnaeus) Druce confirmed that the specimen was *Bruguiera cylindrica* (Linnaeus) Blum. Following the development in mangrove species taxonomy, changes were made to the names of two species, *Ceriops decandra* (Griff.) Ding Hou and *Xylocarpus mekongensis* Pierre; changed as *Ceriops zippeliana* and *Xylocarpus molucensis* respectively.

As can be seen from data in Table 1, there were three species of *A. alba*, *C. philippinense*, and *S. ovata* were typical for mangrove forests in the north coast. Trees of *A. alba* occurred along the mainland coast of the northern part of Bunaken National Park. Trees of this species shared the same habitat with trees of *A. marina*. Trees of *S. ovata* were collected from three locations at Pungkol (southern section of Bunaken National Park) in a dense population, mangrove forest near Tiwoho Village and Mantehage Island in individual trees. Type of habitat of this species is 'seasonally or regularly freshwater influenced landward fringe' (Djamaluddin, 2018). The presence of freshwater seems to be a limiting factor for this species survival at landward fringe. Trees of *C. philippinense* were found only at small isolated location in Mantehage Island (1°42'59.4" N; 124°45'31.2" E). This species is very common in the Philippines (Giaesen *et al.*, 2006). It is also reported to occur in Eastern Kalimantan (Mukhlisi & Sadiyah, 2014) and western coast of Central Sulawesi (Wahyuningsih *et al.* 2012). However, this species may be absent in the south coast of northern Sulawesi (Damanik and Djamaluddin, 2012; Djamaluddin, 2015), as well as in the West Papua (Prawiroatmodjo & Kartawinata, 2014).

Typical species of the south coast included *A. floridum*, *H. globulus*, and *O. octodonta*. Trees of these species were not common in this area, and each of these inhabited very special and isolated habitat.

Detail explanations of the distribution pattern for all mangrove species in the

northern coast of Sulawesi island may need further investigation. However, the distribution limit of *C. philippinense* in the biogeographic region of Indo-Malesia can be explained from the presence of this species in Mantehage Island. It is expected that Mantehage Island is probably the distribution limit of this species in the eastern part of Indonesia. A question that needs to be answered is that why trees of this species were being isolated in a small area in Mantehage Island. Domination of seawater circulation from the northern Pacific Ocean flowing through the southeast of the Philippines may be a significant factor that relates to the presence of *C. philippinense* in Mantehage Island. The arrival of seeds of this species in Mantehage Island may be under influence of this seawater circulation. Another question is for the presence of *O. octodonta* in Panua Reserved (0°27'45.3" N; 121°58'54.5" E) in the south coast. This species shows a disjunctive distribution, since it occurs in considerably separated places in the Philippines, Kalimantan, eastern Malaysia, Palau, and northern Australia.

Implication for species conservation

In the International Union for Conservation of Nature (IUCN) Red List 2010, the species of *C. philippinense* categorized as "an endangered species", the second most severe conservation status for wild population, meaning that this species is more likely to become extinct. In Mantehage Island where this species was found there were only small population of this species, and they were subjected to logging. As an endangered species, trees of this species as well as the habitat needed to be protected. Trees of *S. ovata* were found in three locations in Bunaken National Park, but they were very rare growing in an individual tree at landward areas subjected to sedimentation. This species was categorized as "a near threatened" species in the IUCN Red List that this needed to be protected. Another species that was categorized as "a near threatened" species was *A. floridum*, occurring only at Pulau Pondang (0°25'00.3" N; 124°20'59.8" E) in the south coast. Although its conservation status in the

IUCN Red List was categorized as “a least concern”, trees of *B. cylindrica* were rare, so that this species needed to be protected locally.

CONCLUSION

The mangrove flora in the northern coast of Sulawesi Island was floristically rich with at least 31 species of 13 families. In the context of species composition, there was difference between the north and south coasts. Three species occurred only in the north coast including *A. alba*, *C. philippinense*, and *S. ovata*, meanwhile three other species were only found in the south coast including *Aegiceras floridum*, *H. globulus*, and *O. octodonta*. The presence of *C. philippinense* in Mantehage Island confirmed its distribution limit in the geographic region of Indo-Malesia. Protection was needed for several species including *A. floridum*, *C. philippinense*, and *S. ovata*, due to their conservation status, and for *B. cylindrica*, *H. globules*, and *O. octodonta* due to their locally rare population.

ACKNOWLEDGMENT

Financial support had come in part from various institutions including USAID-NRM, Mangrove Action Project (MAP), Rufford Small Grant (SG), Whitley Fund for Nature (WFN), Global Environmental Fund-Small Grants Programme (GEF-SGP), Balai Taman Nasional Bunaken, IUCN, Susclam Project, Mangrove for Future (MFF), and Australian Agency for International Development (AusAid). I am sincerely indebted to Villagers at Tiwoho Village, Rap-rap Village, Bango Village, Staffs of Perkumpulan KELOLA, Antra-Sulut, my students at the Faculty of Fishery and Marine Science, Kelompok Kerja Mangrove in Tomini Gulf, for their contributions during field works. I also thank to Mr. Afif Kaumbo and Mr. Brama Jabar for helps in providing spatial information and maps.

LITERATURE CITED

- [1] Bingham, FM., & Lukas, R. (1994). The southward intrusion of North Pacific intermediate along the Mindanao coast. *Journal of Physical Oceanography*, 24: 141-154.
- [2] Blasco, F. (1984). Taxonomic considerations of the mangrove species. In: Snedaker, SC. & Snedaker, JG. (Eds.). 'The Mangrove Ecosystem: Research Methods', pp.81-90. UNESCO.
- [3] Claridge, D., & Burnett J. (1992). Mangroves in focus. Marino Lithographics, Queensland.
- [4] Clough, BF. (1979). Mangrove ecosystem in Australia: structure, function and management. Proceedings of the Australian National Mangrove Workshop. Australian Institute of Marine Science, Cape Ferguson, 18-20 April 1979.
- [5] Damanik, R., & Djamaluddin, R. (2012). Mangrove map of Tomini Gulf. Sustainable Coastal Livelihoods and Management Program, CIDA, IUCN, Lestari Canada. [Indonesian]
- [6] Davie, J., Merril, R., & Djamaluddin, R. (1996). The sustainable use and conservation of the mangrove ecosystem of the Bunaken National Park. USAID, Jakarta.
- [7] Ding Hou. (1958). Rhizophoraceae. *Flora Malesiana*, I(5): 429-493.
- [8] Djamaluddin, R. (2004). The dynamics of mangrove forest in relation to die-back and human use in Bunaken National Park, North Sulawesi, Indonesia. [Dissertation]. University of Queensland, Brisbane.
- [9] Djamaluddin, R. (2015). The mangrove flora in Tomini Bay. The 5th International Conference on Plant Diversity, Universitas Jenderal Sudirman, Purwokerto 20 – 21 August 2015.
- [10] Djamaluddin, R. (2018). The mangrove flora and their physical habitat characteristics in Bunaken National Park, North Sulawesi, Indonesia. *Biodiversitas Journal of Biological Diversity*, 19(4):1303-1312.

- [11] Duke, NC., Ball, MC., & Ellison, JC. (1998). Factor influencing biodiversity and distribution gradients in mangrove. *Global Ecology and Biogeography Letters*, 7(1): 27-47.
- [12] Duke, NC. (1992). Mangrove floristics and biogeography. In: Robertson, AI., & Alongi, DM. (Eds.). 'Coastal and Estuarine Studies', pp.63-100. American Geophysical Union, Washington, D.C.
- [13] Duke, N., Kathiresan, K., Salmo III, S.G., Fernando, ES., Peras, JR., Sukardjo, S., Miyagi, T., Ellison, J., Koedam, NE., Wang, Y., Primavera, J., Jin Eong, O., Wan-Hong Yong, J., & Ngoc Nam, V. (2010). *Camptostemon philippinense*. *The IUCN Red List of Threatened Species* 2010: e.T178808A7612909. <http://dx.doi.org/10.2305/IUCN.UK.2010-2.RLTS.T178808A7612909.en>.
- [14] Fernando, ES., & Pancho, JV. (1990). Mangrove trees of the Philippines. *Silvatrop Philipp. Forest Research Journal*, 5(1):35-54.
- [15] Giesen, W., Wulffraat, S., Zieren, M., & Scholten, L. (2006). Mangrove guide book for Southeast Asia. FAO and Wetlands International.
- [16] Hamilton, SE., & Casey, D. (2016). Creation of a high spatio-temporal resolution global database of continuous mangrove forest cover for the 21st century (CGMFC-21). *Global Ecology and Biogeography*, 25(6):729-738.
- [17] Mabberley, CM., Pannel, CM., & Sing, AM. (1995). *Flora Malesiana: Seri I Spermathophyta*, 12(1): 371-81.
- [18] Maxwell GS. (2015). Gaps in mangrove science. *ISME/GLOMIS*, 13(5): 18-38.
- [19] Mukhlisi, Sidiyasa K. 2014. Structure and composition of mangrove species in Berau Mangrove Information Centre (PIM), Kalimantan Timur. *Indonesian Forest Rehabilitation Journal*, 2(1): 25-37. [Indonesian]
- [20] Noor, YR., Khazali, M., & Suryadiputra INN. 2006. Mangrove introduction guide. Ditjen PPHKA-Wetland International, Bogor. [Indonesian]
- [21] Percival, M., & Womersley, JS. (1975). Floristics and ecology of the mangrove vegetation in Papua. *New Guinea Botany Bulletin*, 8.
- [22] Prawiroatmodjo, S., & Kartawinata, K. (2014). Floristic diversity and structural characteristics of mangrove forest of Raja Ampat West Papua, Indonesia. *Reimwardtia*, 14(1): 171-180
- [23] Richards, DR., & Fries, DA. (2016). Rates and drivers of mangrove deforestation in Southeast Asia 2000-2012. *PNAS*, 113(2): 344-349.
- [24] Spalding, MD., Blasco, F., & Field, CD. (1997). *World Mangrove Atlas*. International Society for Mangrove Ecosystems, Okinawa.
- [25] Spalding, MD., Kainuma, M., & Collins L. (2010). *World Atlas of Mangroves*. Earthscan, London.
- [26] Srikanth, S., Lum, SKY., & Chen, Z. (2015). Mangrove root: adaptations and ecological importance. *Tress*, 30(2): 451-465.
- [27] Stennis, CGGJ. Van. (1995-1958). *Flora Malesiana*. Nooordhoff-Kolff NV, Djakarta.
- [28] Thom, BG. (1982). Mangrove ecology – a geomorphological perspective. In: Clough, BF. (Ed). 'Mangrove Ecosystem in Australia: Structure, Function and Management'. AIMS with ANU Press, Canberra.
- [29] Tomlinson, PB. (1986). *The botany of mangroves*. Cambridge University Press, New York.
- [30] Van Bennekom, A. (1988). Deep-water transit times in the eastern Indonesian basins, calculated from dissolved silica in deep and interstitial waters. *Netherland Journal of Sea Research*, 22(4): 341-354.

- [31] Vannucci, M. (1998). The mangrove ecosystem: an overview of present knowledge. *Revista Brasileira De Biologia*, 58:1-15.
- [32] Wahyuningsih, EP., Suleman, SM., & Ramadanil. (2012). Structure and composition of mangrove vegetation in Desa Lalombi Kecamatan Banawa Selatan Kabupaten Donggala. *Biocelebes*, 6(2): 84-100. [Indonesian]

THE MANGROVE FLORA IN THE SOUTH AND NORTH COASTAL AREAS OF THE NORTHERN PART OF SULAWESI ISLAND, INDONESIA: AN IMPLICATION FOR SPECIES CONSERVATION

ORIGINALITY REPORT

11%

SIMILARITY INDEX

4%

INTERNET SOURCES

4%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to Universitas Riau

Student Paper

2%

2

Ni Luh Gede Rai Ayu Saraswati, Yulius -, Agustin Rustam, Hadiwijaya L. Salim, Aida Heriati, Eva Mustikasari. "Kajian Kualitas Air Untuk Wisata Bahari Di Pesisir Kecamatan Moyo Hilir Dan Kecamatan Lape, Kabupaten Sumbawa", Jurnal Segara, 2017

Publication

1%

3

eprints.utas.edu.au

Internet Source

1%

4

Submitted to Higher Education Commission Pakistan

Student Paper

1%

5

Jyoti Srivastava, Vandana Prasad. "Evolution and paleobiogeography of mangroves", Marine Ecology, 2019

1%

- | | | |
|----|--|-----|
| 6 | era.daf.qld.gov.au
Internet Source | <1% |
| 7 | Submitted to University of Aberdeen
Student Paper | <1% |
| 8 | www.salmonriverdata.co.uk
Internet Source | <1% |
| 9 | theses.saurashtrauniversity.edu
Internet Source | <1% |
| 10 | laptop.deh.gov.au
Internet Source | <1% |
| 11 | allafrica.com
Internet Source | <1% |
| 12 | Submitted to El Dorado Union High School
District
Student Paper | <1% |
| 13 | link.springer.com
Internet Source | <1% |
| 14 | Maria Lucia Negreiros-Fransozo. "Chapter 29
Size Variation in the Grapsid Crab Aratus
Pisonii (H. Milne-Edwards, 1837) Among
Populations of Different Subtropical
Mangroves", Springer Nature, 2002
Publication | <1% |
-

15	opac.ll.chiba-u.jp Internet Source	<1%
16	www.iwmi.cgiar.org Internet Source	<1%
17	www.slideshare.net Internet Source	<1%
18	Submitted to Universidade Nova De Lisboa Student Paper	<1%
19	"Mangrove Management, Assessment and Monitoring", Tropical Forestry Handbook, 2015. Publication	<1%
20	Jerry van Dijk, Roland E. van der Vliet, Harm de Jong, Maarten J. Zeylmans van Emmichoven et al. "Modeling direct and indirect climate change impacts on ecological networks: a case study on breeding habitat of Dutch meadow birds", Landscape Ecology, 2015 Publication	<1%
21	Mangrove Ecosystems of Asia, 2014. Publication	<1%
22	Submitted to University of Wales, Bangor Student Paper	<1%
23	Submitted to Universiti Teknologi MARA Student Paper	<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On