

the revision for the paper "Effectivity of Arcangelisia flava as immunostimulant to prevent streptococcosis on Nile tilapia...."

Yahoo/Inbox

gavriloaie ionel claudiu <ionelclaudiu@yahoo.com>

To:maryani@fish.upr.ac.id,Benny Pratasik

Wed, Nov 28, 2018 at 8:31 PM

Dear Dr. Pratasik,

Here you have the revision for this paper. The article is very good regarding the subject and the presentation. The English is also good enough, but the main problem was with the references. I have made lots of corrections, both in list and quotations in text.

For the future papers, please pay much more attention to this aspect.

Work only in attached document, address all the comments and highlight with a bright color all the changes you will do. Leave all the red corrections as they are.


Regarding the publication fee of 200 USD + bank taxes, you already received the variants of payment.

Thank you very much!

Cordially yours,

Claudiu Gavrioloaie

Korespondensi revisi paper.



Effectivity of *Arcangelisia flava* Merr as immunostimulant to prevent streptococcosis on Nile tilapia, *Oreochromis niloticus*

¹Maryani, ¹Sinta S. Monalisa, ¹Rosita, ¹Mohamad Rozik,
²Silvester B. Pratasik

¹ Aquaculture Program, Fisheries Department, Faculty of Agriculture, Palangkaraya University, Jl. Yos Sudarso, Kampus Tunjung Nyaho, Palangkaraya, Central Kalimantan, Indonesia; ² Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl.Kampus Bahu Manado-95115, North Sulawesi, Indonesia. Corresponding author: Maryani, maryani@fish.upr.ac.id

Abstract. This study aims to inform the effectivity of *Arcangelisia flava* Merr. application in feed as immunostimulant on total leucocyte, neutrofil, monocyte, lymphocyte, and survivorship of Nile tilapia *Oreochromis niloticus* infected with *Streptococcus agalactiae*. Data were descriptively presented and qualitatively described in the form of histograms. Results showed that the highest number of leukocytes was found in treatment E, 81.534 cells mL⁻¹ and the lowest in treatment A, 21.379 cells mL⁻¹. The highest number of neutrophils occurred in treatment E, 35.25% and the lowest in treatment A, 30.26%. The highest number of lymphocytes was recorded in treatment E, 80.50% and the lowest in treatment A, 66.50%. The highest number of monocytes count also occurred in treatment E, 13.00%, and the lowest in treatment A, 9.00%. *A. flava* could also affect the survivorship. The highest survival was found in treatment E, 91.66%, and the lowest in treatment A, 76.33%.

Key Words: immunostimulant, leucocyte, neutrophil, monocyte, lymphocyte.

Introduction. Disease infection is a constraint in aquaculture that can reduce the fisheries productivity. According to Park et al (2009), streptococcosis is caused by bacteria *Streptococcus* sp. and one of the diseases that often attack Nile tilapia, *Oreochromis niloticus* and can increase fish mortality up to 50%. The disease comes up when significant temperature fluctuation occurs, so that the immune system of the fish decreases. *Streptococcus* sp. enters the body through digestive system (Irianto 2004). The initial symptoms exhibit that the fish stomach looks bigger. The bacteria will enter the blood vessels reaching the kidney. The fish kidney infected with Streptococcosis, particularly Nile tilapia, will look pale and swell. The fish will show whirling movements when the bacteria *Streptococcus* sp. infects the brain's nerve system. The disease symptom can also be known from swelling and bleeding in the fish eyes (exophtalmia).

Prevention of streptococcosis is still done using antibiotics. However, the effect of continuous use of the antibiotics can yield the resistance of pathogenic microorganisms to the antibiotic and leave some residues in fish and their environment (FAO 2008; Widya 2014). Therefore, safer mitigation and disease prevention alternatives are needed for both fish and their environment.

The use of immunostimulant can be conducted as protection and prevention efforts of disease infections in fish. It can increase the natural and adaptive immune system of the fish (Kani et al 2003). Immunostimulant has higher superiority than other methods, since it is capable of increasing the non-specific immune response of the aquatic animals to prevent various disease infections from bacteria or viruses (Sakai 1999; Galindo-Villegas & Hosokawa 2004; Barman et al 2013; Labh & Shakya 2014).

Arcangelisia flava has been known by Central Kalimantan communities, especially Dayak people as yellow root and utilized as traditional medicine with its ability to cure

Commented [indra1]: no need for this here

Commented [indra2]: what do you mean? May be to estimate?

Commented [indra3]: add few short conclusive remarks

Commented [indra4]: if you say "infected with...", you have to say with *Streptococcus*, which is the bacterial species, not with streptococcosis, which is the disease. It is like you would say "I am infected with a flu" instead "I am infected with a flu virus"

Commented [indra5]: all these have to be sustained with reference(s)

various diseases, one of which is hepatitis. Phytochemical test revealed that *A. flava* was detected to hold secondary metabolites, namely alkaloid, saponin, terpenoid, and flavonoid (Maryani et al 2013). All active compounds contained in *A. flava* enable it to be developed as an immunostimulant.

The objective of the study is to know the effectivity of *A. flava* in the feed as immunostimulant on total leukocytes, neutrophils, monocytes, and lymphocytes, and the survivorship of Nile tilapia infected with *Streptococcus agalactiae*.

Material and Method. This study was carried out from May 10th to June 11th, 2017. Test animals were Nile tilapia taken from Freshwater Aquaculture Development Center in Mandiangin, Banjar Regency, South Kalimantan, with mean size of 8.0±1.2 cm, as many as 20 ind tank⁻¹. Each treatment had 3 replications, so that there were 300 individuals of fish employed. The fish were firstly acclimated in the fiberglass tank for 7 days to guarantee no streptococcosis symptom and other infection. During acclimatization, the fish were fed *ad libitum* twice a day and siphoned every morning before feeding.

A. flava extract was obtained through extraction, as much as 2 kg of *A. flava* was wind-dried. The dry material was blended and then macerated in 8.5 L of 96% ethanol for 24 hours until the supernatant was formed. The supernatant was evaporated using vacuum rotavapour at 40°C at a speed of 120 times per minute. The extract was diluted using distilled water with treatment concentration.

Bacterium *S. agalactiae* used in the test was obtained from Freshwater Aquaculture Development Center of Mandiangin, Banjar Regency, South Kalimantan. Before use, the bacterium was isolated on Brain Heart Infusion Agar (BHIA) sterilized in the autoclave. The isolate of *S. agalactiae* was taken using ose needle and inserted into the agar media using scratch method and incubated at 37°C for 24 hours.

The commercial feed used contained 30-32% protein. It was added with *A. flava* extract with treatment concentrations 0 ppm (A), 50 ppm (B), 100 ppm (C), 150 ppm (D), 200 ppm (E), and 250 ppm (F). Extract addition was done by evenly spraying the feed up to be homogenously mixed and then wind-dried. *A. flava* extract-containing feed was given at 5% of body weight twice a day for 21 days and then exposed to infection of bacterium *S. agalactiae* in the water at the doses of 4 mL with a density of 10⁷ cfu mL⁻¹ for each aquarium.

This study was experimental using Complete Randomized Design with 6 treatments and 3 replications. Parameters observed were total leukocytes, neutrophils, monocytes, and lymphocytes, and the survivorship. The survival rate of Nile tilapia was assessed using Effendie (1979):

$$SR = Nt/No \times 100$$

where: SR = survival rate;

Nt = number of fish at the end of observation;

No = number of fish at initial study.

Supporting parameters measured were water temperature, pH, dissolved oxygen (DO), and ammonia. Measurements of water temperature, pH, and DO were done every morning to control the culture environmental conditions. Ammonia was measured at the end of treatment.

Results and Discussion

Leukocytes. Number of leukocytes at treatment A, B, C, D, and E increases, but it decreases at treatment F (Figure 1). The highest number of leukocytes after treated with *A. flava* extract was 81.534 cells mL⁻¹ at treatment E and the lowest was 21.379 cells mL⁻¹ at treatment A, indicating that treatment E is an appropriate dose of the immunostimulant to support the immune system well. Increased mean total leukocytes could result from the effect of *A. flava* extract administration.

Commented [indra6]: how do you figure this? You have 6 treatments (A to F), each one involving 20 fish, so, you have to have 20x6 = 120 fish for the experiment. Since you had 3 replications for each treatment, you had to have 360 fish in total, not 300.

Commented [indra7]: the whole plant was used?

Commented [indra8]: provide the manufacturer and type

Commented [indra9]: on the section *Water quality* you do not mention at all this parameter




Figure 1. Number of leukocytes in Nile tilapia treated with *A. flava* under different doses after infected with *S. agalactiae*.

Commented [indra10]: on the left side, replace sel/ml with cell mL⁻¹

Immunostimulant is a chemical compound of drugs or other materials that is capable of increasing the specific and non-specific response mechanisms of the fish (Anderson 1992; Galeotti 1998; Mastan 2015). The administration of *A. flava* extract as much as 50 ppm to 250 ppm to the test fish exhibited increment of total leukocytes and gave significantly different effect from that of 0 ppm. An appropriate dose of immunostimulant administration will raise the complimentary activities, opsonin macrophage, phagocytosis, and polymorphonuclear leukocytes (PMN) that help eliminating the entering anti-gene so that fatal infection development does not occur. Increased number of total leukocytes indicates the humoral and cellular leukocyte responses to prevent the presence of bacteria (Erlinger et al 2004; Finlay & McFadden 2006; Magnadottir 2006).

Commented [indra11]: this word is expressing a compliment; praising or approving.

The correct word you should use is complementary - serving to fill out or complete

Leukocytes play important role in fish defensive system against pathogenic infections (Anderson & Siwicki 1995). When infections occur, the leukocytes will be transported to the infected part to give quick protection against the infectious gene (Sadikin 2002). Increased leukocyte population could result from increased cell division activity and mitogenic effect of the immunostimulant used. The mitogenic compounds will activate the defence cells to differentiate that synthesize DNA in the lymphocytic cells and increase the leukocyte population (Faulman et al 1983; Bly et al 1986; Rorstad et al 1993).

Commented [indra12]: rephrase

The present study revealed that total leukocytes rose with addition of *A. flava* extract dose, except slight decline at the dose of 250 ppm (F), but the number of leukocytes was higher than that in 0 ppm (A). It means that *A. flava* extract holds immunostimulant-containing compounds that could induce the body resistance against *S. agalactiae*. According to Maryani et al (2013), *A. flava* contains secondary metabolites, one of which is flavonoid. The working mechanism of flavonoid is to inhibit the function of cell membrane through complex formation against the extracellular protein that disturbs the wholeness of the bacteria cell membrane. It is done by denaturing the bacteria cell protein and by making cell membrane irreparable (Nuria et al 2009; Naiborhu 2002; Shib et al 2015). Other study found that flavonoid mechanism in cell membrane function prevention is to distort the cell membrane permeability, restrain the cell membrane function through cell membrane permeability distortion, and impede the enzymatic binding, such as ATPase and phospholipase (Li et al 2003).

Neutrophils. Figure 2 shows that number of neutrophils increases at treatment B, C, D and E, but declines at treatment F. The highest increment was recorded at treatment E, 35.25 %, and the lowest at treatment A, 30.25%.




Figure 2. Number of neutrophils in Nile tilapia treated with *A. flava* under different doses after infected with *S. agalactiae*.

Neutrophils also rose with increased dose of *A. flava* extract and slightly declined at the treatment of 250 ppm (F), but they were higher than those at control treatment. Neutrophils work to conduct phagocytotic activities against antigens entering the fish body. This study found that increased total leukocytes were followed with increased number of neutrophils. In general, number of neutrophils increases at bacterial infections since they are released from the blood vessel to the infected location. Major function of neutrophils is to destroy alien materials through phagocytosis, i.e. chemotaxis, particle attachment to the cell, particle ingestion by the cell, and particle destruction by lysosome in the phagosome (Tizard 1998; Karuthapandi & Innocent 2010). Releasing neutrophils from the blood vessels when infection occurs could be caused by external chemical stimulation or chemotaxis (Vadstein 1997; Kozinska & Guz 2004; Mishra et al 2006).

Neutrophil is polymorphonuclear (PMN) cell with the shortest life time and will die through apoptosis, approximately 24 hours. Increased number of neutrophils could result from the entry of alien matter in the fish body. This increase could result from immunostimulant administration of *A. flava* extract containing secondary metabolites, one of which is flavonoid. This compound has direct immune effect and stimulates the granulocytes development indicated with increased amount of neutrophils in the fish body (Lou et al 2012). Neutrophils can yield monocyte and macrophage stimulating substance. The secretion can raise the phagocytosis and ROS involved in intracellular killing. This cell is a powerful phagocyte. Phagocytosis is done by approaching the foreign particle and releasing pseudopodia to all directions around the particle. A neutrophil can phagocytize 5-20 bacteria (Tizard 1998).

Lymphocytes. Treatments A, B, C, D, and F resulted in lymphocyte range from 66.50% to 78.25%. The range reflects that the lymphocytes are in normal condition (Galindo-Villegas & Hosokawa 2004), 60-85%, but lower than that of normal fish reported by Smith (2007), > 80%. In treatment E, number of lymphocytes increased up to 80.50%. It means that humoral immune response of the fish is in good condition, so that the resistance to the alien material and antibody formation occur. Increased number of lymphocytes occurs as the lymphocytes that are temporarily in the blood will migrate to various lymph nodes or into the spleen. When they meet the alien materials, the lymphocyte will grow and undergo mitosis of plasma cell that functions as antibody producer (Mishra et al 2009).




Figure 3. Number of lymphocytes in Nile tilapia treated with *A. flava* under different doses after infected with *S. agalactiae*.

This result is supported by Mundriyanto et al (2002) that the role of lymphocytes in the immune system is to release immune substances for body protection by detecting the antigens through specific receptor of the cell membrane. T lymphocyte cells will work if the immune response occurs against the antigens by differentiating to different effector T cells. The cells will then produce different cytokine depending upon the types of the immune responses. Th1 cells produce interferon gamma (IFN- γ) and interleukin-2 (IL-2) that will raise the activity of macrophagic cells, natural killer (NK) cells, and cytotoxic T (Tc) cells. This mechanism will develop quick phagocytotic activities to destroy the antigens and intracellular microorganisms. Th2 cells will produce several types of cytokines, such as IL-4, IL-5, IL-13, that will stimulate the antibody development, activate the mast cells and the eosinophils to make stronger defense against the extracellular antigens (Tizard 1998; Kresno 2001; Abbas et al 2010; Watson & Preedy 2013).

Commented [indra13]: to be added to the references list

Commented [indra14]: ???

Monocytes. The present study found that the highest fish monocytes were recorded in treatment E, 13.00%. This condition could result from the alteration of monocyte to macrophage and migrate to the infected locations to phagocytize the antigens of *S. agalactiae*.




Figure 4. Number of monocytes in Nile tilapia treated with *A. flava* under different doses after infected with *S. agalactiae*.

Increased monocytes during infection, according to Selvaraj et al (2005) stimulate the monocyte to destroy the bacteria. Monocytes are cells in the blood vessel and develop to macrophage when they are activated, in which macrophages possess stronger phagocytic ability than neutrophils even though the granulocytes have higher amount. Maftuch (2007) also describes that inflammation from infection or antigen-antibody reaction will increase monocyte production twice more.

According to Bijanti (2005), monocyte distribution in the blood becomes shorter (1-2 days), and then the cell migrates in the tissues where it differentiates further to be macrophage. Secombes (1990), Anderson & Siwicki (1995), Bastami et al (2009), and Harikrishnan et al (2010) added that proportion of monocytes was low in leukocyte population, but it could rise about 38% in short time when infection occurred.

Survivorship. The present study found that the highest survivorship of fish treated with *A. flava* extract was recorded in treatment E, 91.66%, and the lowest in treatment A, 76.33% (Figure 5). It reveals that treatment E is the most effective compared to other treatments.

High survivorship of the test fish after *A. flava* extract administration reflects the potential of the extract to reduce *S. agalactiae* population. It could be due to the presence of antibacterial substances in *A. flava*, flavonoid. Bacterial infection in fish could occur through wounded body surface, food or gills. Then the bacteria enter the blood vessels, disperse, and yield blood poisoning. *S. agalactiae* can also infect the fish since they can recognize certain cell receptors, then the bacteria kill and break down the host cells by producing the extracellular enzymes, and used the breakdown host cells as nutrients for their growth.




Figure 5. Survivorship of Nile tilapia treated with *A. flava* under different doses after infected with *S. agalactiae*.

The population development of pathogenic bacteria causes inflammation around the infected location and makes the wound get worse to be a haemorrhage. The breakdown of fish body cells in the infected area damages the blood vessel, the pathogenic bacteria enter and disperse through blood circulation to the entire fish body. If these haemorrhages attack the important organs, such as respiration organs, digestive tract, kidney, and liver, they will cause mortality (Jun et al 2010).

The presence of flavonoid in *A. flava* extract as antibacterial is to denature the protein cells, destroy the bacterial cell membrane so that the bacteria cannot grow, and prevent toxin release until the bacteria are killed (Hendra et al 2011; Rijayanti 2014). Besides, flavonoid is an antioxidant as well and capable of increasing the immune system function because the leukocytes as antigens feeder are produced faster and the lymphoid system is also activated faster. The present finding is supported by Susilo (2012) that flavonoid compounds contained in dandelion (*Taraxacum officinale*) extract can raise the

immune activities. According to Suhirman & Winarti (2013), the secondary metabolites of plants, such as flavonoids, are potential as immunostimulant, antibacterial, antiviral or other microorganisms. The potential of flavonoids as immunostimulant has also been reported by Susanti et al (2012) in cinnamon (*Cinnamomum verum*) extract against *Salmonella enteritidis*, by Ulfah (2014) for roselle (*Hibiscus sabdariffa*) flower, and by Kusmardi et al (2007) in tropical shrub (*Cassia alata*) leaf.

Water quality. Poor water quality condition could inhibit growth and cause disease in fish. Water temperature during the study ranged from 27 to 29°C, and it could support the survivorship of Nile tilapia. It is in agreement with Khairuman & Amri (2012) that optimum water temperature for feeding activities ranged from 25 to 27°C. Water pH ranged from 6.90 to 7.30 during the study, highly supported the survivorship of Nile tilapia, and this range belongs to good condition for fish production suggested by Cholik et al (2005). Dissolved oxygen ranged from 5.00 to 6.60 mg L⁻¹ and these are higher than that required by National Standard Bureau of Indonesia (BSN 7550: 2009), 3.0 mg L⁻¹, for culture media of Nile tilapia.

Conclusions. Application of *A. flava* as non-specific immunostimulant in the fish feed could raise the amount of leukocytes, neutrophils, lymphocytes, and monocytes in Nile tilapia infected with *S. agalactiae*. The highest number of leukocytes, neutrophils, lymphocytes and monocytes was found in treatment E, 81.534 cells mL⁻¹, 35.25%, 80.50%, and 13.00% respectively. The administration of *A. flava* in the feed as non-specific immunostimulant affected the survival rate of Nile tilapia infected with *S. agalactiae* with the highest in treatment E, 91.66%, and the lowest in treatment A, 76.33%.

References

- Abbas A. K., Lichtman A. H., Pillai S., 2010 Cellular and molecular immunology. Saunders Elsevier, Philadelphia, 564 pp.
- Anderson D. P., 1992 Fish immunology. TFH Publication Ltd., Hongkong, 239 pp.
- Anderson D. P., Siwicky A. K., 1995 Basic haematology and serology for fish health programs. In: Diseases in Asian Aquaculture II. Shariff M., Arthur J. R., Subasinghe J. P. (eds), Manila, Philippines, pp. 185-202.
- Barman D., Nen P., Mandal S. C., Kumar V., 2013 Immunostimulants for aquaculture health management. Journal of Marine Science: Research and Development 3(3):134.
- Bastami K., Darvish M. A. H., Zaragabadi A. M, Salehi Mir S. V., Shakiba M. M. 2009 Measurement of some haematological characteristics of the wild carp. Comp Clin Pathol 18:321-323.
- Bijanti R., 2005 [Hematology Ikan]. Bagian Ilmu Kedokteran Dasar Veteriner, Fakultas Kedokteran Hewan Universitas Erlangga, Surabaya, 31 pp. [in Indonesian]
- Bly F. E., Cuchens M. M., Clem L. W. 1986 Temperature mediated process in teleost immunity: binding and mitogenic properties of con-canavalin A. with channel catfish lymphocytes. Immunology 58:350-354.
- BSN (National Standard Board) 2009 [Nile tilapia production (*Oreochromis niloticus* Bleeker)]. Kelas Benih Sebar. BSN (Badan Standar Nasional). SNI 7550:2009, 12 pp. [in Indonesian]
- Cholik F., Jagatraya A. G., Poenormo R. P., Jauzi A., 2005 [Aquaculture: future national expectation]. Penerbit Masyarakat Nusantara dengan taman Akuarium Air Tawar, TMII, Jakarta, 94 pp. [in Indonesian]
- Cushnie T. P., Lamb A. J. 2005 Antimicrobial activity of flavonoids. International Journal of Antimicrobial Agents, 26, 343-356.
- Effendie M. I. 1997 [Fisheries biology]. Yayasan Pustaka Nusantara, Yogyakarta, 163 pp. [in Indonesian]
- Erlinger T. P., Platz E. A., Rifai N., Helzlsouer K. J., 2004 C-reactive protein and the risk of incident colorectal cancer. JAMA 291:585-590.

Commented [indra15]: the underlined references were already checked and corrected. So, do not make any changes on them.

Only address the comments on some other references

Commented [indra16]: write the full name

Commented [indra17]: translate in English

Commented [indra18]: not quoted in text

Commented [indra19]: write the full name

- FAO (Food and Agriculture Organization), 2008 Cultured aquatic species information programme *Penaeus monodon* (Fabricius, 1798). <http://www.fao.org/fishery/cultured-species/Penaeusmonodon/en>. 408 pp.
- Faulman E., Cuchens M. A., Lobb C. J., Miller N. W., Clem L. W., 1983 An effective culture system for studying in vitro mitogenic responses of channel catfish lymphocytes. *Transaction of the American Fisheries Society* 112:673-679.
- Finlay B., McFadden G., 2006 Antiimmunology: evasion of the host immune system by bacterial and viral pathogens. *Cell* 124(4):767-782.
- Galeotti M., 1998 Some aspects of the application of immunostimulants and a critical review of methods for their evaluation. *Journal of Applied Ichthyology* 14:189-199.
- Galindo-Villegas J., Hosokawa H., 2004 Immunostimulants: toward temporary prevention of diseases in marine fish. In: *Avances en Nutrición Acuícola VII. Cruz Suárez L. E., Ricque Marie D., Nieto López M. G., Villarreal D., Scholz U., González M., Memorias del VII Simposium Internacional de Nutrición Acuícola, 16-19 Noviembre, 2004, Hermosillo, Sonora, México, pp. 279-319.*
- Harikrishnan R., Balasundaram C., Heo M. S., 2010 Herbal supplementation diets on hematology and innate immunity in goldfish against *Aeromonas hydrophila*. *Fish and Shellfish Immunology* 27:508-515.
- Hendra R., Ahmad S., Sukari A., Shukor M. Y., Oskoueian E., 2011 Flavonoid analyses and antimicrobial activity of various parts of *Phaleria macrocarpa* (Scheff.) Boerl fruit. *Int J Mol Sci.* 12:3422-3431.
- Irianto A., 2004 *Patologi Ikan Teleostei*. Gadjah Mada University Press, Yogyakarta. [in Indonesian]
- Jun J. W., Kim J. H., Gomez D. K., Choresca C. H., Han J. E., Shin P. S., Park S. C., 2010 Occurrence of tetracycline-resistant *Aeromonas hydrophila* in Korean cyprinid loach (*Misgurnus anguillicaudatus*). *African Journal of Microbiology Research* 4(9):849-855.
- Kani R. A., Rantetondok A., Malina A. S., 2003 [The effect of CpG-ODN administration on the body immune system of brown-marbled grouper *Epinephelus fuscoguttatus*]. *Jurnal Pasca Universitas Hasanuddin*, 11(261): 8 pp. [in Indonesian]
- Karuthapandi M., Innocent X. B., 2010 Immune responses of *Tilapia mossambicus* administered with rat serum vaccine for *Vibrio anguillarum*. *Current Biotica* 4(1):63-72.
- Khairuman H., Amri K., 2012 [Rearing Nile tilapia in running water pond]. *Cetakan Pertama, PT Agro Media Pustaka, Jakarta*, 13-25 pp. [in Indonesian]
- Kozinska A., Guz L., 2004 The effect of various *Aeromonas bestiarum* vaccines on nonspecific immune parameters and protection of carp (*Cyprinus carpio* L). *Fish and Shellfish Immunology* 16:437-445.
- Kresno S. B., 2001 [Immunology: laboratory diagnosis and procedures]. Edisi Ketiga, Fakultas Kedokteran, Universitas Indonesia, Jakarta, 437 pp. [in Indonesian]
- Kusmardi, Kumala S., Triana E. E., 2007 [Immunomodulator effect of Chinese shrub (*Cassia alata* L.) leaf extract on macrophage phagocytosis activities and capacity]. *Jurnal Makara Kesehatan* 11(2):50-53. [in Indonesian]
- Labh S. N., Shyam N. L., Shakya S. R., 2014 Application of immunostimulants as an alternative to vaccines for health management in aquaculture. *International Journal of Fisheries and Aquatic Studies* 2(1):153-156.
- Li H., Wang Z., Liu Y., 2003 [Review in the studies on tannins activity of cancer prevention and anticancer]. *Zhong-Yao-Cai* 26(6):444-448. [in Chinese]
- Lou Z., Wang H., Rao S., Sun J., Ma C., Li J., 2012 P-coumaric acid kills bacteria through dual damage mechanisms. *Food Control* 25:550-554.
- Maftuch, 2007 [*Vibrio alginolyticus* exposure on the intestinal histology of humpback grouper (*Cromileptes altivelis*) and the development of number and macrophage cell activities]. *Jurnal Penelitian Perikanan* 10(1):66-70. [in Indonesian]
- Magnadottir B., 2006 Innate immunity of fish (overview). *Fish and Shellfish Immunology* 20:137-151.

Commented [indra20]: write the full name

Commented [indra21]: translate the title in English

Commented [indra22]: provide the pages too

- Maryani, Marsoedi, Nursyam, Maftuch, 2013 The phytochemistry and the anti-bacterial activity of yellow root (*Arcangelisia flava* Merr.) against *Aeromonas hydrophila*. Journal of Biology and Life Science 4:180-1.
- Mastan S. A., 2015 Use of immunostimulants in aquaculture disease management. International Journal of Fisheries and Aquatic Studies 2(4):277-280.
- Mishra S., Sahu N. P., Pal A. K., Xavier B., Mukherjee S. C., 2006 Pre and post challenge immuno-haematological changes in *L. rohita* juveniles fed gelatinized and non gelatinized carbohydrate with n-3 PUFA. Fish and Shellfish Immunology 21:346-356.
- Mishra V. S., Mishra A., Kumari R., Murthy P. N., Naik B. S., 2009 Antibacterial activity of ethanol extracts *Andrographis paniculata*. Fish and Shellfish Immunology 71(4):436-438.
- Naiborhu P. E., 2002 [Extraction and benefit of mangrove (*Sonneratia alba* and *Sonneratia caseolaris*) as natural antibacterial against pathogens of tiger prawns, *Vibrio harveyi*.] [Thesis], Sekolah Pascasarjana IPB, Bogor, 68 pp. [in Indonesian]
- Nuria, Cut M., Faizaitun, Arvin, Sumantri, 2009 Test on antibacterial activity of ethanol extract of *Jatropha* (*Jatropha curcas* L.) leaf ethanol extract against *Staphylococcus aureus* Atcc 25923, *Escherichia coli* Atcc 25922, and *Salmonella typhi* Atcc 1408. Mediagro 5(2):26-37.
- Park Y., Nho S. W., Shin G. W., Park S. B., Jang H. B., Cha I. S., Ha M. A., Kim Y. R., Dalvi R. S., Kang B. J., Jung T. S., 2009 Antibiotic susceptibility and resistance of *Streptococcus iniae* and *Streptococcus parauberis* isolated from olive flounder (*Paralichthys olivaceus*). Veterinary Microbiology 136:76-81.
- Rijayanti R. P., 2014 [Antibacterial activity test of ethanol extract of mango *Mangifera indica* L leaf against *Staphylococcus aureus* in vitro]. Disertasi, Fakultas Kedokteran Universitas Tanjungpura, Pontianak, 167 pp. [in Indonesian]
- Rorstad G., Asjord P. M., Robertsen B., 1993 Adjuvant effect of yeast glucan in vaccines against furunculosis in Atlantic salmon (*Salmo salar* L). Fish and Shellfish Immunology 3:179-190.
- Rustama M. M., Lingga M. A. 2005 Antibacterial activity test of ethanol extract of garlic (*Allium sativum* L.) against negative and positive gram bacteria isolated from penaeid shrimp *Metapenaeus monoceros*, lobster *Panulirus* sp., and mysis shrimp Rebon (*Mysis Acetes*). Jurnal Biotika 5(2):35-40. [in Indonesian]
- Sadikin H. M., 2002 [Blood biochemistry]. Penerbit Widya Medika, Jakarta, pp. 299-316. [in Indonesian]
- Sakai M., 1999 Current research status of fish immunostimulants. Aquaculture 172:63-92.
- Secombes C. J., 1990 Isolation of salmonids macrophages and analysis of their killing activity: In: Techniques in fish immunology. Stolen J. S., Fletcher T. C., Anderson D. P., Robersten B. S., Van Muiswinkel W. B. (eds), SOS Publications fair Haven, pp. 137-154.
- Selvaraj V., Sampath K., Sekar V., 2005 Administration of yeast glucan enhances survival and some specific and non-specific immune parameters in carp (*Cyprinus carpio*) infected with *Aeromonas hydrophila*. Fish and Shellfish Immunology 19:293-306.
- Shib S. S, Sukumaran V., Giri S. S., Park S. C., 2015 Flavonoid fraction of guava leaf extract attenuates lipopolysaccharide-induced inflammatory response via blocking of NF-κB signalling pathway in *Labeo rohita* macrophages. Fish and Shellfish Immunology 1(1):85-92.
- Smith C., 2007 Hematology and cell type differentiation. In: Fish histology and histopathology. Mufford V. (ed), USFWS-NCTC, 437 pp.
- Suhirman S., Winarti C., 2013 [Prospect and function of medicinal plants as immunomodulator]. Buletin Perkembangan Teknologi Tanaman Rempah dan Obat XIX No.2 Available from : http://www.readbag.com/balittro-litbang-deptan-go-id-ind-images-storiesedsus_vol19no2-4obat. [in Indonesian]

Commented [indra23]: is this the original title? Or the paper was written in Indonesian?

Commented [indra24]: what type of thesis: MSc, BSc, PhD...?

Commented [indra25]: is this the complete name?

Commented [indra26]: is this the original title? Or the paper was written in Indonesian?

Commented [indra27]: what kind of: PhD, MSc, or BSc?

Commented [indra28]: not quoted in text

Commented [indra29]: ???

- Susanti P. A., Trisunuwati P., Murwani S., 2012 [Effect of ethanol extract of cinnamon *Cinnamomum burmanii* on GR-1 development expressing IFN- γ and macrophage phagocytotic activity]. Fakultas Kedokteran Hewan Universitas Bratawijaya, Malang, pp. 1-9. [in Indonesian]
- Susilo J., 2012 [Immunomodulator effect of ethyl acetate fraction of dandelion *Taraxacum officinale* Weber et Wiggers leaf on non-specific immune response of male Balb/C rat]. STIKES Ngudi Waluyo, Ungaran, 89 pp.
- Tizard I., 1998 [An introduction to veterinary immunology]. Translated by Masduki and S. Hardjosworo, Pengantar Imunologi Veteriner, Universitas Airlangga, Surabaya, 197 pp. [in Indonesian]
- Ulfah, M. 2014 Immunostimulant activity test of acetate fraction of ethanol extract of Rosella *Hibiscus sabdariffa* L. petals on lymphocyte cell proliferation of Sweden strained rat *In vitro* and its chemical compound identification. Universitas Wahid Hasyim, Semarang, 78 pp. [in Indonesian]
- Vadstein O., 1997 The use of immunostimulation in marine larviculture: possibilities and challenges. *Aquaculture* 155:401-417.
- Watson R. R., Preedy V. R., 2013 Bioactive food as dietary interventions for arthritis and related inflammatory diseases. Academic Press, UK, 680 pp.
- Widya D. R., 2014 [Antimicrobial activity of nymphaea seed (*Nymphaea pubescens* L.) against bacteria *Aeromonas hydrophila*, *Streptococcus agalactiae*, and *Saprolegnia* sp.]. *Aquacoastmarine* 2(1):7-17. [in Indonesian]

Commented [indra30]: I suppose the paper is written in Indonesian. Am I right?

Commented [indra31]: is this a thesis? If yes, what kind of?

Commented [indra32]: write the full name

Received: 02 September 2017. Accepted: 31 October 2018. Published online: ... November 2018.

Authors:

Maryani, Aquaculture Program, Fisheries Department, Faculty of Agriculture, Palangkaraya University, Jl. Yos Sudarso Perum Dosen Muda UPR No. 99 Palangkaraya, Central Kalimantan, Indonesia, e-mail: maryani@fish.upr.ac.id

Sinta Silvia Monalisa, Aquaculture Program, Fisheries Department, Faculty of Agriculture, Palangkaraya University, Jl. Yos Sudarso Perum Dosen Muda UPR No. 99 Palangkaraya, Central Kalimantan, Indonesia, e-mail: mona212101@gmail.com

Rosita, Aquaculture Program, Fisheries Department, Faculty of Agriculture, Palangkaraya University, Jl. Yos Sudarso Perum Dosen Muda UPR No. 99 Palangkaraya, Central Kalimantan, Indonesia, e-mail: rosita.pribadi@yahoo.co.id

Mohamad Rozik, Aquaculture Program, Fisheries Department, Faculty of Agriculture, Palangkaraya University, Jl. Yos Sudarso Perum Dosen Muda UPR No. 99 Palangkaraya, Central Kalimantan, Indonesia, e-mail: rozi_raha@yahoo.com

Silvester Benny Pratasik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl.Kampus Bahu Manado-95115, North Sulawesi, Indonesia, e-mail: spjong07@yahoo.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Maryani, Monalisa S. S., Rosita, Rozik, Pratasik S. B., 2018 **Effectivity of *Arcangelisia flava* Merr as immunostimulant to prevent streptococcosis on Nile tilapia, *Oreochromis niloticus***. *AAFL Bioflux* 11(6): ...-...