

Phenolic content and antioxidant activities of five seaweeds from North Sulawesi, Indonesia

by Grace Sanger 1


Submission date: 07-Jan-2020 04:31PM (UTC+0700)

Submission ID: 1239738410

File name: Journal_Bioflux.pdf (604.83K)

Word count: 5267

Character count: 29097



58

Phenolic content and antioxidant activities of five seaweeds from North Sulawesi, Indonesia

¹Grace Sanger, ²Lexy K. Rarung, ¹Bertie E. Kaseger, ¹Jan R. Assa, ¹Agnes T. Agustin

41

¹ Department of Fishery Product Technology, Faculty of Fishery and Marine Science, Sam Ratulangi University, Manado, Indonesia; ² Department of Fishery Agribusiness, Faculty of Fishery and Marine Science, Sam Ratulangi University, Manado, Indonesia.

Corresponding author: G. Sanger: sanger.grace@yahoo.co.id

40

Abstract. This research aims to determine the composition of phenolic and antioxidant activity of five seaweeds (*Gracilaria salicornia*, *Halymenia durvylae*, *Halimeda macroloba*, *Turbinaria decurens* and *Sargassum olygocystum*) that were collected from North Sulawesi, Indonesia. Seaweeds were dried and extracts were prepared using the maceration method, with methanol solvent. The analyses performed consist of extraction yield, total phenolic content (TPC), scavenging radical activity of DPPH (1,1-diphenyl-2-picrylhydrazyl), ferric reducing antioxidant power (FRAP) and ferrous ion chelating (FIC) activity. The results of analysis present the highest content of TPC in *H. macroloba* (186.80±15.54 µg GAE (gallic acid equivalent) g⁻¹ extract). *T. decurens* had the highest radical DPPH scavenging activity (IC₅₀ 10.01±0.54 mg mL⁻¹). The highest value of FRAP was for *H. macroloba*, 28.52±1.46 µg GAE g⁻¹ extract. *S. olygocystum* and *T. decurens* exhibit higher values of FIC (IC₅₀ 5.18±0.21 and 7.02±0.43 mg mL⁻¹, respectively). The conclusion is that *G. salicornia*, *H. durvylae*, *H. macroloba*, *T. decurens* and *S. olygocystum* extracts are a possible source of phenolic compounds and natural antioxidants.

Key Words: algae, antioxidant activity, extracts, phenol.

Abstrak. Penelitian ini bertujuan untuk menghitung kandungan fenol dan aktifitas antioksidan lima jenis rumput laut (*Gracilaria salicornia*, *Halymenia durvylae*, *Halimeda macroloba*, *Turbinaria decurens* and *Sargassum olygocystum*), yang diambil dari perairan Sulawesi Utara Indonesia. Rumput laut dikeringkan kemudian diekstraksi menggunakan metanol dengan metoda maserasi. Jenis analisis penelitian ini terdiri dari: hasil ekstraksi, kandungan total fenol, aktifitas peredam radikal DPPH (1,1-diphenyl-2-picrylhydrazyl), daya reduksi dan aktifitas pengkelat ion. Hasil analisis menunjukkan bahwa kandungan total fenol tertinggi terdapat pada *H. macroloba* ((186.80±15.54 µg EAG (equivalent asam galat) g⁻¹ ekstrak). *T. decurens* mempunyai aktifitas antioksidan peredam radikal DPPH tertinggi, sebesar IC₅₀ 10.01±0.54 mg mL⁻¹). Nilai Daya reduksi tertinggi pada *H. macroloba* (28.52±1.46 µg EAG g⁻¹ ekstrak). Sedangkan *S. olygocystum* dan *T. decurens* mempunyai nilai pengkelat ion lebih tinggi, dengan nilai masing-masing (IC₅₀ 5.18±0.21 and 7.02±0.43 mg mL⁻¹). Kesimpulan penelitian ini menunjukkan bahwa *G. salicornia*, *H. durvylae*, *H. macroloba*, *T. decurens* and *S. olygocystum* mempunyai potensi sebagai sumber senyawa fenolik dan antioksidan alami.

Kata Kunci: rumput laut, fenol, aktifitas antioksidan, ekstrak.

Introduction. Seaweeds are beneficial to humans, environment and animal nutrition. They are used as fertilizers and soil acclimatizing substances in many countries (Anantharaman et al 2010; Robledo & Freil 1997). Seaweeds contain large amounts of nutrients and natural bioactive compounds, such as carotenoids, dietary fibers, proteins, essential fatty acids, vitamins and minerals (Devi et al 2008; Chandini et al 2008; Mantanjun et al 2008). These compounds are interesting for the pharmaceutical, medical, cosmetic, nutraceutical, food and agricultural industries (Boonchum et al 2011; Murugesan et al 2015; Kelman et al 2012).

Phenolic substances are generally obtained from edible red, green and brown seaweeds. The antioxidant characteristics of these seaweeds have been associated to their phenolic composition. Phenolic compounds or polyphenols have attained substantial attention due to their physiological purposes, counting antioxidant, antimutagenic, antitumor and anticancer activities (Souza et al 2011). Seaweeds could eliminate free

radicals by acting as free radical scavengers (Molyneux 2004) or by donating a hydrogen atom to the free radical (Re et al 1999).

All organisms have multifaceted regularities of antioxidant enzymes, for example thioredoxin enzyme. Some of these enzymes are preserved throughout growth and are needed for a normal development. Antioxidants in biological systems have various purposes, could be a counter for oxidative destruction and have a contribution in cell pathways. The most important action of antioxidants in cells is to inhibit the destruction caused by reactive oxygen groups (Halliwell et al 1992; Borek 1993). Reactive oxygen groups consist of hydrogen peroxide, superoxide anion and free radicals, such as hydroxyl. These molecules are changeable and extremely reactive, and can injure cells by chain reactions, like lipid peroxidation or creation of DNA adducts that could initiate cancer, endorsing mutations or cell mortality. Thus, to decrease or avoid these injuries, all cells constantly and consistently use antioxidants (Halliwell 1991; Aruoma 1993; Reaven & Witzum 1996).

Bioactive compounds from seaweeds possess a wide assortment of pharmacological properties, like anticancer, antibacterial, antifungal, anti-viral, anti-inflammatory, anticoagulant, antioxidant, bone regenerating, hepatoprotective and neuroprotective properties (Liu et al 2012; Chakraborty et al 2013).

Seaweed is one of the potential natural bioactive resources. In Indonesia, some types of macroalgae, such as *Gracilaria salicornia*, *Halymenia durvillae*, *Halimeda macroloba*, *Turbinaria decurens* and *Sargassum olygocystum* grow abundantly and have a high economic value for various industries. Any studies evaluating the antioxidant potential of the Indonesian seaweeds would increase their usefulness rate. Hence, the present study aims to examine the total phenolic content (TPC) and antioxidant ability of these five seaweeds.

Material and Method

Sample preparation and extraction. *G. salicornia*, *H. durvillae*, *H. macroloba*, *T. decurens* and *S. olygocystum* were collected from Nain Village, Manado, North Sulawesi. The samples were rinsed with fresh water to remove their holdfast, epiphytes, salt, debris and shellfish present. Afterwards, the samples were transported to the laboratory of Fisheries and Marine Science Faculty, Unsrat-Manado. Samples were shade-dried for two days and dried in an oven at 50°C for 3 days. After the drying process was complete, the samples were grounded to fine powder.

The samples were extracted using methanol (1:10 w/v) and were incubated overnight at room temperature. Extraction was repeated three times until the sample was colorless. The procedure was carried out in triplicates. The extracts were filtered and concentrated using a vacuum rotary evaporator (Buchi, Inggris) at 40°C. The extracts were stored in colored vial for future analysis.

Chemicals and reagents. Follin-Ciocalteu's phenol, gallic acid, sodium carbonate, methanol, ferric chloride hexahydrate, trichloroacetic acid (TCA), potassium ferricyanide, ferrous sulfate, vitamin C and butylated hydroxytoluene (BHT) were purchased from Merck (Spain). 1-diphenyl-2-picrylhydrazyl (DPPH), ferrozine iron reagent and phosphate buffer were purchased from Sigma-Aldrich (St. Louis, MO, USA). All other solvents and chemicals were of analytical grade.

Total phenolic content (TPC). The TPC of the extracts was determined using the Follin Ciocalteu reagent according to the method of Dev et al (2008), with a modification. Briefly, 75% of Follin Ciocalteu's phenol reagent (1 mL) was added to 5 mL of extract (0.1 g dry sample in 10 mL methanol) and vortexed. Next, 7% Na₂CO₃ (1 mL) was added and the reaction mixture was incubated at room temperature for 30 minutes. The absorbance was read at 750 nm using a spectrophotometer (Shimadzu type 1240, Japan). TPC was expressed as µg gallic acid equivalent (GAE) g⁻¹ dried extract.

DPPH radical scavenging assay. The DPPH-scavenging capacity of the extracts was measured based on the scavenging potential of stable 1,1-diphenyl-2-picrylhydrazyl (DPPH) radicals by the seaweeds antioxidants. The capacity of the extracts to scavenge the DPPH radical was determined by the method of Chew et al (2008) with a modification. Briefly, 2 mL of 0.93 µM DPPH in methanol was added to 0.5 mL extract of varying dilutions. The mixture was then vortexed and left in the dark for 20 minutes at room temperature. The absorbance was measured at 517 nm using a spectrophotometer (Hitachi type 1240, Japan). Vitamin C and BHT were used as control positive. Antioxidant activity was expressed as a percentage of the DPPH scavenging activity relative to the control, using the following equation:

$$\% \text{ radical scavenging activity} = \left(\frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \right) \times 100$$

Ferric reducing antioxidant power (FRAP). The seaweeds extract antioxidant activities at various dilutions were determined by using FRAP assay following the method of Kumar et al 2008, with modifications. 1 mL of 0.2 M phosphate buffer (pH 6.6) and 1 mL of K₃[Fe(CN)₆] (1%) were mixed with 1 mL of extracts. The reaction mixture was incubated at 50°C for 20 minutes, after which 1 mL of 10% TCA was added and centrifuged at 3000 rpm for 10 minutes. 1 mL of solution from the upper layer was mixed with 1 mL of sterile water and 0.5 mL of 0.1 % FeCl₃·6H₂O. The absorbance was measured at 700 nm. BHT was used as a positive control. The FRAP value was expressed as µg GAE g⁻¹.

Ferrous ion chelating (FIC) assay. The chelating of ferrous ions by extracts was investigated following the method described by Chew et al (2008). Briefly, 0.5 mL of seaweed extract at various concentrations, 0.5 mL of 0.1 mM FeSO₄ and 0.5 mL of 0.25 mM ferrozin ion reagent were mixed. The reaction mixtures were incubated for 20 minutes. The absorbance of the solution was measured at 562 nm. The percentage of inhibition of ferrozine-Fe²⁺ complex formation was determined using the following formula:

$$\% \text{ inhibition} = \left(\frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \right) \times 100$$

Statistical analysis. All experiments were conducted in triplicate. The means of parameters, extraction yields, TPC and antioxidant activity are presented as mean±standard deviation, using Microsoft Excel 2007.

Results and Discussion

Extraction yield. Table 1 shows the extraction yield of the five seaweed extracts on a dry basis weight. Among the methanol extract of the five seaweeds, *H. macroloba* had the highest yield extraction (18.45%±0.43), followed by *H. durvillae* (6.34%±0.43), *T. decurens* (5.72%±0.23), *S. olygocystum* (4.95%±0.03) and *G. salicornia* (3.56%±0.12), respectively.

Table 1
Extraction yield of seaweed extracts on a dry weight basis

Seaweeds	Division	Yields (%) w/w
<i>Gracilaria salicornia</i>	Rhodophyta	3.56±0.12
<i>Halymenia durvillae</i>	Rhodophyta	6.34±0.43
<i>Halimeda macroloba</i>	Chlorophyta	18.45±0.43
<i>Turbimaria decurens</i>	Phaeoephyta	5.72±0.23
<i>Sargassum olygocystum</i>	Phaeoephyta	4.95±0.03

Total phenolic content. TPC of methanol extracts for the five seaweeds are presented in Table 2. The results show that *H. macroloba* exhibited the highest amount of TPC ($186.80 \pm 15.84 \mu\text{g GAE g}^{-1}$ extract), whereas the lowest TPC was for *G. salicornia* ($14.73 \pm 1.04 \mu\text{g GAE g}^{-1}$ extract).

25

Table 2

Total phenolic content (TPC) of seaweed extracts

Seaweeds	TPC ($\mu\text{g GAE g}^{-1}$ dry extract)
<i>Gracilaria salicornia</i>	14.73 ± 1.04
<i>Halymenia durvilae</i>	27.60 ± 2.37
<i>Halimeda macroloba</i>	186.80 ± 15.54
<i>Turbinaria decurens</i>	77.20 ± 5.43
<i>Sargassum olygocystum</i>	84.67 ± 6.89

Note: GAE - gallic acid equivalent.

24

DPPH radical scavenging activity. DPPH is a stable nitrogen free radical, which can be successfully scavenged by antioxidants. DPPH have been used comprehensively as a free radical to calculate the activity of antioxidants from plant extracts. A newly ready DPPH solution displays an intense purple color. This purple color usually weakens/vanishes while an antioxidant is present in the solution. Hence, antioxidant compounds can scavenge DPPH free radicals and transform them in a colorless product. Therefore, the faster is the decline of the absorbance, the more strong the antioxidant activity is (Seenivasan et al 2013). The capacity of the extract to scavenge DPPH radicals was established by the decline of absorbance at 517 nm (Molyneux 2004). Figure 1 shows the DPPH scavenging activity of the five seaweed extracts. The most effective activity was presented by *T. decurens*, followed by *G. salicornia*, *H. durvilae*, *S. olygocystum* and *H. macroloba*, with the following IC_{50} values: 10.01 ± 0.53 ; 12.81 ± 0.93 ; 14.17 ± 1.06 ; 15.38 ± 1.17 ; $18.54 \pm 1.25 \text{ mg mL}^{-1}$, respectively.

19

Ferric reducing antioxidant power. The reducing capability is deliberated as an important indicator of potential antioxidant activity of a constituent or a sample. The attendance of reductants (antioxidants) instigates the reduction of the Fe^{3+} /ferricyanide multifarious to the ferrous type. Consequently, by evaluating the establishment of Perl's Prussian blue at 255 nm, the quantity of Fe^{2+} can be observed (Seenivasan et al 2013). Table 3 shows the antioxidant activity of the extracts using FRAP assay. The highest reducing power was observed in the case of *H. macroloba*, followed by *S. olygocystum*, *T. decurens*, *G. salicornia* and *H. durvilae*, with the following values: 28.52 ± 1.46 ; 19.95 ± 1.72 ; 16.77 ± 1.47 ; 7.24 ± 0.61 and $5.39 \pm 0.341 \mu\text{g GAE g}^{-1}$ extract, respectively. The positive control, BHT, showed higher antioxidant activity than all the seaweed samples.

Ferrous ion-chelating activity. *G. salicornia*, *H. durvilae*, *H. macroloba*, *T. decurens* and *S. olygocystum* extracts have ferrous ion-chelating abilities, as presented in Figure 2. The ability of all the five species of seaweeds increased with the concentrations, which increased from 2.5 to 10 mg mL^{-1} . The extracts with a higher ferrous ion-chelating ability were from brown algae *S. olygocystum* and *T. decurens*, followed by *G. salicornia*, *H. durvilae* and *H. macroloba*, the IC_{50} values being 5.18 ± 0.21 , 7.02 ± 0.43 , 13.12 ± 0.54 , 14.39 ± 1.34 and $34.25 \pm 2.78 \text{ mg mL}^{-1}$, respectively.




Figure 1. DPPH radical scavenging activity (%) of seaweed extracts.

The methanol yields of extraction for the seaweeds were different, the same as in other cases (Chew et al 2008; Chandini et al 2008; Ganesan et al 2008). Mantanjun et al (2008) detected a yield of total methanol extract for red algae, *Euchema spinosum*, of 1.88%, for green algae, *Caulepa lentillifera*, of 30.86% and brown algae, *Dictyota dichotoma*, of 40.33%. The difference in the yields of various extracts is ascribed to the polarities of different constituents in the plants.

Table 3

Ferric reducing antioxidant power (FRAP) of seaweeds extracts

Species	FRAP ($\mu\text{g GAE g}^{-1}$ extract)
<i>Gracilaria salicornia</i>	7.24±0.61
<i>Halymenia durvillae</i>	5.39±0.34
<i>Halimeda macroloba</i>	28.52±1.46
<i>Turbinaria decurrens</i>	16.77±1.47
<i>Sargassum oligocystum</i>	19.95±1.72
BHT	41.32±3.87

Note: GAE - gallic acid equivalent.

Halimeda spp. have been examined over the past years as sources of high phenolic compounds with biological activity of diverse types (Table 4). The antioxidant properties of phenols are an effect of their capability to work as reducing agents, hydrogen donors and free radical quenchers. Phenols can also work as metal chelators, which avoid the catalytic purpose of metal in the process of instigating radicals (Devi et al 2008). Many hydrophilic polyphenolic compounds were observed in seaweeds, such as epigallocatechin gallate, epicatechin and phlorotannins (Boonchum et al 2011). *H. macroloba* contains high levels of polyphenolic compounds, up to 28000 μg of epigallocatechin and 1880 μg of catechol. It also presents caffeic acid and hesperidin (oshie et al 2002).

Some studies fervently sustain an involvement of polyphenols in the avoidance of cardiovascular diseases, cancer inflammation and osteoporosis (Das et al 2012). Moreover, polyphenols seem to have a function in the hindrance of neurodegenerative diseases and diabetes mellitus (Scalbert et al 2005). Most likely, cells react to polyphenols primarily over straight contacts with receptors or enzymes implicated in motion transduction, which may produce consequences in the alteration of the redox position of the cell and may activate a sequence of redox-reliant reactions. Antioxidant and prooxidant impressions of polyphenols have been explained with dissimilarity effects

on cell physiologic developments. As antioxidants, polyphenols may recover cell endurance; as prooxidants, they may generate apoptosis and avoid tumor growth (Lambert et al 2005).




Figure 2. Ferrous Ion chelating activity (% inhibition) of the seaweed extracts.

The five species of seaweeds, especially *T. decurens*, show high antioxidant activity as determined by the DPPH assay. Polyphenols are forceful antioxidant components (Boonchum et al 2011). However, the antioxidant activities were not the effect of just phenolic compounds. For example, *T. decurens* had low phenolic content, but presented high antioxidant activity. As it was explained by Kuda & Ikemori (2009), the activities may be initiated by other hydrophilic compounds, for instance peptides, fucoidan and Maillard reaction products and it may be correlated to polysaccharides and pigments (Chattopadhyay et al 2010). Sanger et al (2018) reported that the TPC of ethanol extracts of *T. decurens* was 60.79 ± 4.57 $\mu\text{g GAE g}^{-1}$ extract, the carotene content was 6.948 ± 0.654 $\mu\text{g g}^{-1}$ dry weight and the scavenging radical activity of DPPH was IC_{50} 0.3033 ± 0.023 mg mL^{-1} . According to Chattopadhyay et al (2010), the sulfated polysaccharides responsible for the antioxidant activity of *T. conoides* have been isolated and identified as fucoidan, laminaran, and alginate. The major antioxidants isolated from the Hawaiian specimen of *T. ornata* were carotenoid and fucoxanthin (Kelman et al 2012). Crude and purified sulfated polysaccharides from *T. ornata* and *T. conoides* have been shown to be excellent antioxidants and have been suggested as natural antioxidant agents (Chattopadhyay et al 2010) for using as food supplements to raise the shelf-life of food products and nutraceuticals against oxidative stress diseases (Chakraborty et al 2013). Seaweed polysaccharides have an essential function as free radical scavengers (*in vitro*) and antioxidants in the deterrence of oxidative destruction in live organisms (Yuan et al 2005). The crude polysaccharide of *T. ornata* could be deliberated as a prospective antioxidant and anti-inflammatory substances (Ananthi et al 2010).

The reducing capacity properties show that the antioxidant constituents are electron donors and can decrease the oxidized intermediates of the lipid peroxidation process, so they can act as primary and secondary antioxidants. In terms of antioxidant activity, *H. macroloba*, *T. decurens* and *S. olygocystum* are more active than the red algae *G. salicornia* and *H. durvilae*. Boonchum et al (2011) reported that the reducing power of aqueous extracts from *H. macroloba* (1 mg mL^{-1}) was $0.092 \text{ mg gallic acid g}^{-1}$ dry weight and its TPC was $0.077 \pm 0.001 \text{ mg gallic acid g}^{-1}$ dry weight. *Halimeda* sp. possesses high phenolic content and low quantities of other antioxidants, like ascorbate, β -carotene, chlorophylls and selenium. Phenolic acids are the most important constituents of *Halimeda* spp., the substantial antioxidant activity being exhibited by *Halimeda* spp. extracts, especially by the phenolic acids (salicylic, cinamic, gallic

pyrogallic, ferulic and caffeic acids). The genus *Halimeda* contains phytopharmaceuticals and possess pharmacological properties, such as antimicrobial activity, stimulation of apoptosis, anti-trichomonal and anti-inflammatory activity; it is also valuable in biomedicine, in hepato, neuro and athero protection (Silva et al 2017).

2 Table 4
Total phenolic content (TPC) and biological activity in some species of the genus *Halimeda*

Species	Antioxidant activity	TPC	Reference
53 <i>H. macroloba</i>	DPPH IC ₅₀ 18.54±1.25 mg mL ⁻¹ FIC 34.25±2.78 mg mL ⁻¹	186.80±15.54 µg GAE g ⁻¹ extract	This study
<i>H. macroloba</i>	DPPH (10 mg mL ⁻¹) 57.37±3.13% FRAP 14.89±2.78 g GAE/100 g extract	18.42±0.65 g GAE/100 g extract (70% methanol extract)	14 Sanger et al (2013)
<i>H. macroloba</i>	DPPH 1.0212±0.044 mg mL ⁻¹ FRAP 0.248±0.014 µM Fe ₂ +mg ⁻¹	62.78±5.38 µg GAE g ⁻¹ acetone extract	Sanger et al (2018)
<i>H. macroloba</i>	DPPH 1.9776±0.1402 FRAP 0.214±0.013 µM Fe ₂ +mg ⁻¹	11.58±1.032 µg GAE g ⁻¹ ethanol extract	Sanger et al (2018)
<i>H. macroloba</i>	Anti-lipid IC ₅₀ 155.590±16.129 mg mL ⁻¹ peroxidation activity 0.018 ±0.002b mg trolox g ⁻¹ dry weight	0.085±0.003 mg gallic acid/g dry weight	Boonchum et al (2011)
<i>H. opuntia</i>	β-carotene-linoleic acid system (20 µg phenols) 73.5% DPPH (THF extract) IC ₅₀ 12.8-15.2 mg phenolic compounds	74.3 mg/g dry weight seaweed	Silva et al (2017)
<i>H. monile</i>	β-carotene-linoleic acid system 74.4% (20 µg phenols) IC ₅₀ 7.7-13.2 mg phenolic compounds	66.7mg/g dry weight seaweed	Silva et al (2017)
<i>H. incrassata</i>	DPPH 19-53% Inhib. oxidation-Cu LDL 0.87±0.09mg/mL Inhib. oxidation-AAPH-LDL 0.16±0.01 mg/mL	10-40 µg polyphenolics	Costa-Mugica (2012)
<i>H. incrassata</i>	β-carotene-linoleic acid 95% (10 µg polyphenols)	255 µg/g fresh seaweed	Vidal et al (2011)

31 e: TPC - total phenolic content; DPPH - 1,1-diphenyl-2-picrylhydrazyl; FIC - ferrous ion chelating activity; GAE - gallic acid equivalent; FRAP - ferric reducing antioxidant power; THF - tetrahydrofuran.

The binding of the antioxidant constituents to metal ions was estimated using the FIC assay. An extract with higher binding activity would avoid or restrain reactions, like Fenton's reaction, which produces reactive hydroxyl radicals. In this study, brown algae (*S. Olygocystum* and *T. decurens*) had higher binding activities than red algae (*G. salicornia* and *H. durvilae*) and green algae (*H. macroloba*). As it was shown by Chew et al (2008), brown algae *Padina antillarum* had the highest chelating ability compared with red algae *Kappaphycus alvarezzie* and green algae *Caulerpa racemosa*. Phlorotannins, commonly found in brown seaweeds, are powerful chelators of heavy metals (Toth & Pavia 2000; Sathya et al 2013). In contrast, the TPC values of *S. olygocystum* and *T. decurens* were lower than that of *H. macroloba*. According to Wang et al (2009) and Chin et al (2015), phlorotannins did not seem to be extremely effective metal chelators, hence indicating the existence of some compounds other than phenols. Polysaccharides, pigments, proteins or peptides in the extracts also have the capacity to

chelate metal ions (Chakraborty et al 2013). Polysaccharides (alginates, fucoidan) and phytochelatin are more valuable than phlorotannins regarding detoxification and copper accumulation in *Ascophyllum nodosum* (Wang et al 2009; Chin et al 2012). Actually, compounds counting phenolic acids, flavonoid quercetin and phenolic glycosides are also noticed to chelate transition metal ions like Fe²⁺. These active compounds might have a synergistic effect, contributing with an essential function in antioxidant scavenging activities of seaweed extracts (Kuda & Ikemori 2009).

Conclusions. *G. salicornia*, *H. durvillae*, *H. macroloba*, *T. decurens* and *S. olygocystum* extracts can be used as a source of antioxidants. The highest content of TPC was in *H. macroloba* (186.80±15.54 µg GAE g⁻¹ extract). *T. decurens* had the highest radical DPPH scavenging activity (IC₅₀ 10.01±0.54 mg mL⁻¹). The highest value of FRAP was *H. macroloba*, 28.52±1.459 µg GAE g⁻¹ extract. *S. olygocystum* and *T. decurens* exhibit a strong FIC ability, with IC₅₀ 5.18±0.21 and 7.02±0.43 mg mL⁻¹ values, respectively.

Acknowledgements. The authors are grateful to Prof. Dr. Ir. Remmy Mangindaan M.Sc., from Sam Ratulangi University Manado and Dr. Ir. Dikdik Kurnia, M.Sc, Ms., from Padjadjaran University, Bandung, for their constant support and for providing the research facilities.

References

- Anantharaman P., Karthikaidevi G., Manivannan K., Thirumaran G., Balasubramanian T., 2010 Mineral Composition of marine macroalgae from Mandapam Coastal Regions; Southeast Coast Of India. *Recent Research in Science and Technology* 2(10):66-71.
- Ananthi S., Raghavendran H. R. B., Sunil A. G., Gayathri V., Ramakrishnan G., Vasanthi H. R., 2010 *In vitro* antioxidant and *in vivo* anti-inflammatory potential of crude polysaccharide from *Turbinaria ornata* (marine brown alga). *Food Chemical Toxicology* 48(10):187-192.
- Aruoma I. O., 1993 Antioxidant action of plant foods: use of oxidative DNA damage, as a tool for studying antioxidant efficacy. *Free Radical Research* 30:419-427.
- Boonchum W. Y., Peerapornpisal P., Vacharapiyasophon J., Pekkoh C., Pumas U., Jamjai D., Amornlerdpison T., Noiraksar T., Kanjanapothi D., 2011 Antioxidant activity of some seaweed from the Gulf of Thailand. *International Journal Agriculture Biology* 13:95-99.
- Borek C., 1993 Molecular mechanisms in cancer induction and prevention. *Environmental Health Perspectives* 101(Supplement 3):237-245.
- Chakraborty K., Praveen N. K., Vijayan K. K., Rao G. S., 2013 Evaluation of phenolic contents and antioxidant activities of brown seaweeds belonging to *Turbinaria* spp. (Phaeophyta, Sargassaceae) collected from Gulf of Mannar. *Asian Pacific Journal of Tropical Biomedicine* 3(1):8-16.
- Chandini S. K., Ganesan P., Bashar N., 2008 *In vitro* antioxidant activities of three selected brown seaweeds of India. *Food Chemistry* (107):707-713.
- Chattopadhyay N., Ghosh T., Sinha S., Chattopadhyay K., Karmakar P., Ray B., 2010 Polysaccharides from *Turbinaria conoides*: Structural features and antioxidant capacity. *Food Chemistry* 118:823-829.
- Chew Y. L., Lim Y. Y., Omar M., Khoo K. S., 2008 Antioxidant activity of three Edible seaweeds from two areas in South East Asia. *Food Science and Technology* 41(6):1067-1072.
- Chin Y. X., Lim P. E., Maggs C. A., Phang S. M., Sharifuddin Y., Brian D., Green B. D., 2015 Anti-diabetic potential of selected Malaysian seaweeds. *Journal of Applied Phycology* 27(5):2137-2148.
- Costa-Mugica A., Batista-Gonzalez A. E., Mondejar D., Soto-López Y., Brito-Navarro V., Vázquez A. M., Brömme D., Claudina Zaldívar-Muñoz C., Alexis Vidal-Novoa A., Oliveira e Silva A. M., Mancini-Filho J., 2012 Inhibition of LDL-oxidation and antioxidant properties related to polyphenols content of hydrophilic fractions from

- seaweed *Halimeda incrassata* (Ellis) Lamouroux. Brazilian Journal of Pharmaceutical Sciences 48(1):31-37.
- Das L., Bhaumik E., Raychaudhuri U., Chakraborty R., 2012 Role of nutraceuticals in human health. Journal of Food Science and Technology 49(2):173-183.
- Devi G. K., Manivannan K., Thirumaran G., Rajathi F. A. A., Anantharaman P., 2011 *In vitro* antioxidant activities of selected seaweeds from Southeast Coast of India Asian Pacific Journal of Tropical Medicine 4(3):205-211.
- Devi K. P., Suganthi N., Kesika P., Pandian S. K., 2008 Bioprotective properties of seaweeds: *In vitro* evaluation of antioxidant activity and antimicrobial activity against food borne bacteria in relation to polyphenolic content. BMC Complementary and Alternative Medicine 8(3):882-888.
- Ganesan P., Kumar C. S., Baskar N., 2008 Antioxidant properties of methanol extract and its solvent fraction obtained from selected index red sea weeds. Bioresources Technology 99:2717-2723.
- Halliwell B., 1991 Reactive oxygen species in living systems: source, biochemistry, and role in human disease. American Journal of Medicine 91(3C):14S-22S.
- Halliwell B., Gutteridge J. M. C., Cross C. E., 1992 Free radicals, antioxidants and human disease: where are we now? Journal of Laboratory and Clinical Medicine 119:598-620.
- Kelman D., Posner E. K., McDermid K. J., Tabandera N. K., Wright P. R., Wright A. D. 2012 Antioxidant Activity of Hawaiian Marine Algae. Marine Drugs 10(2):403-416.
- Kuda T., Ikemori T., 2009 Minerals, polysaccharides and antioxidant properties of aqueous solutions obtained from macro alga beach-casts in the Noto Peninsula, Ishikawa, Japan. Food Chemistry 112(3):575-581.
- Kumar K. S., Ganesan K., Subba-Rao P. V., 2008 Antioxidant potential of solvent extracts of *Kappaphycus Alvarezii* (Doty) Doty-an edible seaweed. Food Chemistry 107:289-295.
- Lambert J. D., Hong J., Yang G., Liao J., Yang C. S., 2005 Inhibition of carcinogenesis by polyphenols: evidence from laboratory investigations. American Journal of Clinical Nutrition 81(1 suppl):284S-91S.
- Liu L., Heinrich M., Myers S., Dworjanyn S. A., 2012 Towards a better understanding of medicinal uses of the brown seaweed *Sargassum* in traditional Chinese medicine: A phytochemical and pharmacological review. Journal Ethnopharmacology 142(3):591-619.
- Matanjun P., Mohamed S., Mustapha N. M., Muhammad K., Ming C. H., 2008 Antioxidant activities and phenolics content of eight species of seaweeds from North Borneo. Journal of Applied Phycology 20(4):367-373.
- Molyneux P., 2004 The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. Songklanakarin Journal of Science and Technology 26(2):211-219.
- Murugesan S., Bhuvaneswari S., Selvan T., 2015 Evaluation of antioxidant property of methanolic extract of *red Chondrococcus hornemannii* and *Spyridia fusiformis*. Journal of Chemical and Pharmaceutical Research 7(1):333-337.
- Re R. N., Pellegrini A., Proteggente A., Pannala M., Yang C., Rice-Evans C., 1999 Antioxidant activity applying an improved ABTS radical cation decolorization assay. Free Radical Biology and Medicine 26:1231-1237.
- Reaven P. D., Witzum J. L., 1996 Oxidised LDL in atherogenesis. Role of dietary modification. Annual Review of Nutrition 16:51-71.
- Robledo D., Freile-Pelegrin Y., 1997 Chemical and mineral composition of six potentially edible seaweed species of Yucatan. Journal of Botanica Marina 40:301-306.
- Sanger G., Kaseger B. E., Rarung L. K., Damongilala L. K., 2018 [The potential of seaweeds as a functional food source of natural pigments and antioxidants]. Jurnal Pengolahan Hasil Perikanan Indonesia 21(2):208-218. [in Indonesian].
- Sanger G., Widjanarko S. B., Kusnadi J., Berhimpon S., 2013 Antioxidant Activity of Methanol Extract of Seaweeds Obtained from North Sulawesi. Food Science and Quality Management 19:63-70.

- Sathya R., Kanaga N., Sankar P., Jeeva S., 2013 Antioxidant properties of phlorotannins from brown seaweed *Cystoseira trinodis* (Forsskal) C. Agardh. *Arabian Journal of Chemistry* 10(2) Supplement S2608-S2614.
- Scalbert A., Johnson I. T., Saltmarsh M., 2005 Polyphenols: Antioxidant and beyond. *The American Journal of Clinical Nutrition* 81(1):215S-217S.
- Seenivasan I. H. R., 2013 *In vitro* antioxidant activity of selected seaweeds from Southeast Coast of India. *International Journal of Pharmacy and Pharmaceutical Sciences* 5(2):474-484.
- Silva A. M. O., Novoa A. V., Gutierrez D. D., Mancini-Filho J., 2017 Seaweeds from *Halimeda* Genus as sources of natural antioxidants. *Journal Analytical & Pharmaceutical Research* 5(6):1-5.
- Souza B. W. S., Miguel A., Cerqueira M. A., Joana T., Martins J. T., Mafalda A. C., Quintas M. A. C., 2011 Antioxidant potential of two red seaweeds from Brazilian Coasts. *Journal of Agricultural and Food Chemistry* 59(1):5589-5594.
- Toth G., Pavia A., 2000 Lack of phlorotannins induction in the *Ascophylum nodosum* in response to increased copper concentration. *Marine Ecology Progress Series* 192:119-126.
- Vidal A., Silva de Andrade-Wartha E. R., Fallarero A.L., de Oliveira A. M.S, Silva A.M, Ines M.G, Gonsales A.E. Vuorela P., Costa A, Malcino M-Filho 2011 Antioxidant activity and bioactive components in hydrophilic and lipophilic fractions from the seaweed *Halimeda incrassata*. *Revista Brasileira de Farmacognosia* 21(1):53-57.
- Wang T., Jónsdóttir R., Ólafsdóttir G., 2009 Total phenolic compounds, radical scavenging and metal chelating of extracts from Icelandic seaweeds. *Food chemistry* 116(1):240-248.
- Yoshie Y., Wand W., Hsieh Y. P., Suzuki T., 2002 Compositional Difference of Phenolic Compounds between two Seaweeds, *Halimeda* spp. *Journal of Tokyo University of Fisheries* 88:21-24.
- Yuan H., Zang W., Li X., Lu X., Li N., Gao X. Song J., 2005 Preparation and in vitro antioxidant activity of α -carrageenan oligosaccharides and their oversulfated, acetylated and phosphorylated derivatives. *Carbohydrate Research* 340(4):685-693.

Received: 24 March 2019. Accepted: 15 May 2019. Published online: 1 November 2019.

Authors:

Grace Sanger, Department of Fishery Product Technology, Faculty of Fishery and Marine Science, Sam Ratulangi University, 95115 Manado, Indonesia, e-mail: sanger.grace@yahoo.co.id

Lexy Karel Rarung, Department of Fishery Agribusiness, Sam Ratulangi University, 95115 Manado, Indonesia, e-mail: lexyrarung@unsrat.ac.id

Bertie Elias Kaseger, Department of Fishery product Technology, Faculty of Fishery and Marine Science, Sam Ratulangi University, 95115 Manado, Indonesia, e-mail: bertiekaseger@gmail.com

Jan Rudolf Assa, Department of Agriculture Technology, Faculty of Agriculture, Sam Ratulangi University, 95115 Manado, Indonesia, e-mail: assayance@yahoo.co.id

Agnes Triasin Agustin, Department of Fishery Product Technology, Faculty of Fishery and Marine Science, Sam Ratulangi University, 95115 Manado, Indonesia, e-mail: ata170855@yahoo.com

How to cite this article:

Sanger G., Rarung L. K., Kaseger B. E., Assa J. R., Agustin A. T., 2019 Phenolic content and antioxidant activities of five seaweeds from North Sulawesi, Indonesia. *AAFL Bioflux* 12(6):2041-2050.

Phenolic content and antioxidant activities of five seaweeds from North Sulawesi, Indonesia

ORIGINALITY REPORT

17%

SIMILARITY INDEX

11%

INTERNET SOURCES

12%

PUBLICATIONS

12%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Oktay, M.. "Determination of in vitro antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts", LWT - Food Science and Technology, 200303
Publication 1%
- 2 www.science.gov
Internet Source 1%
- 3 www.j3.jstage.jst.go.jp
Internet Source 1%
- 4 biotech-health.com
Internet Source <1%
- 5 citeseerx.ist.psu.edu
Internet Source <1%
- 6 Marta Mesías, Marta Navarro, Vural Gökmen, Francisco J Morales. "Antiglycative effect of fruit and vegetable seed extracts: inhibition of AGE formation and carbonyl-trapping abilities", Journal of the Science of Food and Agriculture, <1%

2013

Publication

-
- | | | |
|-------|--|-----|
| 7 | Abad, Lucille V., Lorna S. Relve, Charles Darwin T. Racadio, Charito T. Aranilla, and Alumanda M. De la Rosa. "Antioxidant activity potential of gamma irradiated carrageenan", <i>Applied Radiation and Isotopes</i> , 2013. | <1% |
| <hr/> | | |
| 8 | www.eckloniacava.com
Internet Source | <1% |
| <hr/> | | |
| 9 | L, Aranganathan, Suman TY, Remya RR, Gayathri S, and Radhika Rajasree S. R. "Enhanced cytotoxic activity of AgNPs on retinoblastoma Y79 cell lines synthesized using marine seaweed <i>Turbinaria ornata</i> ", <i>IET Nanobiotechnology</i> , 2016. | <1% |
| <hr/> | | |
| 10 | www.scribd.com
Internet Source | <1% |
| <hr/> | | |
| 11 | repositorium.sdum.uminho.pt
Internet Source | <1% |
| <hr/> | | |
| 12 | pdfs.semanticscholar.org
Internet Source | <1% |
| <hr/> | | |
| 13 | How Y Lai, Yau Y Lim, Kah H Kim. "Blechnum Orientale Linn - a fern with potential as antioxidant, anticancer and antibacterial agent", | <1% |

BMC Complementary and Alternative Medicine, 2010

Publication

14

Submitted to Imperial College of Science, Technology and Medicine

Student Paper

<1%

15

E. Conde. "Extraction of natural antioxidants from plant foods", Separation extraction and concentration processes in the food beverage and nutraceutical industries, 2010

Publication

<1%

16

myais.fsktm.um.edu.my

Internet Source

<1%

17

Submitted to Universiti Teknologi MARA

Student Paper

<1%

18

www.jmbfs.org

Internet Source

<1%

19

Submitted to Higher Education Commission Pakistan

Student Paper

<1%

20

Submitted to Kyungpook National University

Student Paper

<1%

21

Submitted to University of Newcastle upon Tyne

Student Paper

<1%

22

www.journalajst.com

Internet Source

<1%

23

Niu, Yuge, Pingping Shang, Lei Chen, Hua Zhang, Lu Gong, Xiaowei Zhang, Wenjuan Yu, Yuhong Xu, Qin Wang, and Liangli (Lucy) Yu. "Characterization of a Novel Alkali-Soluble Heteropolysaccharide from Tetraploid *Gynostemma pentaphyllum* Makino and Its Potential Anti-inflammatory and Antioxidant Properties", *Journal of Agricultural and Food Chemistry*

Publication

<1%

24

www.maxwellsci.com

Internet Source

<1%

25

Guinea, M.. "In vivo UVB-photoprotective activity of extracts from commercial marine macroalgae", *Food and Chemical Toxicology*, 201203/04

Publication

<1%

26

studentsrepo.um.edu.my

Internet Source

<1%

27

eprints.fbme.utm.my

Internet Source

<1%

28

Hemlatha Nagappan, Poh Ping Pee, Sandra Hui Yin Kee, Ji Tsong Ow, See Wan Yan, Lye Yee Chew, Kin Weng Kong. "Malaysian brown

<1%

seaweeds *Sargassum siliquosum* and *Sargassum polycystum* : Low density lipoprotein (LDL) oxidation, angiotensin converting enzyme (ACE), α -amylase, and α -glucosidase inhibition activities", Food Research International, 2017

Publication

29

Lamia Mhadhebi. "Anti-Inflammatory and Antiproliferative Activities of Organic Fractions from the Mediterranean Brown Seaweed, *Cystoseira Compressa* : Anti-inflammatory and antiproliferative activities of *C. Compressa*", Drug Development Research, 03/2012

Publication

30

Muhammad Nursid, Nurrahmi Dewi Fajarningsih, Ekowati Chasanah. "CYTOTOXIC ACTIVITY AND APOPTOSIS INDUCTION OF T47D CELL LINES BY *Turbinaria decurrens* EXTRACT", Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology, 2013

Publication

31

bashanfoundation.org

Internet Source

32

www.ijpbs.com

Internet Source

33

Submitted to Jawaharlal Nehru University (JNU)

Student Paper

<1%

<1%

<1%

<1%

<1%

34

Hajime Inoue, Ken-Ichi Hisamatsu, Miki Yamauchi, Norio Kumagai. "Rapid and simple determination of histamine-N-methyl transferase activity by high-performance liquid chromatography with UV detection", Mediators of Inflammation, 2001

Publication

<1%

35

Li-Jun Wang, Dong Li, Lei Zou, Xiao Dong Chen, Yong-Qiang Cheng, Koji Yamaki, Li-Te Li. "Antioxidative Activity of Douchi (A Chinese Traditional Salt-Fermented Soybean Food) Extracts During Its Processing", International Journal of Food Properties, 2007

Publication

<1%

36

Submitted to National University of Singapore

Student Paper

<1%

37

Submitted to University of Salford

Student Paper

<1%

38

arrow.dit.ie

Internet Source

<1%

39

ijpjournal.com

Internet Source

<1%

40

worldwidescience.org

Internet Source

<1%

41

docslide.us

Internet Source

<1%

42 ulir.ul.ie Internet Source <1%

43 repository.wit.ie Internet Source <1%

44 www.valacta.com Internet Source <1%

45 RAJAURIA, GAURAV, AMIT KUMAR JAISWAL, NISSREEN ABU-GANNAM, and SHILPI GUPTA. "ANTIMICROBIAL, ANTIOXIDANT AND FREE RADICAL-SCAVENGING CAPACITY OF BROWN SEAWEED HIMANTHALIA ELONGATA FROM WESTERN COAST OF IRELAND : ANTIMICROBIAL AND ANTIOXIDANT PROPERTIES OF IRISH SEAWEED", Journal of Food Biochemistry, 2013.
Publication

46 O'Sullivan, Anthony M, Yvonne C O'Callaghan, Michael N O'Grady, David S Waldron, Thomas J Smyth, Nora M O'Brien, and Joseph P Kerry. "An examination of the potential of seaweed extracts as functional ingredients in milk", International Journal of Dairy Technology, 2014.
Publication

47 flex.flinders.edu.au Internet Source <1%

48

darshanpublishers.com

Internet Source

<1%

49

Submitted to Universidade de Sao Paulo

Student Paper

<1%

50

Kuda, T.. "Antioxidant properties of dried product of 'haba-nori', an edible brown alga, *Petalonia binghamiae* (J. Agaradh) Vinogradova", Food Chemistry, 2006

Publication

<1%

51

Submitted to University of Queensland

Student Paper

<1%

52

Submitted to iGroup

Student Paper

<1%

53

Submitted to Macquarie University

Student Paper

<1%

54

Wang, B.G.. "In vitro antioxidative activities of extract and semi-purified fractions of the marine red alga, *Rhodomela confervoides* (*Rhodomelaceae*)", Food Chemistry, 20090415

Publication

<1%

55

Gaurav Rajauria. "Effect of hydrothermal processing on colour, antioxidant and free radical scavenging capacities of edible Irish brown seaweeds : Effect of heat on seaweeds antioxidant power", International Journal of Food Science & Technology, 12/2010

<1%

56 Bin Zhang, Chuan-dong Fang, Gui-juan Hao, Yang-yang Zhang. "Effect of kappa-carrageenan oligosaccharides on myofibrillar protein oxidation in peeled shrimp (*Litopenaeus vannamei*) during long-term frozen storage", Food Chemistry, 2018 <1%
Publication

57 Submitted to Dongeui University Graduate School <1%
Student Paper

58 Balboa, Elena M., Enma Conde, Andres Moure, Elena Falqué, and Herminia Domínguez. "In vitro antioxidant properties of crude extracts and compounds from brown algae", Food Chemistry, 2013. <1%
Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On