

ChemTech

International Journal of ChemTech Research CODEN (USA) : IJCRGG ISSN : 0974-4290, ISSN(Online):2455-9555

Vol. 10, No. 15, 2017

This website is best viewed in 1024 x 768

		INDEX Page-1.			
S.N.(Page Numbers)	Title of Paper and Names of Authors				
(01-08)	Revision of Optimized Hydrogel Design for the Covering of NPK Fertilizers as a Strategy for Sustainable Development Cifuentes Cetina Angie Roco; AvilaViatela Joahana Katherine, Rodrguez Miranda Juan Pablo				
	(09-14)	Analysis of the Environmental Quality Behavior in the Sumapaz River Basin using Data Mining (Colombia)			
Lpez Snchez Wils		on Ricardo; Rodrguez Miranda J	uan Pablo; Ruiz Ochoa Mau	ricio Andres	
Abstract		Full Text PDF	Full Text XML		
(15-20)	Environmental Quality Assessment in the Bogot River Basin (Colombia) using Particle Swagging				
	Rodrguez Miranda Juan Pablo; Garca Ubaque Cesar Augusto; Snchez Cspedes Juan Manuel				
	Abstract	Full Text PDF	Full Text XML		
	Environmental Quality Estimation using Diffuse Logic in the Machet River Hydrographic Basin (Cundinamarca, Colombia)				
(21-25)	Rodrguez Miranda Juan Pablo; Garca Ubaque Cesar Augusto; Snchez Cspedes Juan Manuel				
	Abstract	Full Text PDF	Eull Text XML		
(26-30)	Environmental Quality Estimation in the Minero River Basin (Cundinamarca, Colombia) using Artificial Neuronal Net - Levenberg Marquardt				
	Rodrguez Miranda Juan Pablo; Garca Ubaque Cesar Augusto; Snchez Cspedes Juan Manuel				
	Abstract	Full Text PDF	Full Text XML		
	Analysis of Variables in the River Suarez Basin using Data Mining (Colombia)				
(51-50)	Lpez Suchez Wilson Ricardo; Rodrguez Miranda Juan Pablo; Suchez Cspedes Juan Manuel				
	Abstract	Full Text PDF	Full Text XML		



2

ChemTech

International Journal of ChemTech Research

CODEN (USA) : IJCRGG ISSN : 0974-4290, ISSN(Online):2455-9555

Vol. 10, No. 15, 2017

This website is best viewed in 1024 x 768

INDEX Page-2.

S.N.(Page Numbers)	Title of Paper and Names of Authors				
(108-114)	Influence of sewage sludge and organic composts on different soils under incubation periods: I. Zinc and copper releases Hala Kandil				
	(115-121)	Influence of sewage sludge and organic composts on different soils under incubation periods: II. Heavy metals (Pb, Cd and Ni) releases.			
Hala Kandil					
Abstract		Full Text PDF	Full Text XML		
(122,125)	Level of Knowledge in Reproductive-Age-Women about Iron-Deficiency Anemia and Its Relationship between Age, Education and Occupation at Teladan Health Care Centre Medau				
(122-125)	Sry Suryani Widjaja, Sanny Tan				
	Abstract	Full Text PDF	Full Text XML		
	Synthesis of New coumarin substituted chromone with reference to their Anti-bacterial efficacy and fluorescence properties				
(126-134)	Pankaj S. Chaudhari, Shrikant S. Patil				
	Abstract	Full Text PDF	Full Text XML		
(135-145)	Review on dielectric study of polycrystalline rare earth substituted Tungsten Bronze electroceramics for high temperature applications				
	Shilpi Jindal, Ajay Vasisth, Sheela Devi				
	Abstract	Full Text PDF	Full Text XML		
(146-162)	Spectro-thermal characterization and molecular modeling of bioactive metal complexes of 4 - diethylaminosalicyldeoxime				
	Bibhesh K Singh, Jagdish Chandra, Hament K. Rajor, Pinkee				
	Abstract	Full Text PDF	Full Text XML		



ChemTech

International Journal of ChemTech Research

CODEN (USA) : IJCRGG ISSN : 0974-4290, ISSN(Online):2455-9555

Vol. 10, No. 15, 2017

This website is best viewed in 1024 x 768

INDEX Page-3.

S.N.(Page Numbers)	Title of Paper and Names of Authors				
(239-246)	Synthesis and vibrational study of 2-amino, 3-chloro 1,4-naphthoquinone by DFT				
	Shubhangi V.Kulkarni , Raghunath G. Sarawadekar and Avinash B. Pawar				
	Abstract	Full Text PDF	Full Text XML		
(247-254)	Synthesis and Characterization of Resorcinol based Schiff Base Ligand and Its Complexes of CO(III), NI(II), CU(II) and ZN(II) Metal Ions				
	S. Ramarasan, L. Rakesh Sharma, A. N. Paul Angelo				
	Abstract	Full Text PDF	Full Text XML		
(255-262)	Composition of Pigments and Antioxidant Activity in Edible Red Seaweed Halimenia durvilae Obtained from North Sulawesi				
j	Abstract	Full Text PDF	Full Text XML		
	The Potentiality of Cabomba aquatica DC not Aubletii Ethanol Extract as Larvicide against Aedes aegypti Larvae				
(263-271)	Erina Yatmasari, Soedarto, Sri Subekti Bendryman, Sukardiman				
	Abstract	Full Text PDF	Full Text XML		
	Clinical Outcome Difference of Internally Fixated Distal Radius Fracture Between Young Patients and Elderly In Haji Adam Malik General Hospital				
(272-276)	MH Siregar, N Nasution				
	Abstract	Full Text PDF	Full Text XML		
(277-281)	Physico-Chemical Properties and Fatty Acids Profile of Skipjack (Katsuwonus pelamis) Fish Oil				
(#77201)	Frans Lumoindong				
	Abstract	Full Text PDF	Full Text XML		





International Journal of ChemTech Research CODEN (USA): IJCRGG, ISSN: 0974-4290, ISSN(Online):2455-9555 Vol.10 No.15, pp 255-262, 2017

Composition of Pigments and Antioxidant Activity in Edible Red Seaweed *Halimenia durvilae* Obtained from North Sulawesi

Sanger G.*, Rarung L.K., Kaseger B.E., and Timbowo S.

Faculty of Fishery and Marine Science, Sam Ratulangi University. Manado, Indonesia

Abstract : This study was carried out to identify the pigment and antioxidant activity extracted from *Halimeniadurvilae*. Hexane, acetone and ethanol were used as extraction solvent. The pigments were measured consisting of chlorofhyll a, chlorofhyll b, total chlorofhyll, chlorofhyll c₁+c₂, fucoxathin, carotenoids, phycocyanin and phycoerythrin, while 1,1-diphenyl-2-picrylhydracyl (DPPH) radical scavenging and ferric reducing antioxidant power (FRAP) were used to studied their antioxidant activity. Total Phenolic content (TPC) were also investigated. The result showed hexane extract respectively containing highest level of all pigments. The lowest value of pigment was phycocyanin, recorded for hexane, acetone and etanol extract were 0.7875 ± 0.08 ; 0.1475 ± 0.08 and $0.1565 \pm 0.02 \mu g$ g⁻¹dried weight. Ethanol extract exhibited the lowest TPC ($7,605\pm0.383 \mu g$ GAE(Gallic acid equivalent)g⁻¹.Good value of radical scavenging of DPPH was acetone extract (scavenging activity of IC₅₀ 1.211±0.03 mg ml⁻¹), The highest Reducing power was acetone, 0.17 ± 0.01 uM Fe2+/mg extract, respectively. Thus *H.durvilae* could be used as natural pigment and antioxidant source which is potential to be applied in food product as functional food.

1. Introduction

Seaweeds or marine macro algae are the prospective renewable resource in marine environment. About 6.000 species of seaweeds have been identified and are classified into green algae (Chlorophyceae), brown algae (Phaeophyceae) and red (Rhodophyceae) (Devi *et al.*, 2011,Chandini*et al.*, 2008).The color in case of green seaweeds is due to the attendance of chlorophyll a and b; beta-carotene (a yellow pigment) and variuostypical xanthophylls (Yellowish or brownish pigments). Fucoxanthin is the prominent of the xanthophylls pigment which is responsible for the color of brown seaweeds. This compounds masks the other pigments such as Cholorophyill a and c and other xanthophylls. Phycoerythrin and phycocyanin accountable for the color of red seaweeds and mask the pigments such as Chlorophyll a and beta-carotene (Gupta and Abu-Ghannam, 2011.)Anthocyanins belonging to the flavonoid group are another group of pigments which are responsible for the red, purple, and blue colour Anthocyanins exhibited a high possible as colorants because of their low toxicity (Özkan and Bilek, 2014).

Macroalgae have been reported to have more then 2400 natural poducts of profitable significance in pharmaceutical, biomedical and nutraceutical industries. They have been utilized as ingredients in human and animal food preparations owning to their outstanding source of bioactive compounds which consist of sulfated polysaccharides, polyphenols, diterpenes, protein, essential fatty acids, dietary fiber vitamins and minerals((Chinnadurai et al., 2013, Özkan and Bilek, 2014, Chandihi *et al.*, 2007). Among functional ingredients identified from marine algae, natural pigments have obtained specific attention as they have been found to

showmanyadvantageous biological activities such as antioxidant, anticancer, anti-inflammatory, anti-obesity, anti-angiogenic and neuroprotective activities. (Pangestuti and Kim, 2011).

Humans are impacted by many free radicals especiallyreactive oxygen species (ROS). ROS constitutes superoxide (O2 -), hydroxyl (HO•), hydrogen peroxide (H2O2) and nitric oxide (NO). These molecules are unsteady and highly reactive, and can harm cells by chain reactions, such as lipid peroxidation or configuration of DNA adducts . Extreme amounts of ROS may be injurious because they can damage essential biomolecules: proteins, DNA and lipidsand other cells which consequences various diseases disorders such as cancer, diabetes, stroke, cataract, myocardial infarction, atherosclerotic and Parkinsons diseases (Wu *et al.*, 1988; Chew*et al.*, 2008). In order to diminish or avoid this damage of the human body by ROS antioxidants are believed to be protective, all cells perpectually contain antioxidants (Wu *et al.*, 1998, Halliwell *et al.*, 1995, Tapiero*et al.* 2002).

Natural antioxidants are chosen by consumer due to worry on the toxic and carcinogenic effect of synthetic antioxidant (Mantanjun *et al.*, 2007, Ahn*et al.*, 2003). Natural antioxidant considered safe for use as ingredients in medicine, dietary supplements, nutraceuticals and cosmetics with the intent of improving consumer health, reducing the belongings of damaging diseases and other broader aspects of immune system function (Shahidi, 2009, Pangestuti and Kim 2011, Yip *et al.*, 2014). Additional, there are facts obtainable in the literature to show the potential defensive properties of seaweeds against oxidative stress in target tissues and lipid oxidation in foods.Consequently, consumption of antioxidant and addition of antioxidant in food materials protect the body as well as against oxidative stress.Although seaweeds possess extensive applications in food and pharmaceutical industries, the pigments and antioxidant activities of many types of seaweeds in Indonesian area are still unexplored.Hence, the present study was proposed to explore thepigments and antioxidant properties of *H.durvilae* which grows plentifully in NorthSulawesi.

2. Materials and methods

Sample preparation and extraction

Sufficient amount *of H. durvilae was* collected from Arakan Village, Manado North Sulawesi. The sample was completely washed with seawater and fresh water to eliminate epiphytes, dirt particles and shells. It was then brought tothe laboratory followed with shading-drying for two days and oven-dryed at 50 °C for 3 days and ground using mixer without producing heat and transformed to powder.

The dried sample was extracted using hexane, acetone and ethanol (1 : 10 w/v) The samples were incubated for overnight at room temperature in a dark place. Extraction was repeated three times till the sample became colorless. The procedure was carried out in triplicates. The extracts were filtered and concentrated in vacum rotary evaporator at 40 °C. The extracts were stored in dark glass bottle for future analysis.

Chemicals and reagents

1,1-diphenyl-2-picrylhydrazyl (DPPH) was purchased from Sigma Aldrich, Follin-Ciocalteu's phenol, butylatedhydroxytoluene (BHT), sodium carbonate (Na₂CO₃), potassium dihydrogen phosphate (KH₂PO₄, iron (III) chloride-6-hydrate (FeCl₃), Trichloroacetic acid (TCA) and Potasium ferricyanide K₃Fe(Cn)6 purchased from Merk. All other solvent and chemicals were of analytical grade.

Pigment.

The pigments extracted using different solvents were quantified using UV-Visible spectrophotometer by reading the absorbance at their respective wavelengths and using the formulae given below:

-Chlorophyll Chl a (mg g-1) = $[12.7 (A663) - 2.69 (A645) V] / (1000 \times W)$ (Arnon, 1949).

-Chlorophyll Chl b (mg g^{-1}) = 22.9 x 0.645-4.68-A663)V]/(1000xW) (Arnon, 1949).

-Total Chlorophyll (mg g-1) = $[20.2 (A645) + 8.02 (A663) V] / (1000 \times W)$ (Jeffrey et al, 1961).

-Chlorophyll C1+C2 (mg g-1) = $[24.36 \times A630 - 3.73 \times A664]$ (Jensen and Jensen, 1959 & Duxbury and Yentsh, 1956).

-Carotenoids (μ g g-1) = [7.6 (A480) - 1.49 (A510) V] / (1000×W) (Seely et al, 1972).

-Fucoxanthin (mg g-1) = A470 -1.239 (A631+A581 $-0.3 \times A664$) $-0.0275 \times A664$ /141(Sudhakar*et al.*, 2013).

Total phenolic content (TPC)

The TPC of the extracts was measured using Follin Ciocalteu method as described by Ganesan *et al.*, (2008) with modification. 50% Follin Ciocalteu's phenol reagent (1 ml) was added to 0,1 ml extract and vortexed, Furthermore, added with 7% Na₂CO₃ (1 ml), and the reaction mixture was then incubated at room temperature for 30 min. The absorbance measured at 750 nm.TPC was expressed in terms of g gallic equivalents (μ GAE)/ g dried samples.

1,1-diphenyl-2-picrylhydrazyl (DPPH) assay

DPPH-scavenging potential of different concentrations of extracts was measured based on to test the scavenging ability of stable 1,1-diphenyl-2-picrylhydrazyl (DPPH) radicals by the seaweeds antioxidants. DPPH assay was measured using the method describe by Chew. *et al.*, (2008) with modification. Briefly, 2 ml of 93 μ M DPPH (solution in methanol) were added to 0.5 ml extract at various dilutions. The mixture was then vortexed vigorously and left for 20 minutes at room temperature in dark condition. The absorbance was measured at 517 nm and it activity was expressed as percentage of DPPH scavenging activity relative to the control, using that following equation:

% Inhibition = $[(A \text{ control} - A \text{ sample}) / A \text{ control}] \times 100\%.$

Ferric Reducing Antioxidant Power (FRAP).

Reducing power of seaweed extract was determined by the method prescribed by Chew*et al.*, (2008). The antioxidant activity of the standards was estimated by using the increasing in absorbance caused by the generated ferrous ion. Briefly, 1.0 ml of extracts at various dilutions were mixed with 1ml of 0,2 M phosphate buffer, pH 6,6 and 1 ml potassium ferricyanide (1%). Reaction mixture was incubate at 50°Cfor 20 minutes. After incubation, 1 ml of trichloroacetic acid (10%) was added and centrifuged at 3000 rpm for10 minutes. From the upper layer, 1 ml solution was mixed with 1 ml steril water and 0,5 ml FeCl₃ (0,1%). Absorbance of all solution was measured at 700 nm. The FRAP value was expressed in terms of uM Fe2+/mg extract.

Statistical analysis

All experiments were conducted in triplicate. The means of parameterspigment composition, total phenol content and antioxidant activity presented as mean \pm standart deviation.

3.Result and Discussion

Pigment

Regarding photosynthetic pigment a significant variation in content was recorded among the hexane, acetone and ethanol solvent studied (Tabel 1). The results show that *H. durvilae* contain highest amount of chlorophyll C1+C2 (2.26864 \pm 0.04mg g⁻¹ dried weight) whereas, the concentration of phycoerythrin and phycocyanin were found lowest, in hexane were2.33 \pm 0.253and 0.7875 \pm 0.08(µgg⁻¹). Naziret al., (2013), the concentration of carotenoids, phycoerythrin and phycocyaninin red seaweeds were found higher than green seaweeds, Pereira *et al*, 2012 reported that red seaweeds *Gracilaria dumingencis* possesses more phycocyanin than green strain. Red seaweeds observed are insignificant variation in chlorophyll a and phycocyanin (Plastino *et al.*, 2004).Pereira *et al*, 2012, seaweeds contain three main photosynthetic pigments i.e. chlorophylls, carotenoids and phycobilins. These pigments provide protection against high light intensity and also support in light absorption and energy transfer to the reaction centre.

Parameter	Hexane	acetone	ethanol
Chlorofil a (mg g ⁻¹)	0.181382 ± 0.01	0.01713 ±0.01	0.02181 ± 0.05
Chlorofil b (mg g^{-1})	0.038453 ± 0.001	0.00315 ± 0.001	0.01918 ± 0.006
Total Cholorofil (mg/g ⁻¹)	0.219787 ± 0.05	0.02038 ±0.001	0.04097 ± 0.002
Total chlorofil $c1+c2 (mg g^{-1})$	2.26864 ± 0.04	0.08763 ± 0.001	0.90726 ± 0.22
Carotenoid (µg g ⁻¹)	28.654 ±0.212	2.642 ±0.035	$5.96 \pm 0,3241$
Fucoxantin (mg g ⁻¹)	-0.37143 ± 0.03	0.023399 ± 0.005	-0.0198 ± 0.04
Pycoerytrin (µg g ⁻¹)	2.33 ± 0.253	0.48 ± 0.018	1.13 ±0.081
Phycocyanin (µg g ⁻¹)	0.7875 ± 0.08	0.1475±0.002	0. 1565 ±0.08

Table 1. Quantification of photosynthetic pigments of H.durvilae in different solvents

All the values are mean \pm SD of triplicates

Carotenoids have highest content in hexane extract, $28.654 \pm 0.212 \mu g^{-1}$. Carotenoids scavenge oxygen radicals and reduce oxidative stress. Thus, they denote antioxidant activity. The carotenoids are used in food, nutraceutical, and pharmaceutical preparations by their applications as colorants and their provitamin A activity. One of the most consumed carotenoid groups of pigments are responsible for the yellow, orange, and red colourof many foods, maintain to be intensely explored mainly because of their health-promoting properties. Carotenoids are unstable when they're exposed to light or oxygen because of their properties as highly conjugated and powerfully coloredisoprenoid plant compounds (Özkan and Bilek, 2014).

Variation in the pigment concentration is a reply to environmental variations that allows an organism to adjust under a specific habitat. Increased temperature, heavy metal accumulation and ill-inclined light due to extreme exploitation of natural resources and unrestrained anthropogenic activities, induces too much production of ROS which causes injury to biological membranes and unfavorably affect a number of plant physiological processes (Nasir *et al.*, 2015). Pigment shows several capability to maintain immune system, to help prevent cancer and is being utilized in cancer therapy, to aid to invigorate and energize the bodydetoxification of the liver, to normalize blood pressure and to struggle bad odors, bad breath as well as body odor by reason of the magnesium salts that it contains(Ferruzzi and Blakeslee, 2007).

Total Phenol Content

Various studies haveconcentrated on the biological activities or phenolic compounds, which are potential antioxidants and free radical-scavenger. Early researchreported that marine seaweeds extracts, particularly their polyphenols, have antioxidant activity. The most important active compounds in different seaweeds extracts have been revealed to the phlorotanins. The total polyphenol content (expressed as gallic acid equivalent) of *H.durvilae* extracts is shows in Fig.1. It was observed that hexane, aseton and etanol extracts had TPC 23.235±1,011; 22.151±1.967; 7.605±0.383µg GAEg⁻¹.Duan*et al.*, (2006) observed TPC (73,7 mg GAE g⁻¹) in ethyl acetate soluble fraction of red algae *P.urceolata*. the TPC in brown algae, *Papenfussiella kurono* was 0.18 mg cathechin equivalent/g in ethanolic extract (Kuda*et al.*, 2005).



Fig. 1. Total Phenolic Content of H. Durvilae extract. Data are expressed as mean ±SD (n=3).

Marine alge contain phloroglucinol phenol (phlorotannins) which are probably excelent antioxidant, since plant phenolics can perform as ROS scavengers metal chelators and enzyme modulator and inhibit lipid peroxidation (Rodrigo and Bosco 2006).Phenol, 2-[(1-phenylethyl)thio] characterize a diverse group of pigment, extensively distributed in nature. They serve as accessory pigments to harvest light photosynthesis. Moreover, these type of pigments can give rise to rich in polyphenol compounds. Norisoprenoids resulting from the oxidative cleave of carotenoid are signal in development, provide as antifungal and antibacterial agents and donate to their flavor and aroma. the norisoprenoid derivatives detected were present in the Rhodophyta, namely α -ionone, geranyl acetone, β -ionone,2,3-epoxy- β -ionone, dihydroactinidiolide and 2,3-Epoxy-b-ionone(Valentão*et al.*, 2010).

DPPH radical scavenging activity

DPPH is a compound that possesses a nitrogen free radical and is readily destroyed by a free radical scavenger. It has been used extensively as a free radical to assess reducing substances and is a valuable reagent for investigating the free radical scavenging activities of compounds. DPPH radical scavenging of *H.durvilae* are presented on Fig. 2.and expressed as presentation reduction of the initial DPPH• absorption by the tested compound.



Fig 2. DPPH radical scavenging activity of *H. durvilae* extract. Data are expressed as mean ±SD (n=3).

The best radical scavenging obtained in the aseton extract. The radical scavenging of hexane, acetone and ethanol extracts were IC₅₀ 15.940 \pm 1.63; 1.1676 \pm 1.99 and 8.017 \pm 2.31 mg ml⁻¹. The beneficial effect is could beside the high content of phenol, it may also be related to the high content of mineral, dietary fiber and pigment. The red seaweeds are a diverse eukaryotic lineage, characterized by accessory photosynthetic pigments phycoerythrin, phycocyanin and allophycocyanins arranged in phycobilisomes The extracts of *H.durvilae* showed better radical scavenging activity than ethanol extract of Kappaphycusalvarezii (Doty) $(IC_{50}3.03 \text{ mg ml}^{-1})$ (Kumar et al., 2008) and the extract of Palmaria palmita (dulse) (IC₅₀ 12.5 mg ml⁻¹) (Yuan, Carrington & Walsh, 2005). Seaweeds are low in fat but have vitamins and bioactive compounds such asterpenoids, sulfated polysaccharides and polyphenol compounds, the later being a potential natural antioxidant not found in land plants (Chew et al., 2008). Algae polysaccharides participate essential function as free radical-scavengers in-vitro and antioxidant for the avoidance of oxidative damage in living organism. Their activity depends on numerous structural parameter, such as the amount of sulfation, the molecular weight, sulfation position, type of sugar and glycosidic branching. Furthermore, several reports expose that the sulfate and phosphate group in the polysaccharides cause the differences in their biological activities (Juan et al. 2005a). Dietary natural antioxidant are reported to help in preventing aging and other diseases. There are various evidences that seaweeds contain compounds with a moderately high antioxidant and antiproliferative activity(Yuan, Carrington & Walsh, 2005).

Ferric reducing antioxidant power (FRAP).

The antioxidant activity in FRAP was measured based on the capability of the antioxidant compounds in the sample to reduce ferric (III) to ferrous (II) in a redox-linked colourimetric reaction that involve single electron (Chew *et al.*, 2008). FRAP assay was electron donor and it finished the oxidation chain reaction by reducing the oxidized intermediates into the stable form (Tachakittirungrod *et al.* 2007)

Fig. 3.shows that the highest reducing power of *H. durvilae* was acetone of 0.17 ± 0.01 uM Fe²⁺mg⁻¹ extract. The reducing ability of acetone extract considerably as same as TPC. There was strong correlation (R²=0.96) between the reducing power and the TPC of the seaweeds methanolic extracts expressesed as phologlicinol equivalents. (Matanjun *et al.*, 2008).



Fig.3. Ferric Reducing Antioxidant Power (FRAP) of *H.durvilae* extracts. Value are the mean ± SD (n=3)

The reducing power of red algae *Palmaria palmate* in 1-butanol extract was4.48 µg ascorbic acid equivalent (AAC) g⁻¹ (Yuan & walsh, 2006). Polyphenols are reducing agent, and together with other dietary reducing agent such as vitamin C, E and carotenoid, referred to as antioxidant, protect the body's tissues against oxidative stress and connected pathologies such as cancer, coronary heart disease and inflammation (Tapiero *et al.*, 2002). The ethanolic extract of *K. alvarezii* showed higher inhibitory effect than did the positive control, BHT. This might be due to the presence of ascorbic acid and Vitamin A (β -carotene) content in the extract of *K. alvarezii* (Fayaz*et al.*, 2005). The reducingpower property indicates that the antioxidant compounds are electron donor and can decrease the oxidized intermediates of the lipid peroxidation process, so that they can perform as primary and secondary antioxidants (Yen and Chen 1995).

Invitro antioxidant activity of κ -carrageenan oligosaccharides and their oversulfated, acetylated and phosphorilated derivatives was investigated by Juan *et al.* 2005. They are also reported that phosphorylated and sulfated glucans exhibited better antioxidant capacity than did glucans or other neutral polysaccharides, which indicated that polyelectrolytes, such as glucans sulfate or phosphate, might have enhanced scavenging activity. Moreover the sulfate content from *Porphyrayezoensis* was reported to contribute to the antioxidant activity. The cell walls of *K.avareziiis* known to be constitued of carrageenan, a sulfate polysaccharides, which may contribute to its antioxidant potential in addition to the presence of ascorbic acid, vitamin A and various phenolics (Kumar *et al.*, 2008).

Conclusions.

In the present investigation the various solvent extracts of *H.durvilae* exhibited content of chlorophyll carotenes, phycoerythrine and phycocyanin. The highest content of pigments are in hexane solvent. The antioxidant activity by DPPH assay and reducing power showed the acetoneextract is highest. Thus *H.durvilae* could be used as natural pigment and source of antioxidant which is potential to be applied in food product as functional food. Future study is required for identification of the active compound in acetone extract which is responsible for highest antioxidant activity.

Acknowledgments.

The authors are grateful to Prof. Dr. Ir. Remmy Mangindaan M.Sc. from Sam Ratulangi University Manado, the head of Laboratory Chemistry of Natural product and Dr. Ir. Dikdik Kurnia, M.Sc, Ms. The head of MIPA laboratory from Padjadjaran University Bandung for their constant support and provide research facilities..

References

- 1. Ahn, C.B., JoonY.J.and Kang D.S. 2004. Free Radical Scavenging Acrivityof Enzymatic Extract from a Brown Seaweed Scytophonlomentaria by Electron Spin Resonance Spectrometry. Journal of Food Research International 37: 253-258.
- 2. Arnon D., 1949. Copper Enzymes in Isolated Chloroplasts. Polyphenoloxidase in Beta Vulgaris.Plant Physiol. 24, 1-15.
- 3. Beer S., Eshel A. 1985. Determining Phycoerythrin and Phycocyanin Concentrations in Aqueous Crude Extracts Of Red Algae. Australian Journal of MarineAnd Freshwater Research. 36: 785 792.
- 4. Chandini S.K., GanesanP. and Bhaskar N. 2008. *In vitro* Antioxidant Activities of Three Selected Brown Seaweeds of India. *Food Chem.*,107: 707–713.
- 5. Chew Y.L., Lima Y.Y., Omara M., Khoob K.S. 2008. Antioxidant Activity of Three Edible Seaweeds from Two Areas in South East Asia.LWT.41: 1067-1072
- 6. Duxbury A.C., And Yentsch C.S. 1956. Plankton Pigment Nomographs. J. Mar. Res., 16: 92-101.
- 7. Devi G.K., Manivannan K.1., ThirumaranG., Rajathi F.A.A, Anantharaman P. 2011. In vitro Antioxidant Activities of Selected Seaweeds from Southeast Coast of India. Asian Pacific Journal of Tropical Medicine.205-211.
- 8. Duan, X., Zhang, W., Li, X. and Wang, B. 2006. Evaluation of Antioxidant Property of Extract and Fractions Obtained from A Red Algae, *PolysiphoniaUrceolata*. *Food Chem.*, 95: 37–43.
- 9. Fayaz M, Namitha K.K., Chidambara Murthy K.N., Swamy M., Sarada R., Khanam S, *et al.* 2005.Chemical Composition, Iron Bioavailability, and Antioxidant Activity of Kappaphycusalvarezzi (Doty).J.Agric Food Chem.53:792-797.
- 10. FerruzziM.G.and Blakeslee J. 2007. Digestion, Absorption, and Cancer Preventative Activity of Dietary Chlorophyll Derivatives. Nutrition Research 27:1-12.
- Ganesan P., Chandini S., Kumar N.Bhaskar. 2008. Antioxidant Properties of Methanol Extract and Its Solvent Fractions Obtained from Selected Indian Red Seaweeds. BiresourceTechnology .99: 2717-2723.
- 12. Gupta S. and Abu-Ghannam N. 2011. Bioactive Potential and Possible Health Effects of Edible Brown Seaweeds.Trends Food Sci.Technol.2011; 22: 315-326.
- 13. Halliwell, B. and C.E. Cross, 1994. Oxygen-Derived Species: Their Relation to Human Disease And Environmental Stress. *Environ. HealthPerspect.*, 102: 5–12.
- 14. Jeffry W. 1961. Paper-Chromatographic Separation of Chlorophyll And Carotenoids from Marine Algae. Biochem. J., 1961, 80, 336-342.
- 15. Jensen A. and Jensen S.L.1959. Quantitative Paper Chromatography of Carotenoids. Acta Chem. Scand. 13: 1863-68.
- 16. Kuda T.T., Sunekawa M., Goto H., Araki Y. 2005. Antioxidant Properties of Four Edible Algae Harvested in The Noto Peninsula, Japan. JFood Compos Anal.18:625-633.
- 17. Kumar, K.S., Ganesan, K. and Subba-Rao, P.V. 2008. Antioxidant Potential of Solvent Extracts of *KappaphycusAlvarezii* (Doty) Doty-An Edible Seaweed. Food Chem., 107: 289-295.
- Matanjun P., Suhaila Mohamed S., Noordin Mohamed stapha, Kharidah Muhammad, Cheng Hwee Ming. 2008. Antioxidant Activities and Phenolics Content of Eight Species of Seaweeds from NorthBorneo. J Appl.Phycol.20:367-373.
- Nasir K.M., Mobin M., Abbas Z.K., 2015. Variation in Photosynthetic Pigments, Antioxidant Enzymes and Osmolyte Accumulation in Seaweeds of Red Sea. Int J Plant Biol. Res. 3(1): 1028.
- 20. Özkan G. and BilekS.E. 2014. Microencapsulation of Natural Food Colourants. Int. J. of Nutrition and Food Sciences.3(3): 145-156.
- 21. Pangestuti, R. and Kim S.K. 2011. Biological Activities and Health Benefit Effects of Natural Pigments Derived from Marine Algae. *Journal of Functional Foods* 3: 255-266.

- 22. Pereira DC., Trigueiro TG., Colepicolo P., Marinho-soriano E.2012. Seasonal Changes in The Pigment Composition of Natural Population of *Gracilaria dumingensis (Graciliares*, Rhodophyta); Brazillians J. of Pharmacognosy. 22:874-880.
- 23. Plastino E.M., Ursi S., Fuijii M.T.2004. Color Inheritance, Pigment Caractrization and Growth of A Rare Light Green Strain of *Gracilaria birdiae* ((*Graciliares*, Rhodophyta) phycological Research. 52:45-52.
- 24. Rodrigo R. Bosco C., 2006. Oxidative Stress and Protective Effect of Polyphenol: Comparative Studies in Humans and Rodent Kidney. A review. Comparatives Biochem. Physiol. Part C (142:317-327).
- 25. Seely G.R., Duncan M.J., & Vidaver W.E. 1972. Preparative And Analytical Extraction of BrownAlgae with Dimethyl Sulfoxide. Marine Biology, 12: 184-188.
- 26. Shahidi F., 2009. Nutraceuticals and Functional Foods: Whole Versus Processed Foods. *Trends Food Sci. Tech.*, 20: 376–387.
- 27. Sudhakar M.P., Ananthalaksmi J.S. and Nasir B.B. 2013. Extraction and Purification and Study on Antioxidant Properties of Fucoxantin from Brown Seaweeds. J. of Chemical and PharmaceuticalResearch 5(7): 169-75.
- 28. Tapiero H., Tew K.D., Nguyen Ba G., Mathe'G., (2002). Polyphenols: Do They Play A Role in The Prevention of Humans Pathologies? Review.Biomed.Pharmaco. 56:200-207.
- 29. Tachakittirungrod S, Okonogi S, Chowwanapoonpohn S., 2007. Study on Antioxidant Activity of Certain Plant in Thailand: Mechanism of Antioxidant Action of Guava Leaf Extract. Food Chem., 103:381-388.
- 30. Valentão P., Trindade P., Gomes D., Guedesde Pinho P., Mouga T. and Andrade P.B. 2010. *Codium tomentosum* and *Plocamium cartilagineum*: Chemistry and Antioxidant Potential J. FoodChemistry119.1359–1368.
- 31. Wu, Y., C. Hong, S. Lin, P. Wu and M. Shiao, 1998. Increase of Vitamin E Content in LDL and Reduction of Atherosclerosis in Cholesterol-FedRabbits by a Water-Soluble Antioxidant-Rich Fraction of *Salvia miltiorrhiza*. *Arterioscler*. *Thomb.Vasc. Biol.*, 18: 481–486.
- 32. Yen G.C. Chen H.J., 1995. Antioxidant Activity of Various Tea Extract in Relation to Their Antimutagenicity.J. Agryc. Food Chem.43 27-32.
- 33. Yip W.H., Lim Seng Joe L.S. Mustapha W.A.W., Maskat M.Y. And Said M., 2014Characterisation and Stability of Pigments Extracted from *Sargassum binderi* Obtained from Semporna, Sabah. SainsMalaysiana 43(9): 1345–1354
- 34. Yokoya NS, Necchi O, Martins AP, Gonzalez SF, Plastino EM.2007. Growth Responses and Photosynthetic Characteristics of Wild and Pycoerythrin-Deficient Strains of *Hypnea musciformis*(Rhodophyta). Journal of Applied Phycology. 19: 197-205.
- 35. Yuan A., Zhang W., Li X., Lu X., Li N., GaoK., *et al.*, 2005. Preparation and In-Vitro Antioxidant Activity of k-Carrageenan Olygosaccharides and Their Oversulfated, Acetylated and Phosphorylated Derivatives. Carbohydrate Research, 300: 685-692.
- 36. Yuan Y.V., Carrington M.F., Walsh N.A., 2005. Extracts from Dulse (Palmaria palmata) are Effective Antioxidants and Inhibitors of Cell Proliferation in vitro. Food ChemToxicol.43:1073-1081.
- 37. Yuan, Y .and Walsh, N.A. 2006. Antioxidant and Antiproliferative Activity of Extraxts from A Variety of Edible Seaweeds. Food And Chemical Toxicology, 44:1144-1150.
