

Phytochemical Characterization of Cucumber (*Cucumis sativus* L.) Seeds as Candidate of Water Additive for Organic Broiler Chickens

by Jet Mandey 7

Submission date: 07-Oct-2021 12:03PM (UTC+0700)

Submission ID: 1667525459

File name: Phytochemical_Characterization_of_Cucumber.pdf (246.51K)

Word count: 3110

Character count: 17672

Phytochemical Characterization of Cucumber (*Cucumis sativus* L.) Seeds as Candidate of Water Additive for Organic Broiler Chickens

Jet S. Mandey, Fenny R. Wolayan, Cherly J. Pontoh, and Bonie F. J. Sondakh
Animal Husbandry Faculty, Sam Ratulangi University, Manado, Indonesia

Email: jetsm_fapet@yahoo.co.id, feny_wolayan@unsrat.ac.id, cherlypontoh61@gmail.com, bfjsondakh@yahoo.com

Abstract—The research was conducted to evaluate the phytochemical characterization of cucumber (*Cucumis sativus* L.) as candidate of natural water additive for organic broiler chickens. Research was done by laboratory analysis to phytochemicals, proximate on chemical compounds, β -carotene and antioxidant potency of cucumber seeds. Data were analysed by descriptive method. Results showed that phytochemical screening by color visualization showed the presence of flavonoid, tannin, saponin and steroid. And, the quantitative analysis by spectrophotometry found the flavonoid total 0.36% (w/w), phenol total 0.40% (w/w), and analysis by titrimetry found tannin 2.82%. Proximate analysis on cucumber seeds were dry matter 90.53%, crude protein 26.68%, crude fat 14.14%, crude fiber 32.27%, Ca 3.024%, P 0.807% and gross energy 4817.72 Kkal. β -carotene compound analysed by TLC Scanner was 2.82% and antioxidant IC 50% analyzed by DPPH/Spectrophotometry was 6555.55 ppm. It can be concluded that cucumber seeds can be used as an alternative water additive because of nutrients and antioxidant bioactive potency.

Index Terms—bioactives, cucumber, feed additive, nutrients, water additive

I. INTRODUCTION

Plants are the principal source of nutrients for animals, and sometimes as an unwelcome source of anti-nutritional factors that interfere with an animal's ability to maximize utilization of ingested nutrients. The present review considers the background of interest in the use of plants and their extracts as alternative performance enhancers, and the non-nutrient bioactive compounds of plants [1].

Phytochemicals are primary and secondary compounds. Chlorophyll, proteins and common sugars are included in primary constituents and secondary compounds have terpenoid, alkaloids and phenolic compounds [2]. According to Mallikharjuna, *et al.* [3] phytochemicals are bioactive non-nutrient plant compounds that have protective or disease preventive property. They confer plants with odour (terpenoids), pigmentation (tannins and quinines), and flavour (capsacin). These bioactive components are said to be responsible for the antimicrobial effects of plant extracts

in vitro. They are grouped as flavonoids, alkaloids, glycosides, saponins, tannins, terpenoids, carbohydrates, and sterols. Tang, *et al.*, [4] have isolated the antimicrobial sphingolipids from cucumber stem.

Cucumber (*Cucumis sativus* L.), belongs to family Cucurbitaceae, is most widely cultivated vegetable crop all over the world [5]. Cucumber is the fourth most important vegetable crop after tomato, cabbage, and onion [6], though its calorie and nutritional value is very low, it is a primary source of vitamins and minerals in the human diet [7]. Due to high content of potassium (50-80 mg/100g), cucumber can highly be useful for both high and low blood pressures [8].

Cucumber is a widely cultivated plant of gourd family which is eaten in the unripe, green form. Its fruit extract has shown free radical scavenging and analgesic activities in mice [9], carminative and antacid property [10]. Studies of Gill, *et al.* [11] have shown the antioxidant and antiulcer effect of *C. sativus* in rats. The vegetables contain several phytochemicals possessing antioxidant activity, and the major groups of phytochemicals include vitamin A, C, E and K, carotenoid, terpenoid, flavonoids, polyphenols, saponins, enzymes and minerals.

Recent researchs have identified a vast majority of antioxidants from vegetables and fruits, like vitamin A, C, E, β -carotene, glutathione precursors like selenium, vitamin B2 (Riboflavin), B3 (Niacin), B6 (Pyridoxin), B9 (Folic acid), B12 (Cyanocobalamine). Bioflavonoids are very rich in all vegetables with vivid colors stand as prophylactic [12]. Tang, *et al.* [4] have isolated the antimicrobial sphingolipids from cucumber stem. Abiodun and Adeleke [13] reported that the seeds of the plant served as good source of protein, fat, minerals and calcium. Seeds are cooling, tonic, diuretic and anthelmintic [14]. There is a lack information about the research of cucumber seeds, so, this research was conducted to evaluate the phytochemical characterization of cucumber seeds as candidate of natural water additive for organic broiler chickens.

II. MATERIALS AND METHODS

A. Sample Collection

The sample of *Cucumis sativus* seeds were washed, dried and stored in cold dark conditions. Powdered

Manuscript received June 30, 2018; revised November 22, 2018.

seeds were defatted with hexane. Then extraction was carried out with solvents of increasing polarity such as methanol by maceration process for 16 h at room temperature.

Determination ³⁰ *Proximate Chemical Composition:* Determination of moisture, total ash, crude protein, crude fiber, crude fat, were determined by standard method [15].

B. Phytochemical Screening

The preliminary phytochemical screening of the extracts was carried out to know the different constituents present as per the standard procedures. The extracts were tested for alkaloids, sterols, triterpenes, saponins, flavonoids, quinon and tannins [16], [17].

C. In-vitro Antioxidant Activity

DPPH (2,2-Diphenyl 1-picryl hydrazyl) assay method: The DPPH free radical is reduced to a corresponding hydrazine when it reacts with hydrogen donors. The DPPH radical is purple in colour and upon reaction with hydrogen donor changes to yellow colour. It is a decoloration assay, which is evaluated by the addition of the antioxidant to a DPPH solution in methanol and the decrease in absorbance was measured at 490nm.

Reagents: 2,2-Diphenyl 1-picryl hydrazyl solution (DPPH, 100µM): 22mg of DPPH was accurately weighed and dissolved in 100 ml of methanol. From this stock solution, 18 ml was taken and diluted to 100 ml using methanol to obtain 100 µM DPPH solution [18].

III. ³⁶ RESULTS AND DISCUSSION

A. Phytochemical Screening and Flavonoid Estimation

³⁵ in this study, the qualitative phytochemical analysis revealed the presence of bioactive compounds of cucumber seed (Table I). Results showed that phytochemical screening by color visualization showed the presence of flavonoid, tannin, saponin and steroid. The quantitative analysis spectrophotometry found flavonoid total 0.36% (w/w), phenol total 0.40% (w/w), and, the by titrimetry found tannin 2.82%. Proximate analysis on cucumber seeds were dry matter 90.53%, crude protein 26.68%, crude fat 14.14%, crude fiber 32.27%, Ca 3.024%, P 0.807% and gross energy 4817.72 Kcal. The value of β-carotene compound analyzed by TLC Scanner was 2.82% and antioxidant IC 50% analyzed by DPPH/Spectrophotometry was 6555.55 ppm.

The cucumber seed contained flavonoid that can be considered to a che¹⁵ source of flavonoids. According to Saxena, *et al.* [19], flavonoids have been reported to exert multiple biological property including antimicrobial, antioxidant, cytotoxicity, anti-infla²⁰tory, as well as antitumor activity. Flavonoids, mainly present as colouring pigments in plants also function a¹⁹ potent antioxidants at various levels [20], [21]. Phenolic compounds comprise a large group of organic substances and flavonoids are an important subgroup. This group constitutes the majority of dietary antioxidants [22].

The cucumber seeds in this research also have tannin, saponin and steroid. This research was in line with Njoku and Akumefula [23], who reported that tannins have been found²⁴ the extracts of *Cucumis sativa* (Cucumber). Tannins have astringent properties, hasten the healing of wounds and inflamed mucous membrane. Tannins are potential metal ion chelator, proton precipitating agents and biological antioxidant [24]. Phytosterols have been found in the extracts of cucumber, also was reported by Castro, *et al.* [25], who stated that phytosterols have a significant hypocholesterolemic effect. Saponins have the property of precipitating and coagulating red blood [26]. They exhibit foaming properties and cell membrane-permeabilizing properties. Their soapy character is due to their surfactant properties [27].

TABLE I. PHYTOCHEMICAL SREENING OF CUCUMBER SEEDS

Phytochemical	Content	Unit	Technique Analyzed
Flavonoid	positive	-	Color visualization
Alkaloid	Wagner	negative	-
	Mayer	negative	-
	Dragendorf	negative	-
Tannin	positive	-	-
Saponin	positive	-	-
Quinon	negative	-	-
³¹ bid	positive	-	-
Triterpenoid	negative	-	-
Total Flavonoid	0.36	% (w/w)	Spectrophotometry
Total Phenol	0.40	% (w/w)	Spectrophotometry
Tannin	0.01	%	Titrimetry
β-carotene	2.82	%	TLC Scanner
Antioxidant IC 50%	6555.55	ppm	DPPH/Spectrophotometry

Cucumber fruits have been evaluated for a wide spectrum of activity including diuretic, antihyperglycemic, antioxidant, amyolytic, anticancer and analgesic activities *in vitro* and *in vivo* models [28]-[30]. There were only several reports to the potency of cucumber seeds. Gill, *et al.* [11] reported that methanolic extract of cucumber seed have triterpenoid, saponin, phytosterol, tannin, that possessed significant antiulcer potential which could be due to its antioxidant activity. Vetriselvan, *et al.* [31] reported that the seeds of the plant *Cucumis sativus* has anti-inflammatory effect that support to the traditio¹² use of this plant for the treatment of related diseases. The seed extracts were found effective on controlling the loss of body weight in diabetic rats and against tapeworms [32].

Safe alternative to endogenous antioxidants is natural antioxidants. Fruits, vegetables and their products are dietary sources of antioxidants for maintaining the cellular functions in best way. Due to excellent potential of these compounds against chronic diseases, different reports have shown the vital role of antioxidants of fruits and vegetables to maintain body function in normal mode.

Antioxidant compounds like phenolic acids, polyphenols and flavonoids scavenge free radicals such as peroxide, hydroperoxide or lipid peroxy and thus inhibit the oxidative mechanisms that lead to degenerative diseases [22]. Gill and Bali [14] stated that

triterpenoid glycosides present in cucumber seeds maybe responsible for the antioxidant and antiulcer activity.

Among the various natural pigments, carotenoids comprise a large family of more than 700 structures [33]. Most carotenoids can be derived from a 40-carbon basal structure, which includes a polyene chain contains 3 to 15 conjugated double bonds. The pattern of conjugated double bonds in the polyene backbone of carotenoids determines their light absorbing properties and influences the antioxidant activity of carotenoids [34], [35].

B. Nutritional Evaluation

Chemical composition of cucumber seed meal is shown in Table II. Result showed that investigations on nutritional evaluation of cucumber revealed that their seeds are rich in crude protein, crude fat, gross energy and mineral Ca, followed by crude fiber. Abulude, *et al.* [36] reported chemical composition of cucumber, such as, protein in mesocarp 1.68%, in epicarp 3.84% and in endocarp 0.22%. Fat in epicarp 0.56% and in endocarp 0.02%. In this research, these plants resulted as a good source of nutrients (proteins, fats, carbohydrates, fiber and minerals) and can be used as alternative feed additive/water additive in either of these nutrients for organic broiler chickens.

TABLE II. CHEMICAL COMPOSITION OF CUCUMBER SEED MEAL

Ingredients	Content
Dry matter (%)	90.53
Crude Protein (%)	26.68
Crude Fiber (%)	32.27
Crude Fat (%)	14.14
Ca (%)	3.024
P (%)	0.807
Gross Energy (Kcal)	4817.72

IV. CONCLUSION

The present study reveals that these cucumber seeds served as good source of protein, fat, minerals, calcium and secondary compound which can be used for additive. It can be concluded that cucumber seeds can be used as an alternative water additive because of nutrient and antioxidant bioactive potency.

ACKNOWLEDGEMENT

The authors wish to thank to the "Program Bantuan Seminar Luar Negeri, Ditjen Penguatan Riset dan Pengembangan, Kemenristekdikti" for their financial supported and encouragement.

REFERENCES

[1] H. Greathead, "Plants and plant extracts for improving animal productivity," *Proc. of the Nutrition Society*, vol. 62, pp. 279-290, 2003.
 [2] D. Krishnaiah, R. Sarbatly, and A. Bono, "Phytochemical antioxidants for health and medicine: A move towards nature," *Biotechnol. Mol. Biol. Rev.*, vol. 1, pp. 97-104, 2007.
 [3] P. B. Mallikharjuna, L. N. Rajanna, Y. N. Seetharam, and G. K. Sharanabasappa, "Phytochemical studies of *Strychnos potatorum* L.f. - A medicinal plant," *E-J. of Chemist.*, vol. 4, no. 4, pp. 510-518, 2007.

[4] J. Tang, *et al.*, "Antimicrobial activity of sphingolipids isolated from the stems of Cucumber (*Cucumis sativus* L.)," *Molecules*, vol. 15, pp. 9288-9297, 2010.
 [5] H. S. Paris, M. C. Daunay, and J. Janick, "Occidental diffusion of cucumber (*Cucumis sativus*) 500-1300 CE: Two routes to Europe," *Annals of Botany*, pp. 1-10, 2011.
 [6] T. Tatlioglu, "Cucumber (*Cucumis sativus* L.)," in *Genetic Improvement of Vegetable Crops*, G. Kallo and B. Bergn, Eds., Oxford: Pergamon Press, 1993, pp. 197-227.
 [7] S. Y. Mah, "An effective fungicide for the control of downy mildew on cucumber," *MAPPs Newsletter*, vol. 12, no. 4, p. 40, 1989.
 [8] W. Kashif, Q. M. Kamran, and M. S. Jillani, "Effect of different nitrogen levels on growth and yield of cucumber (*Cucumis sativus* L.)," *J. Agric. Res.*, vol. 46, no. 3, pp. 259-266, 2008.
 [9] D. Kumar, S. Kumar, J. Singh, Narender, Rashmi, B. D. Vashistha, and N. Singh, "Free radical scavenging and analgesic activities of *Cucumis sativus* L. fruit extract," *J. of Young Pharmacist.*, vol. 2, no. 4, pp. 365-368, 2010.
 [10] S. Shama, J. Dwivedi, and S. Paliwal, "Evaluation of antacid and carminative properties of *Cucumis sativus* under simulated conditions," *Scholars Research Library, Der Pharmacia Lettre*, vol. 4, no. 1, pp. 234-239, 2012.
 [11] N. S. Gill, M. Garg, R. Bansal, S. Sood, A. Muthuraman, M. Bali, and P. D. Sharma, "Evaluation of antioxidant and antiulcer potential of *Cucumis sativum* L. seed extract in rats," *Asian J. of Nutritional Nutr.*, vol. 1, no. 3, pp. 131-133, 2009.
 [12] J. S. Godber, "Nutrient bioavailability in humans and experimental animals," *J. Food Quality*, vol. 13, pp. 21-36, 1990.
 [13] O. A. Abiodun and R. O. Adeleke, "Comparative studies on nutritional composition of four melon seeds varieties," *Pakistan J. Nutrition*, vol. 9, no. 9, pp. 905-908, 2010.
 [14] N. S. Gill and M. Bali, "Evaluation of antioxidant, antiulcer activity of 9-beta-methyl-19-norlanosta-5-ene-type glycosides from *Cucumis sativus* seeds," *Res. J. of Medicinal Plant*, vol. 6, no. 26 p. 309-317, 2012.
 [15] AOAC, "Official methods of analysis," Association of Official Analytical Chemist, Washington, D.C., 2000.
 [16] J. B. Harborne, *Phytochemical Methods*, London: Chapman and Hall Ltd., 1973, pp. 14-188.
 [17] J. B. Harborne, *Phytochemical Method: A Guide to Modern Techniques of Plant Analysis*, 3rd ed., New Delhi; Springer Pvt. Ltd, 2005, pp. 5-16, 22.
 [18] N. Muruganatham, S. Solomon, and M. M. Senthamselvi, "Anti-oxidant and anti-inflammatory activity of *Cucumis sativus* (cucumber) flowers," *Int. J. of Pharm. Sci. & Res.*, vol. 7, no. 4, pp. 1740-1745, 2016.
 [19] M. Saxena, J. Saxena, R. Nema, D. Singh, and A. Gupta, "Phytochemistry of medicinal plants," *J. of Pharmacog. and Phytochem.*, vol. 1, no. 6, pp. 168-180, 2013.
 [20] H. Sies, *Antioxidants in Disease, Mechanisms and Therapy*, New York, USA: Academic Press, 1996.
 [21] E. Cadenas and L. Packer, *Hand Book of Antioxidants*, New York: Plenum Publishers, 1996.
 [22] G. Nahak, M. Suar, and R. K. Sahu, "Antioxidant potential and nutritional values of vegetables: A review," *Res. J. of Medicinal Plant.*, vol. 8, no. 2, pp. 50-81, 2014.
 [23] P. C. Njoku and M. I. Akumefula, "Phytochemical and nutrient evaluation of *Spondias mombin* leaves," *Pakistan J. of Nutr.*, vol. 6, no. 6, pp. 613-615, 2007.
 [24] S. I. Okonkwo, "Isolation and characterization of tannin metabolites in *Spondias mombin* (Linn) (Anacardiaceae)," *Natural Sci. Appl. Sci. J.*, vol. 10, no. 1, pp. 21-29, 2009.
 [25] I. A. Castro, L. P. Barros, and P. Sinnecker, "Functional foods for coronary heart disease risk reduction: A meta-analysis using a multivariate approach," *American J. of Clinical Nutr.*, vol. 82, pp. 32-40, 2005.
 [26] D. E. Okwu and C. Josiah, "Evaluation of the chemical composition of two Nigerian medicinal plants," *African J. of Biotech.*, vol. 5, no. 4, pp. 357-361, 2006.
 [27] Y. F. Chu, J. Sun, X. Wu, and R. H. Liu, "Antioxidant and antiproliferative activities of common vegetables," *J. Agric. Food Sci.*, vol. 50, no. 23, pp. 6910-6916, 2002.
 [28] I. M. Villasenor, M. K. Simon, and A. M. Villanueva, "Comparative potencies of nutraceuticals in chemically induced

skin tumor prevention," *Nutr. Cancer*, vol. 44, no. 1, pp. 66-70, 2002.

- [29] M. K. Patil, A. D. Kandhare, and S. D. Bhise, "Pharmacological evaluation of ameliorative effect of aqueous extract of *Cucumis sativus* L. fruit formulation on wound healing in Wistar rats," *Chron. Young Sci.*, vol. 2, pp.207-213, 2011.
- [30] S. Vetrivelvan, U. Subasini, C. Velmurugan, T. Muthuramu, J. Shankar, and Revathy, "Anti-inflammatory activity of Cucumis sativus seed in carrageenan and xylene induced edema model using albino wistar rats," *Int. J. of Biopharmaceutics*, vol. 4, no. 1, pp. 34-37, 2013.
- [31] E. E. Elisha, H. A. A. Twaij, N. M. Ali, J. H. Tarish, M. M. Al-Omari, and S. Karim, "The anti-helminthic activity of some Iraqi plants of the *Cucurbitaceae*," *Pharma. Bio.*, vol. 25, pp. 153-157, 1987.
- [32] G. Britton, S. Liaanen-Jensen, and H. Pfander, *Carotenoids Handbook*, Boston, USA: Birkhanser Verlag, 2004.
- [33] T. W. Godwin, "Carotenoids," in *Encyclopedia of Plant Physiology*, E. A. Bell and B. V. Charlwood, Eds., vol. 8, Spenger-Verlag-Berlin, 1980, pp. 257-281.
- [34] G. Britton, S. Liaanen-Jensen, and H. Pfander, *Carotenoids, Biosynthesis and Metabolism*, vol. 3, Birkhanser, Verlag, Brusel, Switzerland.
- [35] F. O. Abulude, Y. S. Akinyagunla, T. Abe, B. E. Awanlemhen, and O. Afoladi, "Proximate composition, selected mineral, physical characteristic and *in vitro* multienzyme digestibility of cucumber (*Cucumis sativus*) fruit from Nigeria," *American J. of Food Tech.*, vol. 2, no. 3, pp. 196-201, 2007.



Jet Saartje Mandey was born in Manado, North Sulawesi, Indonesia in 1953. She posses the Master (1987) degree in Public Nutrition and Family Resources from Bogor Agricultural Institute and Doctor (2013) degree in Animal Nutrition from Brawijaya University. Professor Mandey's specialization area in Animal Nutrition. She joined the services of Animal Husbandry Faculty, Sam Ratulangi University as a lecturer since 1980.

Her current research interests include: poultry nutrition with emphasis agro-industrial by product, and use of various feed and water additives (herbal and local spices).



Fenny Rinay Wolayan was born in Treman, North Sulawesi, Indonesia in 1962. She possessed the Master degree (1998) and Doctor degree (2012) degree in Feed Technology from Padjadjaran University. Dr. Fenny's specialization area in Feed Technology. She joined the services of Animal Husbandry Faculty, Sam Ratulangi University as a lecturer since 1990. Her current research interests include: feed technology with emphasis agro-industrial by product.



Cherly Joula Pontoh was born in Tomohon, North Sulawesi of Indonesia in 1961. She possessed the Master Degree (2006) in Agronomy from Sam Ratulangi University. She joined the services of Animal Husbandry Faculty, Sam Ratulangi University as a lecturer since 1987. Her current research interests include animal nutrition.



Bonie F. J. Sondakh was born in Manado, North Sulawesi of Indonesia in 1954. He possessed the Master Degree (1985) in Rural Sociology from Bogor Agricultural Institute. He joined the services of Animal Husbandry Faculty, Sam Ratulangi University as a lecturer since 1980. His current research interests include: woman's roles in rural livestock.

Phytochemical Characterization of Cucumber (*Cucumis sativus* L.) Seeds as Candidate of Water Additive for Organic Broiler Chickens

ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

16%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1	Stahl, W.. "Antioxidant activity of carotenoids", Molecular Aspects of Medicine, 200312 Publication	1%
2	krishikosh.egranth.ac.in Internet Source	1%
3	waset.org Internet Source	1%
4	scialert.net Internet Source	1%
5	www.drnatura.com Internet Source	1%
6	"7th International Phytocosmetics and Phytotherapy Congress (India - IPPC2018)", International Journal of Phytocosmetics and Natural Ingredients, 2018 Publication	1%
7	Yung-Hsing Peng, Chin-Shun Hsu, Po-Chuang Huang. "An investigation of spacial	1%

approaches for crop price forecasting in different Taiwan markets", 2015 Conference on Technologies and Applications of Artificial Intelligence (TAAI), 2015

Publication

8

Henry Greathead. "Plants and plant extracts for improving animal productivity", Proceedings of The Nutrition Society, 05/2003

Publication

1 %

9

innovareacademics.in

Internet Source

1 %

10

Azar Hosseini, Hossein Hosseinzadeh. "Antidotal or protective effects of Curcuma longa (turmeric) and its active ingredient, curcumin, against natural and chemical toxicities: A review", Biomedicine & Pharmacotherapy, 2018

Publication

1 %

11

doi.org

Internet Source

1 %

12

www.ipindia.nic.in

Internet Source

1 %

13

C. Baskaran, V. Ratha bai, S. Velu, Kubendiran Kumaran. "The efficacy of Carica papaya leaf extract on some bacterial and a fungal strain by well diffusion method", Asian Pacific Journal of Tropical Disease, 2012

Publication

1 %

14	core.ac.uk Internet Source	1 %
15	documents.mx Internet Source	1 %
16	jmp.ir Internet Source	1 %
17	pharmacophorejournal.com Internet Source	<1 %
18	www.scitechnol.com Internet Source	<1 %
19	www.bib.irb.hr Internet Source	<1 %
20	www.biotech-asia.org Internet Source	<1 %
21	www.pakbs.org Internet Source	<1 %
22	citeseerx.ist.psu.edu Internet Source	<1 %
23	files.eric.ed.gov Internet Source	<1 %
24	jbasicphyres-unizik.org Internet Source	<1 %
25	valleyinternational.net Internet Source	<1 %

26	www.sciencepub.net Internet Source	<1 %
27	Paloma Benito, Dennis Miller. "Iron absorption and bioavailability: An updated review", Nutrition Research, 1998 Publication	<1 %
28	Sang-Mo Kang, Muhammad Hamayun, Gil-Jae Joo, Abdul Latif Khan, Yoon-Ha Kim, Sang-Kuk Kim, Hyung-Jin Jeong, In-Jung Lee. "Effect of Burkholderia sp. KCTC 11096BP on some physiochemical attributes of cucumber", European Journal of Soil Biology, 2010 Publication	<1 %
29	tohumcu.org Internet Source	<1 %
30	journalrepository.org Internet Source	<1 %
31	TUTIK WRESDIYATI, SITI SA'DIAH, ADI WINARTO, VENNY FEBRIYANI. "Alpha-Glucosidase Inhibition and Hypoglycemic Activities of Sweitenia mahagoni Seed Extract", HAYATI Journal of Biosciences, 2015 Publication	<1 %
32	publications.drdo.gov.in Internet Source	<1 %

33 Pasquale Crupi, Rosa Anna Milella, Donato Antonacci. "Simultaneous HPLC-DAD-MS (ESI+) determination of structural and geometrical isomers of carotenoids in mature grapes", Journal of Mass Spectrometry, 2010
Publication <1 %

34 globaljournals.org
Internet Source <1 %

35 Chan, Chim Kei, Bey Hing Goh, Muhamad Noor Alfarizal Kamarudin, and Habsah Abdul Kadir. "Aqueous Fraction of Nephelium ramboutan-ake Rind Induces Mitochondrial-Mediated Apoptosis in HT-29 Human Colorectal Adenocarcinoma Cells", Molecules, 2012.
Publication <1 %

36 Md Abdul Alim, Md Farzanoor Rahman, Tamim Ahsan, Toufiqul Islam, Md Morshedul Alam, Mohammad Nazir Hossain. "Phytochemical profiling, antioxidant potentiality and antibacterial activity of the ethanolic extracts of Roseningea sp. of Bay of Bengal", Journal of Bangladesh Academy of Sciences, 2021
Publication <1 %

37 Yigal Cohen, Kyle M. Van den Langenberg, Todd C. Wehner, Peter S. Ojiambo et al. " <1 %

Resurgence of : The Causal Agent of Cucurbit Downy Mildew ", Phytopathology®, 2015

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off