
Bukti Corespondent Author Judul: identification of bats...



ANNOUNCEMENT FOR ORAL/POSTER PRESENTERS

Dear IC-WICE 2019 presenters,

We would like to gladly thank you for your participation in the International Conference

of Wildlife Trade and Utilization in Wallacea Region (IC-WICE 2019) with the theme:

"Harmonizing Human-Wildlife Relationship for Sustainable Utilization in Wallacea

Region" that was held in August 22nd, 2019. As previously informed, we invited some

interested participants to publish their papers at the conference's proceeding indexed by

Scopus (IOP Conference Series-Earth and Environmental Science). However, the

number of presenters interested in publishing with our IOP proceeding was below 50

articles which also below the contract limit that we offered to IOP Conference Publishing

(>50 articles).

Therefore, as part of our commitment, we would like to publish your article under

different conference's IOP proceeding in 2020. We have collaborated with

International Conference on Food Security and Sustainable Agriculture in the Tropics

(ICFSSAT) (see the Attachment and Select for a Topic) that was held in September 2nd

2019. The costs of publication and editing/formating in the IOP proceeding of ICFSSAT

is IDR 1,500,000 and the payment can be sent to:

BANK BNI 46

Account Name : IC WICE 2019

No:0827895810

before 25th October 2019. The article will be published in 2020. If you required to

publish earlier (still within 2020), we can arrange you to be published under GIESED

2019 IOP Proceeding.

We only require you to submit the article in English with IOP format standards on

15th October 2019 at the latest to: icwice2019@gmail.com. Please fill the form below

and also check your e-mails regularly for further revisions from the reviewers.

Thank you for your cooperation.

Sincerely,

The Chairman of IC-WICE 2019

FORM OF SUBMISSION IOP PROCEEDING

Authour:

Tiltje Andrethe Ransaleleh¹, Meis Jacinta Nangoy¹, Indyah Wahyuni¹, Agustinus Lomboan¹, Ronny Koneri², Suryo Saputro³, Joko Pamungkas³,₄, Alice Latinne⁵

AFFILIATION :

- 1 Departement of Animal Production, Sam Ratulangi University, Manado, Indonesia
 2 Departement of Biology, Sam Ratulangi University, Manado, Indonesia
 3 Primate Research Center, Bogor Agricultural University, Indonesia
 4 Departement of Animal Disease and Veterinary Public Health, Bogo Agricultural University, Indonesia
 5 EcoHealth Alliance, New York

TITLE OF PRESENTATION:

IDENTIFICATION OF BATS ON TRADITIONAL MARKET IN DUMOGA DISTRICT, NORTH SULAWESI

CORRESPONDENCE AUTHOR:

mnangoy@unsrat.ac.id/081241399445, taransaleleh@unsrat.ac.id/ 085211587521

						2-
CLIDANT	TO IOP	Proceeding	of	(Select	one	V)

GIESED IC-FSSAT

Signature

: Tiltje Andretha Ransaleleh Full name

: 30 September 2019 Date





INVITATION FOR CONFERENCE PRESENTER

We would gladly inform you that your abstract for the International Conference of Wildlife Trade and Utilization in Wallacea Region 2019 with the title:

Bats Identification at tradisional Market in Dumoga District North

By

Dr. Ir. Tiltje Andrethe Ransaleleh, M.Si.

Universitas Sam Ratulangi E-mail: taransaleleh@gmail.com

Has been accepted for **Oral/Poster Presentation**. Please bring **Entry Card and Invitation Letter or letter of Acceptence** than give it to the committee for a seminar kit and registration at the International Conference of Wildlife Trade and Utilization in Wallacea Region (IC-WICE) 2019 on August, 22nd 2019 in Dalton Hotel and Convention Makassar. Revised Full Paper is required to be re-submitted on August, 16th-20th 2019 at the latest.

Thank you very much for your participation and looking forward to seeing you at the Conference.



Dr. Risma Illa Maulany, S.Hut, M.Nat.Rest.NIP. 19770317 200501 2 001

Committee Secretariat:
Jln. Perintis Kemerdekaan Km. 10. Kampus Unhas Tamalanrea.
Forestry Faculty, Hasanuddin University Makassar 90245 South Sulawesi, Indonesia.
Telephone/Fax: (+62-411) 589592
Email: icwice2019@gmail.com
http://icwice2019.unhas.ac.id

Sponsored by:









IDENTIFICATION OF BATS ON TRADITIONAL MARKET IN DUMOGA DISTRICT, NORTH SULAWESI

TA Ransaleleh¹, MJ Nangoy¹, I Wahyuni¹, A Lomboan¹, R Koneri¹ S Saputro², J. Pamungkas^{2,3}, A Latinne⁵

¹ Sam Ratulangi University, Manado, Indonesia

Email: taransaleleh@unsrat.ac.id, mnangoy@unsrat.ac.id ssaputra83@gmail.com, latinne@ecohealthallience.org correspondence Author: mnangoy@unsrat.ac.id

Abstract. Fruit bats are used as food by Minahasan tribe in North Sulawesi. This is evident in some traditional markets where various species of fruit bats are sold. As a consequence, bats are continually over hunted for human consumption, threatening bat populations. This study aims to identify bat species sold in traditional markets in Dumoga district, North Sulawesi. Field surveys were performed and data collected in three traditional markets namely Ibolian, Imandi and Dumoga, located nearby the Nani Warta Bone National Park. The collected samples were identified using morphometric measurements and physical characteristics, then analyzed by descriptive method. From these markets were identified and consisted of Acerodon celebensis, Dobsonia exoleta, Neopteryx frosti, Styloctenium wallacei, Rousettus amplexicaudatus, Thoopterus nigrescens, Nyctimene chephalotes, and unknown species. Five of these bat species are endemic to Sulawesi (Acerodon celebensis, Dosonia exoleta, Neopteryx frosti, Styloctenium wallacei, and Thoopterus nigrescens). According to the International Union for Conservation of Nature (IUCN) Red-List, two species have declined status (Styloctenium wallacei Near Threatered (NT), and Acerodon celebensis Vulnarable (VU)), and one species is threatened with extinction status (Neopteryx frosti endangered (EN)).

Key Words: Traditional Market, Bats, Identification

INTRODUCTION

The Minahasa community in North Sulawesi has a culture that consumes bush meat such as warty pigs, rats, and bats (Onibala and Laatung (2007), especially on special occasions and holidays. According to Sheherazade and Susan (2015), around 500 tons of bats are imported to North Sulawesi from other provinces to meet consumer demand. Currently, the consumption of bush meat is popularized with the aim of attracting tourists with "extreme culinary" slogans.

Bats play an important role in the ecosytems and to maintain diversity in forest. They are important animals with ecological functions as seed dispersers, plant pollinators (Liu et al., 2002., Hodgkison et al., 2003., Dumont and O'Neal, 2004., Singaravelan and Marimuthu, 2004) and insect predators. According to Suyanto (2001), there are 205 species in Indonesia, corresponding to 21% of all types of bats in the world. They consist of 72 species from the

² Primate Research Center, Bogor Agricultural University, Indonesia

³ Departement of Animal Disease and Veterinary Public Health, Bogor Agricultural University, Indonesia

⁵EcoHealth Alliance, New York

suborder Megachiroptera (fruit-eating bats, leaves, nectar, and pollen) and 133 species from Microchiroptera suborder (bats that mostly eat insects). There are 23 species of the suborder Megachiroptera found on the island of Sulawesi and surrounding small islands and 11 of these species are endemic (Suyanto 2001, Maryanto & Yani 2003). The results of the study of Ransaleleh et al (2013) on fruit bats in North and Central Sulawesi obtained five species. Two species are endemic namely; *Acerodon celebensis* and *Thoopterus nigrescens*.

Dumoga District, North Sulawesi Province is one of the districts located beside the Nani Warta Bone National Park. The Minahasan people who live in this place consume bat meat as protein resource on a daily basis. Various types of bats can be found in traditional markets. This shows that bats are continually over hunted for human consumption and this activity is one of the major factors threatening bat populations, which will have an impact on the ecosystem balance. Therefore a bat survey has been carried out at the Dumoga District traditional market to identify the types of bats traded. This research is important to implement bat conservation measures, especially for endemic species whose populations are threatened with extinction. We expect that the data and information from this study will be the basis for establishing the rules and policies for bat conservation in Dumoga District by the local government.

RESEARCH METHODS

Surveys in three markets of East Dumoga Subdistrict, namely Dumoga, Imandi, and Ibolian were conducted in this study. The survey was conducted in February 2018 and May 2019. Good communication and relationships were established with the seller so that the seller allows the researcher to identify the bats before they are sold. Parameters measured were body weight, body length, ear length, arm's length under the wings, length of the hind legs of the bat. Observation and documentation for body color and shape. Furthermore bats were identified using a bat identification key from Suyanto (2001).

RESULTS AND DISCUSSION

A total of 170 bats have been identified in three markets, namely Ibolian, Imandi and Dumoga. Identification based on morphometry and color characteristics and body shapes resulted in 9 species from 8 genera: *Acerodon* Jouran, 1837, *Dobsonia* Palmer, 1898, *Nyctimene* Borkhausen, 1797, *Neopteryx* Hayman, 1946, *Stylotecnium* Matschie, 1899, *Cynopterus* Cuvier, 1898, Rousettus Gray, 1821, and *Thoopterus* Matschie, 1899). All of them belong to the family Pteropodidae. Descriptions of 9 species are described as follows:

1. Acerodon celebensis

Body weight 290.21-312.47 g. Body length 180.14-187 mm. Head length 66.59-76.24 mm. Ear length is 30.18-30.29 mm. The length of the forearm is 122.86-123.81 mm. The length of the rear leg is 37.95 mm. Golden yellow color on the feathers of all parts of the body and wing fingers. Brown color on the snout, wings and toes. (Figure 1). This species has no tail.

Comment [AL1]: You have to add the dates when the surveys were performed.



Figure 1. Acerodon celebensis

2. Dobsonia exoleta.

Body weight 282.21-320 g. Body length 171-220 mm. Head length of 52 mm. Ear length 25-26.66 mm. The length of the forearm is 110-116.67 mm. Ear length 11-14.66 mm, Length of the rear leg 33-33.33 mm. Greenish yellow color on body hair. Black color on wing skin. Hairless back. (Figure 2). This species has a tail and no claws on the fingers of the two wings.



Figure 2. Dobsonia exoleta.

3. Neopteryx frosti

Body weight 230 g. Body length 175 mm. Head length of 65 mm. Ear length 21 mm. The length of the forearm is 110 mm. The back foot length is 30 mm. Light brown color on the feathers on the body and wings. Shaped mesh on the dark brown stripe-shaped wing. In the head of the lateral frontal side white lines are found from the bottom of both eyes to the mouth (Figure 3). This type is tailed and has no claws on the wings. According to Bergmans and Rozendal (1988) cited by Suyanto (2001), this type weighs 164-190 grams. Head length from 50.7 to 55.8 mm, wing length under wing 104.9-110.6 mm.



Figure 3. Neopteryx frosti

4. Styloctenium wallacei

Body weight 170-210 g. Body length 160 mm. Head length 54-56 mm. Ear length 21-25 mm. The length of the forearm under 100-110 mm. The back foot length is 30 mm. Reddish brown color (cinnamon brown) on the entire body except the wing ears and legs dark brown. There are short white lines above the eyes and the circumference of the eyes, nose, snout and mouth of the left and right sides. (Figure 4).



Figure 4. Styloctenium wallacei

5. Nyctimene cephalotes

Body weight 57.71-58.75 g. Body length 84.44-86.75 mm. Head length 30.22-31.5 mm. Ear length 14.22-14.5 mm. The length of the forearm is 65.55-66.5 mm. Tail length 20.66-21.5. Length of back legs 14.11-14.5 mm. Grayish brown color on body hair. Yellow spots on the ears, wings, and fingers of the wings. Tubular protruding nose. Thin brown line along the back. Short tail (Figure 5).



Figure 5. Nyctimene cephalotes

6. Cynopterus minutus

Body weight of 45-50 g. Body length 80-85 mm. Head length 26.5-28 mm. Ear length 12.5-14 mm. The length of the forearm is 55 mm. Back foot length 10-12 mm. Grayish brown color on body hair. Short snout. White line at the edge of the ear (Figure 6).



Figure 6. Cynopterus minutus

7. Thoopterus nigrescens

Body Weight 81.32-93.69 g. Body length 110.03-112.69 mm. Head length from 41.5 to 42.92 mm. Ear length 14.74-15.23 mm. The length of the forearm is 73.21-74.77mm. The length of the rear leg is 16,19-16,31 mm. Gray black body, no tail, short snout (Figure 7)



Figure 7. Thoopterus nigrescens

8. Rousettus amplexicaudatus

Body weight ranges from 72.75 to 75.63 g. Body length 109.25-114.69 mm. Head length of 45.59-47.63 mm. Ear length of 15.86-16.16 mm. The length of the forearm is 71.66-72.75 mm. Tail length 18.94 mm. The length of the rear leg is 17.66-18.38 mm. The body is grayish brown, the snout is long, has a blackish-brown tail, wings (Figure 8).



Figure 8. Rousettus amplexicaudatus

9. Rousettus sp.

Body weight ranges from 150-216.66 g. Body length 95-165 m. Head length of 39-53.3 mm. Ear length is 25.3-26 mm. The length of the forearm is 85-96.66 mm. Head length of 22-29.66 mm. According to body weight and morphometric size, this bat is not *Rousettus amplexicaudatus*. However, some other physical features are similar to *Rousettus amplexicaudatus* (Figure 9)



Figure 9. Rousettus sp.

Identification results show seven species of small fruit bats, namely: *Dobsonia exoleta, Neopteryx Frosti, Styloctenium wallacei, Rousettus amplexicaudatus, Thoopterus nigrescens, Nyctimene cephalotes,* and *Cynopterus minutus*. The seven bats are hunted in and around the Nani Warta Bone forest area and the Gunung Moosi community forest estate, near Dumoga sub-district. Table 1 shows the numbers and percentages of bats observed in these markets. *Thopterus nigrescens, Rousettus amplexicaudatus, Acerodon celebensis, Nyctimene cephalotes* were the most common species (30%, 28%, 25% and 7.64%, respectively) whereas, *Dobsonia exoleta, Cynopterus minutus, Styloctenium wallacei, Neopteryx frosti* and the unidentified species of *Rousettus* were less frequently observed (2.35%, 1.76%, 1.17%, 0.5%, and 2.94%, respectively).

Table 1. Total, Percentage, Conservation Status (IUCN_Red List) and endemic status for each species

No	Species	Total	%	Conservation Status	Remaks
		Individu		(IUCN Red-List)	
1	Acerodon celebensis	43	25	Vulnerable (VU)	Endemic
2	Dobsonia exoleta	4	2,35	Least Concern (LC)	Endemic
3	Neopteryx frosti	1	0,59	Endangered (EN)	Endemic
4	Styloctenium wallacei	2	1,17	Near Threatened (NT)	Endemic
5	Rousettus amplexicaudatus	48	28,23	Least Concern (LC)	
6	Thoopterus nigrescens	51	30,00	Least Concern (LC)	Endemic
7	Nyctimene cephalotes	13	7,64	Least Concern (LC)	
8	Cynopterus minutus	5	1,76	Least Concern (LC)	
9	Rousettus sp	4	2,94	Least Concern (LC)	

DISCUSSION

Of the nine species identified in this study, five are endemic to Sulawesi (Acerodon celebensis, Dobsonia exoleta, Neopteryx frosti, Styloctenium wallacei and Thoopterus nigrescens). According to the International Union for Conservation of Nature (IUCN) Red-List 2019, Styloctenium wallacei is Near Threatered (NT), Acerodon celebensis is Vulnerable (VU), and Neopteryx frosti is endangered (EN). The existence of the bat trade in the Dumoga subdistrict traditional market shows that people's understanding of the role of bats in the ecosystem is not yet apparent. Actions need to be taken to carry out bat conservation activities, especially for endemic species and species thought to be endangered. Conservation education for school-age children and continuous education for hunters, sellers, and bat-eaters is necessary. Another way to avoid declining bat populations would to breed them and legalize the trade of the cultivated bat meat following the prohibition on the trade of wild bats from hunting.

CONCLUSION

Eight species were identified in Dumoga traditional markets: Acerodon celebensis, Dobsonia exoleta, Neopteryx Frosti, Styloctenium wallacei, Rousettus amplexicaudatus, Thoopterus nigrescens, Nyctimene cephalotes, and Cynopterus minutus and one unidentified species from the Rousettus genus. Five of these species, A.Celebensis, Dobsonia exoleta, N. frosti, S.wallacei and T.nigrescens, are endemic to Sulawesi. The population of Neopteryx frosti is threatened.

SUGGESTIONS

Promoting campaigns and education about the ecological functions of bats through various social media as well as school, church, and local government media and related agencies needs to be done. We also propose to establish a capture quota and a ban on wild capture by the competent authority.

ACKNOWLEDGEMENT

This work was supported by Fundamental Research Sam Ratulangi University Grant 2019 (Dr. Meis Jacinta Nangoy and Tim by PNBP Unsrat). Travel Grant for International conference of Wildlife Trade and Utilization in Wallacea Regian supported by PREDICT Indonesia under the EPT-2 Program.

REFERENCE

_____The IUCN Red List of Threatened spesies. 2019 https://www.iucnredlist.org/species/14560/115122474. di download 7 Agustus 2019

Dumont RE, O'Neal R 2004 Food hardness ang feeding behavior in old world fruit bats (Pteropodidae). *Journal of Mammalogy* **85**(1) 8-14

Hodgkison R, Balding ST, Zubaid A, Kunz TH 2003 Fruit bats (Chiroptera: Pteropodidae) as seed dispersers and pollinator in a lowland Malaysian rain forest. *Biotropica* **35** 491-502

Liu A-Z, Li D-Z, Wang H, Kress WJ 2002 Ornithophilous and chiropterophilous pollination in musa itinerans (Musaceae), a pioneer species in tropical rain forest of Junnan, Southwestern China. *Biotropica* **34** 254-260

Maryanto I and Mohamad Y 2003. A new spesies of *Rousettus* (Chiroptera: Pteropodidae) from Lore Lindu, Central Sulawesi *Mammal Study* **28** 111-120

Onibala JSIT, Laatung S 2007. Bushmeat hunting in North Sulawesi and related conservation strategies (a case study at the Tangkoko Nature Reserve) *I. Agric. Rul. Dev. Trop. Subtrop.* **90** 110-116

Ransaleleh TA, Maheswari RRA, Sugita P, Manalu W 2013 Identifikasi kelelawar pemakan buah asal Sulawesi berdasarkan morfometri *Jurnal Veteriner* **14**(4) 485-494

Sheherazale and Tsang SM 2015 Quantifying the bat bushmeat trade in North Sulawesi, Indonesia, with suggetions for conservation action *Global Ecology Conservation* **3** 324-330

Singaravelan N and Marimuthu G 2004 Nectar feeding and pollen carrying from ceibe pentandra by pteropodid bats *Journal of Mammalogy* **85**(1) 1-7

Suyanto A 2001 *Kelelawar Di Indonesia Seri Panduan Lapangan* Pusat Penelitian dan Pengembangan Biologi LIPI Bogor Indonesia chapter 6 pp 31-57