

# Molecular identification of bacteria isolated from culture medium of the gold-lipped pearl oyster *Pinctada maxima* larvae

*by* Elvy Like Ginting 23

---

**Submission date:** 01-Mar-2021 04:04PM (UTC-0800)

**Submission ID:** 1521753008

**File name:** edium\_of\_the\_gold-lipped\_pearl\_oyster\_Pinctada\_maxima\_larvae.pdf (365.88K)

**Word count:** 6344

**Character count:** 35102

## Molecular identification of bacteria isolated from culture medium of the gold-lipped pearl oyster *Pinctada maxima* larvae

STENLY WULLUR<sup>1,\*</sup>, HATOPAN NAPITUPULU<sup>1</sup>, ELVY LIKE GINTING<sup>1</sup>,  
NOLDY GUSTAF FRANS MAMANGKEY<sup>1</sup>, LETHA LOUISIANAWANTANIA<sup>1,2</sup>,  
RADOSLAV SMOLAK<sup>3</sup>, ERICK OGELLO<sup>4</sup>

<sup>1</sup>Faculty of Fisheries and Marine Science, Universitas Sam Ratulangi, Jl. Kampus, Bahu, Malalayang, Manado 95115, North Sulawesi, Indonesia  
Tel./fax.: +62-431-868027, \*email: stenlywullur@unsrat.ac.id

<sup>2</sup>Ichthyology section, Zoologisches Forschungsmuseum Alexander Koenig, Adenauerallee 160, D-53113 Bonn, Germany

<sup>3</sup>Department of Ecology, Faculty of Humanities and Natural Sciences, Presov University, 081 16, Presov, Slovakia

<sup>4</sup>Department of Fisheries and Natural Resources, Maseno University, Kondele Kisumu-Busia Rd Maseno, Kisumu, Kenya

Manuscript received: 31 July 2020. Revision accepted: 20 October 2020.

**Abstract.** Wullur S, Napitupulu H, Wantania LL, Ginting EL, Mamangkey NGF, Smolak R, Ogello E. 2020. Molecular identification of bacteria isolated from culture medium of the gold-lipped pearl oyster *Pinctada maxima* larvae. *Biodiversitas* 21: 5291-5297. This study was conducted for the molecular identification of bacteria species isolated from culture medium of the gold-lipped pearl oyster, *Pinctada maxima* larvae. The pearl oysters were cultured using live-microalgae (*Isochrysis* sp) and fish waste diet (FWD) as food sources. Bacteria were isolated from the oyster larvae and identified based on the 16S rRNA gene sequence. The isolated bacteria were grown on agar plate and incubated at 37°C for 24 to 48 hours. Representative colonies of the bacteria were selected and cultured for molecular analysis. The 16S rRNA genes of the bacteria were amplified and the sequences were matched with the NCBI GenBank database. Seven different colonies were observed based on morphological characters. Similarity test by conducting the Basic Local Alignment Search Tool (BLAST) in the NCBI GenBank database, using the 16S rRNA gene sequences showed that the seven isolates colonies possess high similarity to five bacteria species i.e. *Pseudomonas pachastrellae*, *Vibrio alginolyticus*, *Bacillus filamentosus*, *Bacillus cereus* and *Idiomarina fontislapidosi* belonging to four different genera i.e. *Bacillus*, *Staphylococcus*, *Vibrio*, and *Alteromonas*.

**Keywords:** 16S rRNA, bacteria, fish-waste diet, *Pinctada maxima*, gold-lipped pearl oyster

### INTRODUCTION

Pearl oyster culture industry in Indonesia mostly uses oyster seeds collected from the wild. If not managed well, the wild harvesting can potentially destroy oyster population and may affect the sustainability of the national pearl industry (Manez et al. 2010). This is demonstrated by the Segara Anakan lagoon in South Central Java, Indonesia, where the flourishing pearl oysters *Pinctada maxima*/*Pinctada margaritifera* are depleting due to overharvesting (Manez, 2010). In order to maintain stock of oyster in the wild, several private hatcheries commenced the production of the larvae (*P. maxima*), and the juveniles are restocked to the wild for further adult collection at suitable size prior to seeding for pearl production (Kvingedal et al. 2010).

In the process of culturing, however, hatcheries use live microalgae and are cultivated on-site as exclusive food nutrition for the larvae. In fact, the production cost of the live microalgae culture is about 30–50% of hatchery operational cost. Also, technical sources and skilled personnel are required for successful production of live microalgae with adequate quantity and high quality. However, the unavailability of these requirements is common problem for oyster hatchery operations (Tanyaros et al. 2016; Southgate et al. 2016). For this reason, cheap and easily handled food sources based on baker's yeast were

introduced. This resulted in stunted growth and high mortality of the oyster because of nutritional deficiency of the yeasts. This resulted in the growth and high mortality of oysters as there was nutritional deficiency of yeast (Tanyaros et al. 2016). Furthermore, non-living microalgae such as powder and paste were used for larval rearing of oysters (Wassnig and Southgate, 2016). The use of non-living microalgae promotes growth and survival of reared oyster larvae, which is equivalent to live-microalgae (Southgate et al. 2016). However, individual cells of the non-living microalgal-based food are non-motile, negatively buoyant, and eventually settled onto the bottom which is the source of culture crash. Also, the powder microalgae insufficient nutrition due to drying process, while the paste product has short shelf life (Wassnig and Southgate, 2016). The prices of these two non-living microalgal-based foods are expensive, therefore they are preferred by hatcheries in developing countries (Wullur et al. 2020). This led to the development of an inexpensive and easy diet as an alternative food for microalgae, the fish waste diet (FWD) (Ogello et al. 2018; Napitupulu et al. 2019; Ogello et al. 2019, 2020; Wullur et al. 2019, 2020). The FWD was developed from the composition of fish waste and probiotic bacteria. Attempts to feed rotifers with this diet were shown to have a high level of acceptance, as they were harvested at densities of about 2000 to 3000 ind./mL every week during semi-continuous culture

(Ogello et al. 2018; 2019, 2020; Wullur et al. 2017, 2019). In addition, it was suspected that fish waste contains bacteria used by rotifers as their nutritional source (Wullur et al. 2020).

Pearls oysters are reported to feed on bacteria in natural environment. However, bacteria are present in the rearing water of oyster and microalgae culture in hatcheries, it provides metabolic requirements for oyster's larvae by making direct or direct provision of organic molecules and exoenzyme required for the growth and survival of larvae (Karim et al. 2013). On the other hand, the presence of opportunistic bacteria often causes disease massive larval mortality in oyster hatcheries (Saulnier et al. 2010). However, the use of probiotics, which provides health benefits and protection against bacterial pathogens, is essential to maintain a healthy culture environment (Karim et al. 2013). Today, the bacterial population dynamics in live cultures for *P. maxima* is not clear. Furthermore, the bacterial population identification analysis should be more important when additional diets such as FWD are used. The knowledge on bacteria present in living cultures is important for designing nutritional strategies for oyster culture. In this study, a molecular approach was used to identify the species of bacteria present in rearing water of oyster culture. Also, a small-ribosomal subunit 16S rRNA gene was sequenced to identify bacterial species.

## MATERIALS AND METHODS

The larvae of pearl oyster *P. maxima* (14-days old) were cultured in 700 mL seawater (salinity 35 ppt) at a density of about 33 ind./mL for two weeks, using transparent plastic containers. Each container was equipped with aeration of 0.66 mL/min and placed in a room at 25-26°C. The larvae were fed with live-microalgae *Isochrysis* sp. (PBM) and a FWD, which was previously soaked with EM4® (EM) for 24 hours. The preparation of the FWD was adapted as described in Patent No. P00201609066 registered in Indonesia.

The bacteria were isolated from the rearing water of oyster culture at the end of rearing period, with a sample of approximately 1 mL serially diluted (10 to 1000-fold) in sterile seawater. The samples were spread on 2 % nutrient agar and incubated at 37°C for 24-48 hours to observe bacterial colonies. A distinct colony was further selected according to their phenotype characters and grown in nutrient broth for 24 - 48 hours to obtain pure bacterial strains. Bacteria were harvested and centrifuged at 14,000 rpm for 5 minutes for molecular analysis.

The Qiaprep Miniprep Kit (Qiagen, Herten, Germany) was used to extract genomic DNA and 16S rRNA gene from the genomic DNA of bacteria was amplified using two primers (8F; forward, 5'-AGAGTTTGTATCCTGGCTG-3' and 1492R; reverse 5'-ACCTTGTACGACTT-3'). The PCR reaction was prepared with a total volume of 25 µl which consist of 1 µl sample, 5 µl of 5x Hot fire pool, 1 µl of each primer and 17 µl of ddH<sub>2</sub>O. It was then placed in a Professional Thermocycler (Biometra, Analytik Jena) for 35 cycles

amplification at 95°C for 6 min (1 cycle), 95°C for 30 sec, 52°C for 30 sec, and 72°C for 30 sec (34 cycles), followed by 72°C for 10 min as final cycle.

The amplicons and the primer pairs were sent to First-Base Co., Selangor, Malaysia to sequence the 16S rRNA gene. Sequencing procedures were conducted bi-directionally using a BigDye® Terminator v.3.1 Cycles Sequencing Kit (Applied Biosystems, USA) and further read with an ABI PRISM® 377 automated DNA sequencer. Its quality was accessed using the Sequence Scanner version 2.0 software (Applied Biosystem). Subsequently, sequences were trimmed, assembled, and manually edited using Geneious Prime version 2020 (<http://www.geneious.com>, Kearse et al. 2012) prior to the application of BLAST (Basic Local Alignment Tools) analysis at The National Center for Biotechnology Information (NCBI, <https://www.ncbi.nlm.nih.gov/>).

## RESULTS AND DISCUSSION

In this study, five species of bacteria (*Pseudomonas pachastrellae*, *Vibrio alginolyticus*, *Bacillus filamentosus*, *Bacillus cereus*, and *Idiomarina fontislapidosi*) were identified, belongs to 4 genera (*Pseudomonas*, *Vibrio*, *Bacillus*, and *Idiomarina*). These species obtained from rearing water of oyster larval culture were analyzed from seven isolates by conducting nucleotide BLAST using 16S rRNA gene at NCBI. The percent identity of the first top nearest species of 5 isolates (Table 1) was PBM21233 (99.07%), PBM31432 (99.64%), EM114133 (99.21%), EM11441 (99.01%) and EM11442 (98.87%), above the threshold value, while 2 isolates EM114233 (98.30%) and EM31042 (98.58%) were slightly below the threshold value. According to Kim et al. (2014), the threshold value to differentiate bacterial species using 16S rRNA gene has 98.65% sequence similarity/percent identity.

Three isolates namely PBM21233, EM114233, and EM11441 collected from the two feeding treatments (PBM and EM) were identified as *P. pachastrellae* (NR 040991.1) belongs to the genus of *Pseudomonas*. The members of this genus inhabit a variety of environments, including terrestrial and aquatic habitats. According to Özen and Ussery (2012), the total number of species currently included in the genus *Pseudomonas* is around 202 species, which are important in medical or biotechnological applications. Several species such as *Pseudomonas aeruginosa* (Priyaja, 2012; Giri et al. 2012; Vinoj et al. 2013), *Pseudomonas fluorescens* (Nour and El-Ghiet, 2011; Eissa et al. 2014), *Pseudomonas aestumarina*, *Pseudomonas synxantha*, *Pseudomonas chlororaphis* (Sánchez et al. 2014) have reported as a potent probiotic microorganism for marine or freshwater aquaculture. In addition, *P. aeruginosa* improved the innate immunity as well as survival of carp *Labeo rohita* against *Aeromonas trophila* infection (Giri et al. 2012) and enhanced survival of pearl oyster *Pinctada mazatlanica* (Macias et al. 2010). Also, *P. fluorescens* reported to improve survival of rainbow trout *Oncorhynchus mykiss* after challenging the fish with pathogen *Vibrio anguillarum*, while *P.*

48  
*aestumarina* conferred protection to white shrimp *Litopenaeus vannamei* against vibriosis (Sánchez et al. 2014). The combination of *P. synxantha* and *P. aeruginosa* improved growth, survival and the immune responses of western king prawns *Penaeus latissulcatus* juveniles, while *P. chlororaphis* enhances resistance of European perch *Perca fluviatilis* against *Aeromonas sobria* infection (Sánchez et al. 2014). On the contrary, *Pseudomonas plecoglossicida* (Nakai, 2010), *Pseudomonas gessardii* (Hamza et al. 2018), *Pseudomonas alcaligenes* (Xu et al. 2015), *Pseudomonas beatica* (Lopez et al. 2012), *Pseudomonas koreensis* (Shahi and Mallik, 2014) and *Pseudomonas putida* (Mao et al. 2012) caused infection in aquaculture. Nakai (2010) and 10  
 g et al. (2019) reported on the mass mortality of Ayu *Plecoglossus*

*altivelis* and large yellow croaker *Pseudosciaena crocea* resulting from *P. plecoglossicida* attacks, while *P. gessardii* and *P. alcaligenes* facilitated the total mortality of freshwater crayfish *Pacifastacus leniusculus* (Korkut et al. 2018) and Chinese sturgeon *Acipenser sinensis* (Xu et al. 2015), respectively. The *P. beatica* was reported to be present in disease outbreak for flatfish *Dicologlossa cuneata* (Lopez et al. 2012), while *P. koreensis* is present in eye infections of golden mahseer *Tetraodon lineatus* (Shahi and Mallik, 2014). Furthermore, *P. putida* has been reported as an important fish pathogen of rainbow trout *Oncorhynchus mykiss*, European eel *Anguilla anguilla*, oyster toadfish *Opsanus tau*, and large yellow croaker *Pseudosciaena crocea* (Mao et al. 2012).

**Table 1.** Five top hits of the nearest species of the bacterial isolates after nucleotide BLAST against 16S rRNA gene in the NCBI GenBank using rRNA type strains/prokaryotic 16S ribosomal RNA (bacteria and archaea) database

Isolates	Nearest species (accession number)	Score	Query cover	E-value	Identity
PBM21233	<i>Pseudomonas pachastrellae</i> (NR040991.1)	2516	100%	0.0	99.07%
	<i>Pseudomonas oceani</i> (NR152090.1)	2516	100%	0.0	97.79%
	<i>Pseudomonas aestusnigri</i> (NR126210.1)	2412	99%	0.0	97.79%
	<i>Pseudomonas jilinensis</i> (NR169366.1)	2289	100%	0.0	96.16%
	<i>Pseudomonas sabulinigri</i> (NR044415.1)	2259	100%	0.0	95.79%
PBM31432	<i>Vibrio alginolyticus</i> (NR113781.1)	2571	99%	0.0	99.64%
	<i>Vibrio alginolyticus</i> (NR117895.1)	2569	99%	0.0	99.64%
	<i>Vibrio alginolyticus</i> (NR121709.1)	2564	99%	0.0	99.57%
	<i>Vibrio alginolyticus</i> (NR118258.1)	2564	99%	0.0	99.57%
	<i>Vibrio natriegens</i> (NR117890.1)	2564	99%	0.0	99.57%
EM114133	<i>Bacillus filamentosus</i> (NR134701.1)	2521	98%	0.0	99.21%
	<i>Bacillus endophyticus</i> (NR025122.1)	2518	99%	0.0	98.73%
	<i>Bacillus shackletonii</i> (NR025373.1)	2224	99%	0.0	95.00%
	<i>Bacillus circulans</i> (NR112632.1)	2211	99%	0.0	94.92%
	<i>Bacillus circulans</i> (NR104566.1)	2207	99%	0.0	94.98%
EM114233	<i>Pseudomonas pachastrellae</i> (NR040991.1)	2468	99%	0.0	98.30%
	<i>Pseudomonas oceani</i> (NR152090.1)	2362	99%	0.0	97.02%
	<i>Pseudomonas aestusnigri</i> (NR126210.1)	2353	99%	0.0	96.88%
	<i>Pseudomonas jilinensis</i> (NR169366.1)	2235	99%	0.0	95.39%
	<i>Pseudomonas sabulinigri</i> (NR044415.1)	2202	99%	0.0	95.02%
EM11441	<i>Pseudomonas pachastrellae</i> (NR040991.1)	2527	100%	0.0	99.01%
	<i>Pseudomonas oceani</i> (NR152090.1)	2425	100%	0.0	99.74%
	<i>Pseudomonas aestusnigri</i> (NR126210.1)	2409	98%	0.0	97.99%
	<i>Pseudomonas jilinensis</i> (NR169366.1)	2298	100%	0.0	97.12%
	<i>Pseudomonas sabulinigri</i> (NR044415.1)	2266	99%	0.0	97.81%
EM11442	<i>Bacillus cereus</i> (NR115526.1)	2516	100%	0.0	98.87%
	<i>Bacillus proteolyticus</i> (NR157735.1)	2516	100%	0.0	98.87%
	<i>Bacillus wiedmannii</i> (NR152692.1)	2516	100%	0.0	98.87%
	<i>Bacillus cereus</i> (NR114582.1)	2516	100%	0.0	98.87%
	<i>Bacillus cereus</i> (NR115714.1)	2516	100%	0.0	98.87%
EM31042	<i>Idiomarina fontislapidosi</i> (NR029115.1)	2481	99%	0.0	98.58%
	<i>Idiomarina baltica</i> (NR027560.1)	2442	99%	0.0	98.08%
	<i>Idiomarina aquatica</i> (NR144584.1)	2364	99%	0.0	97.09%
	<i>Idiomarina seosinensis</i> (NR025826.1)	2263	98%	0.0	96.57%
	<i>Idiomarina abyssalis</i> (NR024891.1)	2254	99%	0.0	95.66%

So far there are no reports about the role of *P. pachastrellae* in aquaculture. It was first isolated from marine sponge specimen and assigned as a new species in 2005 (Kennedy et al. 2014). *P. pachastrellae* was further identified in the polluted sand samples after a marine oil spill (Lamendella et al. 2014), as well as deep seawater (depth of 1390 m) (Wang and Sun, 2016), and also in soil samples collected from an oily environment in close proximity to the seashore (Kaskatepe et al. 2017). Apart from having highest sequences identities with *P. pachastrellae*, the isolates PBM21233, EM114233 and EM11441 share high sequences identities with *Pseudomonas oceani* (97.79, 97.02, and 99.74% identities, respectively (Table 1)). Wang et al. (2020) placed the two closest relate *Pseudomonas* species in the group of *Pseudomonas pertucinogena* lineage. The *P. oceani* inhabiting deep seawaters (Wang and Sun, 2016) and oily polluted environment (Li et al. 2020) which is similar with *P. pachastrellae*. This study showed for the first time the presence of *P. pachastrellae* in the rearing water of the gold-lipped pearl *P. maxima* larvae which was found in both PBM and EM treatments.

Isolate PBM31432 collected from PBM treatment showed percent identity value above the threshold which referred to *Vibrio alginolyticus* (NR113781.1), a species of genus *Vibrio*. The species in this genus inhabit the aquatic environment. They are judged to be responsible for vibriosis, implicated in economic losses to the aquaculture industry, and is also frequently associated with disease-causing microorganism in humans (Ashrafudoulla et al. 2020). In addition, *V. alginolyticus* is implicated in severe soft tissue infections, sepsis, and other extraintestinal infections in human, through water exposure or the consumption of contaminated seafood (Slifka et al. 2017). This species together with *Vibrio parahaemolyticus*, *Vibrio harveyi*, *Vibrio owensii* and *Vibrio campbelli* were predominantly associated as a causative agent of vibriosis on farmed aquatic animals (Ina et al. 2019). Specifically, *V. alginolyticus* were more frequently reported in the prevalence of many farmed fish species including olive flounder *Paralichthys olivaceus*, black rockfish *Sebastes schlegelii*, red sea bream *Pagrus major*, and sea bass *Micropterus japonicus*, kelp grouper *Epinephelus bruneus*, Orange-spotted grouper *Epinephelus coioides* (Oh et al. 2011; Harikrishnan et al. 2012; Wang et al. 2014). The *V. alginolyticus* was also reported in the infection of farmed shrimp (i.e. white shrimp *Litopenaeus vannamei*, shrimp *Marsupenaeus japonicus* (Kitikew et al. 2013; Zhu et al. 2016), abalone (i.e. small abalone *Haliotis diversicolor*, red abalone *Haliotis rufescens* (Wu et al. 2011) and farmed pearl oyster (i.e. Akoya pearl oyster *Pinctada fucata*, *Pinctada martensii*, *Pinctada maxima* (Wang et al. 2012; 2016; Adzigbli et al. 2020)). The *V. alginolyticus* in this study was only identified in PBM and not in EM treatments. Based on the consideration as a pathogen in the competitive processes, this species is suggested to be possibly displaced by beneficial bacteria in EM.

Two isolates, EM114133 and EM11442, which were collected from EM feeding treatment showed percent

identity value above threshold referring to species of *Bacillus filamentosus* (NR134701.1) and *Bacillus cereus* (NR 115526.1) respectively. Species of bacteria in this genus are often referred to as probiotic microorganisms, as they provide health benefits to the host. According to Cutting (2011), genus *Bacillus* was used for decades as probiotic medicine or dietary supplements for human or veterinary use. In aquaculture, they play an important role in modulating hepatic indexes, gene expression, antioxidant and digestive enzymes. They also enhance water quality, disease resistance and immunity to promote better growth and survival of the reared fish or larvae (Kuebutomye et al. 2019). Particularly, *B. cereus* has generally been recognized as a human pathogen due to their role in food poisoning. In aquaculture, however, *B. cereus* has a significant role in enhancing immune responses, growth, or disease resistance to cultured fish (i.e. rainbow trout *Oncorhynchus mykiss*, silver catfish *Rhamdia quelen*, catfish *Clarias gariepinus*, Asian sea bass *Lates calcarifer*, Crucian Carp *Carassius auratus*, tambaqui *Colossoma macropomum*, Pengze crucian carp *Carassius auratus* var. *Pengze*, grouper *Epinephelus lanceolatus* and *Epinephelus fuscoguttatus* (Orsod et al. 2012; Souza et al. 2012; Gisbert et al. 2013; Hapsari, 2016; He et al. 2017; Zhang et al. 2019; Yang et al. 2019), cultured shrimp (i.e. giant tiger prawn *Penaeus monodon*, pacific white shrimp *Litopenaeus vannamei*, pink shrimp *Farfantepenaeus brasiliensis*, whiteleg shrimp *Penaeus vannamei* (De Souza et al. 2012; Chandran et al. 2014; Hao et al. 2014; Khademzade et al. 2020)), and other aquatic organisms (i.e. sea cucumber *Apostichopus japonicus*, hooded oyster *Saccostrea cucullata*, mud crab *Scylla paramamosain* (Umayaparvathi et al. 2013; Wu et al. 2014; Zhao et al. 2016). However, there have been no reports of the presence of *B. filamentosus* in aquaculture. It was first isolated from a marine sediment in India and was assigned as a new species in 2015 (Sonalkar et al. 2015). Further, the *B. filamentosus* was reported as a plant growth promoting bacteria on wheat *Triticum aestivum* grown in a saline environment (Khalifa et al. 2020). Interestingly, two species of *Bacillus* i.e. *B. cereus* and *B. filamentosus* in this study were only found in EM treatment, which shows an opportunity for using probiotic supplements to supply good bacteria as probiotics in oyster larvae culture.

Isolate EM31042 from EM feeding treatment showed percent identity value lower than the threshold referring to species *Idiomarina fontislapidosi* (NR 029115.1), belongs to the genus *Idiomarina*. This genus is a member of the family *Idiomarinaceae* proposed by Innova et al. (2000), to accommodate true marine bacterial species in the group of *Alteromonas*-like  $\gamma$ -*Proteobacteria*. The genus *Idiomarina* currently consists of about 27 species and are typically euryhalophiles thriving in wide salinity ranges of marine environment (Liu et al. 2018). These include coastal and oceanic waters and sediments (Zhang et al. 2012; Hameed et al. 2016), saline-alkaline soil (Leon et al. 2015), inland hypersaline wetlands (Zhong et al. 2014), solar salt-making works (Lee et al. 2015) and deep sea sediment (Du et al. 2015). In aquaculture, bacteria in this genus live in a

recirculating aquaculture system with other heterotrophic bacteria (Michaud et al. 2009; Garcia et al. 2019; Teitge et al. 2020). They are also found as microbiome in fish gut (Yukgehnash et al. 2020), in brine shrimp *Artemia franciscana* (Riddle et al. 2013), in a microalgal mass culture for finfish hatchery (Sandhya et al. 2017). However, their role in aquaculture, whether as a pathogen or probiotic is not well known.

In conclusion, this study shows that the rearing water of oyster larval culture contains at least 5 different species of bacteria, which are *P. pachastrellae* (PBM21233, EM114233, and EM11441), *V. alginolyticus* (PBM31432), *B. filamentosus* (EM114133), *B. cereus* (EM11442) and *I. fontislapidosi* (EM31042) belonging to the genus of *Pseudomonas*, *Vibrio*, *Bacillus* and *Idiomarina*. The species of bacteria in the treatment of PBM were *P. pachastrellae* and *V. alginolyticus*, while those in the treatment of EM were *P. pachastrellae*, *B. filamentosus*, *B. cereus* and *I. fontislapidosi*. In addition, EM treatment is suspected to involve bacterial species in the process of diet decomposition and subsequently became a food source or decomposed organic material as nutrition for the oyster larvae

#### ACKNOWLEDGEMENTS

The authors are grateful to the Indonesian Ministry of Research and Technology (Kemenristek)/National Agency for Research and Innovation (BRIN) Indonesia and Sam Ratulangi University, Manado, Indonesia for funding this study, and also to the pearl oyster hatchery, PT. Arta Samudera Bitung, for providing the pearl oyster larvae.

#### REFERENCES

- Adzighli L, Hao R, Jiao Y, Deng Y, Du X, Wang Q, Huang R. 2020. Immune response of pearl oysters to stress and diseases. *Rev Aquacult* 12: 513-523.
- Ashrafudoulla M, Mizan MFR, Park SH, Ha SD. 2020. Current and future perspectives for controlling *Vibrio* biofilms in the seafood industry: a comprehensive review. *Crit Rev Food Sci Nutr*. DOI: 10.1080/10408398.2020.1767031
- Cutting SM. 2011. *Bacillus* probiotics. *Food Microbiol* 28: 214-220.
- De Souza DM, Suita SM, Leite FPL, Romano LA, Wasielesk W, Ballester ELC. 2012. The use of probiotics during the nursery rearing of the pink shrimp *Farfantepenaeus brasiliensis* (Latreille, 1817) in a zero exchange system. *Aquacult Res* 43 (12): 1828-1837.
- Dias JA, Abe HA, Sousa NC, Couto MV, Cordeiro CA, Meneses JO et al. 2018. Dietary supplementation with autochthonous *Bacillus cereus* improves growth performance and survival in tambaqui *Colossoma macropomum*. *Aquaculture Research* 49: 3063-3070.
- Du J, Lai Q, Liu Y, Du Y, Liu X, Sun F, Shao Z. 2015. *Idiomarina atlantica* sp. nov., a marine bacterium isolated from the deep sea sediment of the North Atlantic Ocean. *Antonie Van Leeuwenhoek* 107: 393-401.
- Eissa N, Abou El-Gheit N, Shaheen AA. 2014. Protective effect of *Pseudomonas fluorescens* as a probiotic in controlling fish pathogens. *Am J Biosci* 2: 175-181.
- Garcia MME, Cáceres J, Vásquez R, Cruz, R. 2019. Bacteriological water quality of recirculating aquatic systems for maintenance of yellowtail amberjack *Seriola lalandi*. *J World Aquacult Soc* 50: 934-953.
- Giri SS, Sen SS, Sukumaran V. 2012. Effects of dietary supplementation of potential probiotic *Pseudomonas aeruginosa* VSG-2 on the innate immunity and disease resistance of tropical freshwater fish, *Labeo rohita*. *Fish Shellfish Immunol* 32: 1135-1140.
- Gisbert E, Castillo M, Skalli A, Andree KB, Badiola I. 2013. *Bacillus cereus* var. *toyoi* promotes growth, affects the histological organization and microbiota of the intestinal mucosa in rainbow trout fingerlings. *J Anim Sci* 91: 2766-2774.
- Hameed A, Lin SY, Lai WA, Shahina M, Liu YC, Hsu YH, Young CC. 2016. *Idiomarina tyrosinivorans* sp. nov., isolated from estuarine surface water. *Intl J Syst Evol Microbiol* 66: 5384-5391.
- Hamza F, Kumar AR, Zinjarde S. 2018. Efficacy of cell free supernatant from *Bacillus licheniformis* in protecting *Artemia salina* against *Vibrio alginolyticus* and *Pseudomonas gessardii*. *Microb Pathogen* 116: 335-344.
- Hao K, Liu JY, Ling F, Liu XL, Lu L, Xia L, Wang GX. 2014. Effects of dietary administration of *Shewanella haliotis* D4, *Bacillus cereus* D7 and *Aeromonas bivalvium* D15, single or combined, on the growth, innate immunity and disease resistance of shrimp, *Litopenaeus vannamei*. *Aquaculture* 428: 141-149.
- Hapsari F. 2016. The effect of fermented and non-fermented biofloc inoculated with bacterium *Bacillus cereus* for catfish (*Clarias gariepinus*) juveniles. *Aquacult Aquar Conserv Legis* 9: 334-339.
- Harikrishnan R, Kim JS, Balasundaram C, Heo MS. 2012. Immunomodulatory effects of chitin and chitosan enriched diets in *Epinephelus bruneus* against *Vibrio alginolyticus* infection. *Aquaculture* 326: 46-52.
- Huang L, Zuo Y, Jiang Q, Su Y, Qin Y, Xu X, et al. 2019. A metabolomic investigation into the temperature-dependent virulence of *Pseudomonas plecoglossicida* from large yellow croaker (*Pseudosciaena crocea*). *J Fish Dis* 42: 431-446.
- Ina SMY, Alsaari N, Mohamad A, Mursidi FA, MohdAris A, Amal MNA, et al. 2019. Vibriosis in fish: a review on disease development and prevention. *J Aquat Anim Health* 31: 3-22.
- Jiang Y, Zhou S, Chu W. 2019. The effects of dietary *Bacillus cereus* QSI-1 on skin mucus proteins profile and immune response in Crucian Carp (*Carassius auratus gibelio*). *Fish Shellfish Immunol* 89, 319-325.
- Karim M, Zhao W, Rowley D, Nelson D, Gomez CM. 2013. Probiotic strains for shellfish aquaculture: protection of eastern oyster, *Crassostrea virginica*, larvae and juveniles against bacterial challenge. *J Shellfish Res* 32: 401-408.
- Kaskatepe B, Yildiz S, Gumustas M, Ozkan S. 2017. Rhamnolipid production by *Pseudomonas putida* IBS036 and *Pseudomonas pachastrellae* LOS20 with using pulps. *Curr Pharm Anal* 13: 138-144.
- Kennedy J, Flemer B, Jackson SA, Morrissey JP, O'Gara F, Dobson AD. 2014. Evidence of a putative deep-sea specific microbiome in marine sponges. *PLoS One* 9: e91092. DOI: 10.1371/journal.pone.0091092.
- Khademzade O, Zakeri M, Haghi M, Mousavi SM. 2020. The effects of water additive *Bacillus cereus* and *Pediococcus acidilactici* on water quality, growth performances, economic benefits, immunohematology and bacterial flora of whiteleg shrimp *Penaeus vannamei* Boone, 1931) reared in earthen ponds. *Aquacult Res* 51: 1759-1770.
- Khalifa AY, Metwally A, Ammar RB, Farghaly FA. 2020. ACC Deaminase-Containing Rhizobacteria from Rhizosphere of *Zygophyllum coccineum* Alleviate Salt Stress Impact on Wheat (*Triticum aestivum* L.). *Sci J King Faisal Univ Basic Appl Sci* 21: 89-102.
- Kim M, Oh HS, Park SC, Chun J. 2014. Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. *Intl J Syst Evol Microbiol* 64: 346-351.
- Kitikiew S, Chen JC, Putra DF, Lin YC, Yeh ST, Liou CH. 2013. Fucoidan effectively provokes the innate immunity of white shrimp *Litopenaeus vannamei* and its resistance against experimental *Vibrio alginolyticus* infection. *Fish Shellfish Immunol* 34: 280-290.
- Korkea ATL, Heikkinen J, Thompson KD, Von Wright A, Austin B. 2011. *Pseudomonas* sp. M174 inhibits the fish pathogen *Flavobacterium psychrophilum*. *J Applied Microbiol* 111:266-277.
- Korkut GG, Söderhäll I, Söderhäll K, Noonin C. 2018. The effect of temperature on bacteria-host interactions in the freshwater crayfish, *Pacifastacus leniusculus*. *J Invert Pathol* 157: 67-73.
- Kuebutornye FK, Abarike ED, Lu Y. 2019. A review on the application of *Bacillus* as probiotics in aquaculture. *Fish Shellfish Immunol* 87: 820-828.
- Kvingedal R, Evans BS, Lind CE, Taylor JJ, Dupont NM, Jerry DR. 2010. Population and family growth response to different rearing location,

- heritability estimates and genotype×environment interaction in the silver-lip pearl oyster (*Pinctada maxima*). *Aquaculture* 304: 1-6.
- Lamendella R, Strutt S, Borglin S, Chakraborty R, Tas N, Mason OU, Hultman J, Prestat E, Hazen TC, Jansson JK. 2014. Assessment of the Deepwater Horizon oil spill impact on Gulf Coast microbial communities. *Front Microbiol* 5: 130.
- Lee JC, Kim YS, Yun BS, Whang KS. 2015. *Idiomarina halophila* sp. nov., isolated from a solar saltern sediment. *Intl J Syst Evol Microbiol* 65: 1268-1273.
- León MJ, Martínez-Checa F, Ventosa A, Sánchez-Porro C. 2015. *Idiomarina aquatica* sp. nov., a moderately halophilic bacterium isolated from salters. *Intl J Syst Evol Microbiol* 65: 4595-4600.
- Li J, Wang LH, Xiang FG, Ding WL, Xi LJ, Wang MQ, et al. 2020. *Pseudomonas phragmitis* sp. nov., isolated from petroleum polluted river sediment. *Intl J Syst Evol Microbiol* 70: 364-372.
- Liu Y, Lai Q, Shao Z. 2018. Genome-based analysis reveals the taxonomy and diversity of the family Idiomarinaeae. *Front Microbiol* 9: 2453. DOI: 10.3389/fmicb.2018.02453.
- López JR, Diéguez AL, Doce A, De la Roca E, De la Herran R, Navas JJ, et al. 2012. *Pseudomonas baetica* sp. nov., a fish pathogen isolated from wedge sole, *Dicologlossa cuneata* (Moreau). *Intl J Syst Evol Microbiol* 62: 874-882.
- Macias AOL, Ramirez OJJ, Cordova CAI, Saucedo PE. 2010. Evaluation of natural and commercial probiotics for improving growth and survival of the pearl oyster, *Pinctada mazatlanica*, during late hatchery and early field culturing. *J World Aquac Soc* 41: 447-454.
- Manez KS. 2010. Java's forgotten pearls: the history and disappearance of pearl fishing in the Segara Anakan lagoon, South Java, Indonesia. *J Histor Geogr* 36: 367-376.
- Mao Z, Qiu Y, Zheng L, Chen J, Yang J. 2012. Development of a visual loop-mediated isothermal amplification method for rapid detection of the bacterial pathogen *Pseudomonas putida* of the large yellow croaker (*Pseudosciaena crocea*). *J Microbiol Methods* 89: 179-184.
- Michaud L, Lo Giudice A, Troussellier M, Smedile F, Bruni V, Blancheton JP. 2009. Phylogenetic characterization of the heterotrophic bacterial communities inhabiting a marine recirculating aquaculture system. *J Appl Microbiol* 107: 1935-1946.
- Nakai T. 2010. Application of bacteriophages for control of infectious diseases in aquaculture. In: Parviz M, Sabour MW (eds). *Bacteriophages in the Control of Food-and Waterborne Pathogens*. American Society of Microbiology, Washington, DC.
- Napitupulu HG, Rumengan IF, Wullur S, Ginting EL, Rimper JR, Toloh BH. 2019. *Bacillus* sp. as a decomposition agent in the culture of *Brachionus rotundiformis* using raw fish as a source of nutrition. *Jurnal Ilmiah Platax* 7 (1): 158-169. [Indonesian].
- Nour E, ElGhiet ENA. 2011. Efficacy of *pseudomonas fluorescens* as biological control agent against *Aeromonas hydrophila* infection in *Oreo chromis niloticus*. *World J Fish Marine Sci* 3: 564-569.
- Ogello EO, Wullur S, Hagiwara A. 2019. Blending fishwastes and chicken manure extract as low-cost and stable diet for mass culture of freshwater zooplankton, optimized for aquaculture. *IOP Conf Ser Mater Sci Eng* 567 (1): 012022. DOI: 10.1088/1757-899X/567/1/012022.
- Ogello EO, Wullur S, Sakakura Y, Hagiwara A. 2018. Composting fishwastes as low-cost and stable diet for culturing *Brachionus rotundiformis* Tschugunoff (Rotifera): Influence on water quality and microbiota. *Aquaculture* 486: 232-239.
- Ogello EO, Wullur S, Yoshitaka S, Hagiwara A. 2020. Dietary value of waste-fed rotifer *Brachionus rotundiformis* on the larval rearing of Japanese whiting *Sillago japonica*. E3S Web of Conferences. DOI: 10.1051/e3sconf/202014701005.
- Oh EG, Son KT, Yu H, Lee TS, Lee HJ, Shin S, et al. 2011. Antimicrobial resistance of *Vibrio parahaemolyticus* and *Vibrio alginolyticus* strains isolated from farmed fish in Korea from 2005 through 2007. *J Food Prot* 74: 380-386.
- Orsod M, Joseph M, Huyop F. 2012. Characterization of exopolysaccharides produced by *Bacillus cereus* and *Brachybacterium* sp. isolated from Asian sea bass (*Lates calcarifer*). *Mal J Microbiol* 8: 170-174.
- Özen AI, Ussery DW. 2012. Defining the *Pseudomonas* genus: where do we draw the line with *Azotobacter*?. *Microbial Ecol* 63: 239-248.
- Preetha R, Vijayan KK, Jayaprakash NS, Alavandi SV, Santiago TC, Singh IB. 2015. Optimization of culture conditions for mass production of the probiotics *Pseudomonas* MCCB 102 and 103 antagonistic to pathogenic *Vibrios* in aquaculture. *Probiotics Antimicrob Proteins* 7: 137-145.
- Priyaja P. 2012. Pyocyanin (5-methyl-1-hydroxyphenazine) produced by *Pseudomonas aeruginosa* as antagonist to *vibrios* in aquaculture: over expression, downstream process and toxicity. [Dissertation]. Cochin University of Science and Technology, India.
- Richards GP, Watson MA, Needleman DS, Church KM, Häse CC. 2015. Mortalities of Eastern and Pacific oyster larvae caused by the pathogens *Vibrio coralliilyticus* and *Vibrio tubiashii*. *Appl Environ Microbiol* 81: 292-297.
- Riddle MR, Baxter BK, Avery BJ. 2013. Molecular identification of microorganisms associated with the brine shrimp *Artemia franciscana*. *Aquat Biosyst* 9: 7. DOI: 10.1186/2046-9063-9-7.
- Sánchez PT, Zarzuela RI, de Blas I, Balcázar JL. 2014. Probiotics in aquaculture: a current assessment. *Rev Aquacult* 6: 133-146.
- Sandhya SV, Preetha K, Vijayan KK. 2017. Phylogenetic diversity of culturable bacteria in *Chaetoceros gracilis* mass culture system of a marine finfish hatchery. *J Mar Biol Assoc India* 59: 12-18.
- Saulnier D, De Decker S, Haffner P, Cobret L, Robert M, Garcia C. 2010. A large-scale epidemiological study to identify bacteria pathogenic to Pacific oyster *Crassostrea gigas* and correlation between virulence and metalloprotease-like activity. *Microb Ecol* 59: 787-798.
- Shahi N, Mallik SK. 2014. Recovery of *Pseudomonas koreensis* from eye lesions in golden mahseer, *Tor putitora* (Hamilton, 1822) in Uttarakhand, India. *J Fish Dis* 37: 497-500.
- Sitdhipol J, Visessanguan W, Benjakul S, Yukphan P, Tanasupawat S. 2013. *Idiomarina piscisalsi* sp. nov., from fermented fish (pla-ra) in Thailand. *J Gen Appl Microbiol* 59: 385-391.
- Slifka KJ, Newton AE, Mahon BE. 2017. *Vibrio alginolyticus* infections in the USA, 1988–2012. *Epidemiol & Infection*, 145: 1491-1499.
- Somalkar VV, Mawlankar R, Ramana VV, Joseph N, Shouche YS, Dastager SG. 2015. *Bacillus filamentosus* sp. nov., isolated from sediment sample. *Antonie van Leeuwenhoek* 107: 433-441.
- Southgate PC, Beer AC, Ngalaufe P. 2016. Hatchery culture of the winged pearl oyster, *Pteria penguin*, without living microalgae. *Aquaculture* 451: 121-124.
- Souza DMD, Martins GB, Piedras SRN, Pouey JLOF, Robaldo RB, Leite FPL. 2012. Probiotic actions of *Bacillus cereus* var. *toyoi* and *Saccharomyces boulardii* in silver catfish (*Rhamdia quelen*) larvae culture. *Revista Brasileira de Zootecnia* 41 (3): 815-819.
- Tanyaros S, Sujarit C, Jansri N, Tarangkoon W. 2016. Baker's yeast as a substitute for microalgae in the hatchery rearing of larval and juvenile tropical oyster (*Crassostrea belcheri*, Sowerby 1871). *J Appl Aquacult* 28: 35-46.
- Teige F, Pepler C, Steinhagen D, Jung SV. 2020. Effect of disinfection with peracetic acid on the microbial community of a seawater aquaculture recirculation system for Pacific white shrimp (*Litopenaeus vannamei*). *J Fish Dis* 43: 991-1017.
- Umayaparvathi S, Meenakshi S, Arumugam M, Balasubramanian T. 2013. Purification and characterization of protease from *Bacillus cereus* SU12 isolated from oyster *Saccostrea cucullata*. *Afr J Biotechnol* 12:40.
- Vinoj G, Vaseeharan B, DavidJayaseelan B, Rajakumaran P, Ravi C. 2013. Inhibitory effects of *Bacillus licheniformis* (DAB1) and *Pseudomonas aeruginosa* (DAP1) against *Vibrio parahaemolyticus* isolated from *Fenneropenaeus indicus*. *Aquacult Intl* 21: 1121-1135.
- Wang JW, Cai M, Nie Y, Hu B, Yang Y, Wu XL. 2020. *Pseudomonas jilinenis* sp. nov., isolated from oil production water of Jilin oilfield in China. *Curr Microbiol*. DOI: 10.1007/s00284-019-01798-2.
- Wang MQ, Sun L. 2016. *Pseudomonas oceani* sp. nov., isolated from deep seawater. *Intl J Syst Evol Microbiol* 66: 4250-4255.
- Wang Y, Fu D, Luo P, He X. 2012. Identification of the immune expressed sequence tags of pearl oyster (*Pinctada martensii*, Dunker 1850) responding to *Vibrio alginolyticus* challenge by suppression subtractive hybridization. *Compar Biochem Physiol Part D: Genom Proteom* 7: 243-247.
- Wang YD, Huang SJ, Chou HN, Liao WL, Gong HY, Chen JY. 2014. Transcriptome analysis of the effect of *Vibrio alginolyticus* infection on the innate immunity-related complement pathway in *Epinephelus coioides*. *BMC Genomics* 15: 1102. DOI: 10.1186/1471-2164-15-1102.
- Wang Z, Wang B, Chen G, Jian J, Lu Y, Xu Y, Wu Z. 2016. Transcriptome analysis of the pearl oyster (*Pinctada fucata*) hemocytes in response to *Vibrio alginolyticus* infection. *Gene* 575: 421-428.
- Was 6 g M, Southgate PC. 2016. The effects of stocking density and ration on survival and growth of winged pearl oyster (*Pteria penguin*)

- larvae fed commercially available microalgae concentrates. *Aquacult Rep* 4: 17-21.
- Wu CJ, Wang H, Chan YL, Li TL. 2011. Passive immune-protection of small abalone against *Vibrio alginolyticus* infection by anti-Vibrio IgY-encapsulated feed. *Fish Shellfish Immunol* 30: 1042-1048.
- Wu HJ, Sun LB, Li CB, Li ZZ, Zhang Z, Wen XB, et al. 2014. Enhancement of the immune response and protection against *Vibrio parahaemolyticus* by indigenous probiotic *Bacillus strains* in mud crab (*Scylla paramamosain*). *Fish Shellfish Immunol* 41: 156-162.
- Wullur S, Ginting EL, Waraow V, Rumengan IFM, Ogello EO, Hagiwara A. 2019. Growth response of rotifers on a bacterial-based diet made from fishwastes. *IOP Conf Ser: Mater Sci Eng* 567: 012030. DOI: 10.1088/1757-899X/567/1/012030.
- Wullur S, Napitupulu H, Wantania LL, Ginting EL, Warouw V, Tilaar S, Rumengan IFM. 2020. Molecular identification of bacteria isolated from culture medium of rotifer fed on fishery waste diet. *Biodiversitas* 21: 2735-2740.
- Xu J, Zeng X, Jiang N, Zhou Y, Zeng L. 2015. *Pseudomonas alcaligenes* infection and mortality in cultured Chinese sturgeon, *Acipenser sinensis*. *Aquaculture* 446: 37-41.
- Yang G, Cao H, Jiang W, Hu B, Jian S, Wen C, et al. 2019. Dietary supplementation of *Bacillus cereus* as probiotics in Pengze crucian carp (*Carassius auratus* var. *Pengze*): Effects on growth performance, fillet quality, serum biochemical parameters and intestinal histology. *Aquacult Res* 50: 2207-2217.
- Yukgehnai K, Kumar P, Sivachandran P, Marimuthu K, Arshad A, Paray BA, Arockiaraj J. 2020. Gut microbiota metagenomics in aquaculture: factors influencing gut microbiome and its physiological role in fish. *Rev Aquacult*. DOI: 10.1111/raq.12416.
- Zhang YJ, Zhang XY, Zhao HL, Zhou MY, Li HJ, Gao ZM, et al. 2012. *Idiomarina maris* sp. nov., a marine bacterium isolated from sediment. *Intl J Syst Evol Microbiol* 62: 370-375.
- Zhao Y, Yuan L, Wan J, Sun Z, Wang Y, Sun H. 2016. Effects of potential probiotic *Bacillus cereus* EN25 on growth, immunity and disease resistance of juvenile sea cucumber *Apostichopus japonicus*. *Fish Shellfish Immunol* 49: 237-242.
- Zhong ZP, Liu Y, Liu HC, Wang F, Song L. et al. 2014. *Idiomarina planktonica* sp. nov., isolated from a saline lake. *Intl J Syst Evol Microbiol* 64: 3411-3416.
- Zhu F, Wang Z, Sun BZ. 2016. Differential expression of microRNAs in shrimp *Marsupenaeus japonicus* in response to *Vibrio alginolyticus* infection. *Dev Comp Immunol* 55: 76-79.



# Molecular identification of bacteria isolated from culture medium of the gold-lipped pearl oyster *Pinctada maxima* larvae

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://biodiversitas.mipa.uns.ac.id">biodiversitas.mipa.uns.ac.id</a> Internet Source	2%
2	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	1%
3	Elvy Like Ginti, Gladys G. Poluan, Letha L. Wantani, Emma Mauren Mok, Veibe Warouw, Mayse S. Siby, Stenly Wullur. "Screening and Identification of Sponge-Associated Chitinolytic Bacteria by Forming Chitosan from Manado Bay, Indonesia", <i>Pakistan Journal of Biological Sciences</i> , 2021 Publication	1%
4	"The Prokaryotes", Springer Science and Business Media LLC, 2014 Publication	1%
5	<a href="http://othes.univie.ac.at">othes.univie.ac.at</a> Internet Source	1%
6	Qiang Chen, Xiamin Jiang, Qingxi Han, Peng Sheng, Yonghui Chai, Ruibing Peng, Maowang	1%

Jiang, Zhongjian Mao, Shengtao Wu. "Growth, calcium content, proximate composition, and fatty acid composition of triangle sail mussel (*Hyriopsis cumingii*) fed five different microalgal diets", *Aquaculture*, 2020

Publication

---

7	<a href="http://koreascience.or.kr">koreascience.or.kr</a> Internet Source	1%
8	<a href="http://www.coursehero.com">www.coursehero.com</a> Internet Source	1%
9	<a href="http://archive.org">archive.org</a> Internet Source	1%
10	<a href="http://www.frontiersin.org">www.frontiersin.org</a> Internet Source	<1%
11	<a href="http://raw.githubusercontent.com">raw.githubusercontent.com</a> Internet Source	<1%
12	<a href="http://moam.info">moam.info</a> Internet Source	<1%
13	<a href="#">Pepen Supendi, Sri Widiyantoro, Abdul Muhari, Nicholas Rawlinson et al. "Potential megathrust earthquakes and tsunamis off the southern coast of West Java, Indonesia", <i>Research Square</i>, 2020</a> Publication	<1%
14	<a href="http://www.yumpu.com">www.yumpu.com</a> Internet Source	<1%

---

---

15	<a href="#">Pseudomonas Molecular and Applied Biology, 2016.</a> Publication	<1%
16	<a href="#">academicjournals.org</a> Internet Source	<1%
17	<a href="#">Submitted to ICAR - Central Institute of Fisheries Education</a> Student Paper	<1%
18	<a href="#">Brett A. Froelich, Dayle A. Daines. " In hot water: effects of climate change on –human interactions ", Environmental Microbiology, 2020</a> Publication	<1%
19	<a href="#">Haoda Ye, Zhijin Xu, Zhen Tao, Weiye Li, Yanyan Li, Amin Yang, Wei Wang, Xiaolong Yin, Xiaojun Yan. "Efficacy and safety of Pseudomonas plecoglossicida mutant ΔtssD-1 as a live attenuated vaccine for the large yellow croaker (Larimichthys crocea)", Aquaculture, 2021</a> Publication	<1%
20	<a href="#">journals.plos.org</a> Internet Source	<1%
21	<a href="#">theses.gla.ac.uk</a> Internet Source	<1%
22	<a href="#">fishexchange.mpeda.gov.in</a> Internet Source	<1%

---

---

23 [fas.biomedcentral.com](https://fas.biomedcentral.com) Internet Source <1%

---

24 [eprints.cmfri.org.in](https://eprints.cmfri.org.in) Internet Source <1%

---

25 [ueaeprints.uea.ac.uk](https://ueaeprints.uea.ac.uk) Internet Source <1%

---

26 [orbi.ulg.ac.be](https://orbi.ulg.ac.be) Internet Source <1%

---

27 Hai, Ngo Van. "The use of probiotics in aquaculture", *Journal of Applied Microbiology*, 2015. Publication <1%

---

28 [digitalcommons.uri.edu](https://digitalcommons.uri.edu) Internet Source <1%

---

29 Zhenghua Deng, Haijun Wei, Wang Zhao, Mingqiang Chen, Gang Yu, Jing Sun, Youning Li, Yu Wang. "Embryonic development and larval cultivation of *Paphia schnelliana* (Dunker), a unique economic species of the Beibu Gulf", *Aquaculture*, 2021 Publication <1%

---

30 Kathleen Schwerdtner Máñez. "Java's forgotten pearls: the history and disappearance of pearl fishing in the Segara Anakan lagoon, South Java, Indonesia", *Journal of Historical* <1%

# Geography, 2010

Publication

31

[amsdottorato.unibo.it](http://amsdottorato.unibo.it)

Internet Source

<1%

32

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

<1%

33

Gouranga Biswas, Masahiro Sakai. "Loop-mediated isothermal amplification (LAMP) assays for detection and identification of aquaculture pathogens: current state and perspectives", *Applied Microbiology and Biotechnology*, 2014

Publication

<1%

34

"*Marine Microbiology*", Wiley, 2013

Publication

<1%

35

Hongkuan Zhang, Karsoon Tan, Shengkang Li, Hongyu Ma, Huaiping Zheng. "Genome-wide analysis of TRAF gene family and its response to bacterial infection in noble scallop *Chlamys nobilis* with different carotenoids content", *Aquaculture*, 2020

Publication

<1%

36

Ben Yang, Shangyu Zhai, Xin Li, Jing Tian, Qi Li, Hongwei Shan, Shikai Liu. "Identification of *Vibrio alginolyticus* as a causative pathogen associated with mass summer mortality of the

<1%

# Pacific Oyster (*Crassostrea gigas*) in China", Aquaculture, 2021

Publication

---

37

Extremophiles Handbook, 2011.

Publication

---

<1%

38

Zhijuan Mao, Yangyu Qiu, Lei Zheng, Jigang Chen, Jifang Yang. "Development of a visual loop-mediated isothermal amplification method for rapid detection of the bacterial pathogen *Pseudomonas putida* of the large yellow croaker (*Pseudosciaena crocea*)", Journal of Microbiological Methods, 2012

Publication

---

<1%

39

Brian Austin, Dawn A. Austin. "Bacterial Fish Pathogens", Springer Science and Business Media LLC, 2012

Publication

---

<1%

40

[library.wur.nl](http://library.wur.nl)

Internet Source

---

<1%

41

Shurong Xiang, Tandong Yao, Lizhe An, Zhen Li, Guangjian Wu, Youqing Wang, Baiqing Xu, Junxia Wang. "Change of bacterial community in the Malan ice core and its relation to climate and environment", Chinese Science Bulletin, 2004

Publication

---

<1%

"Cellular Ecophysiology of Microbe:

- 
- 42 "Hydrocarbon and Lipid Interactions", Springer Science and Business Media LLC, 2018 <1%
- Publication
- 
- 43 Southgate, Paul C., Andrew C. Beer, and Poasi Ngaluafe. "Hatchery culture of the winged pearl oyster, Pteria penguin, without living micro-algae", Aquaculture, 2016. <1%
- Publication
- 
- 44 Junjing Xue, Kaikai Shen, Yi Hu, Yajun Hu, Vikas Kumar, Gang Yang, Chungeng Wen. "Effects of dietary Bacillus cereus, B. subtilis, Paracoccus marcusii, and Lactobacillus plantarum supplementation on the growth, immune response, antioxidant capacity, and intestinal health of juvenile grass carp (Ctenopharyngodon idellus)", Aquaculture Reports, 2020 <1%
- Publication
- 
- 45 "Probiotics and Prebiotics in Animal Health and Food Safety", Springer Science and Business Media LLC, 2018 <1%
- Publication
- 
- 46 Greeshma James, Bini C. Das, Seena Jose, Rejish Kumar V.J. "Bacillus as an aquaculture friendly microbe", Aquaculture International, 2021 <1%
- Publication
-

47

Sathyajith Chandhini, Vattiringal Jayadradhan  
Rejish Kumar. "Transcriptomics in aquaculture:  
current status and applications", Reviews in  
Aquaculture, 2018

Publication

---

<1%

48

"Microbial Communities in Aquaculture  
Ecosystems", Springer Science and Business  
Media LLC, 2019

Publication

---

<1%

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On