

Re: revised paper
Yahoo/Inbox

gavriloaie ionel claudiu <ionelclaudiu@yahoo.com>

To: Yahoo

Tue, Mar 31, 2020 at 12:14 AM

Dear Professor Pratasik,

Regarding the comments on the paper of professor Lalamentik, both reviewers and myself had to search on the Internet for the validity of coral Latin names. We used marinespecies.org this time. For example, as response to our comments, professor Lalamentik said that Dipsastraea is probably not a coral genus (we said that Favia pallida is actually classified as Dipsastraea pallida - see this: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=758233>, and this: Hoeksema, B. W.; Cairns, S. (2020). World List of Scleractinia. *Dipsastraea pallida* (Dana, 1846). Accessed through: World Register of Marine Species at: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=758233> on 2020-03-30.). So, we have no doubt that professor Lalamentik is a really great scientist and an expert in the coral species. However, we cannot know 100% a scientist is aware about the latest changes in the taxonomy. The books of dr. Veron were published in 2000. How many changes occurred in this field in the last 20 years?

And thus we are at the same point: how much can we trust the databasis?

For example, for fish species I know many Romanian and foreigner scientist and I often ask them via the phone or e-mail about one aspect or another regarding the taxonomy and systematics. But I do know anybody in Romania who knows much about the corals, since we only have few species in the Black Sea.

So, this is a day by day challenge for us here on Bioflux. All the advices are much appreciated.

I apologise for sending you such extended message. Thank you very much for your time and consideration.

Cordially yours,

Claudiu G.

On Saturday, March 28, 2020, 4:22:25 AM GMT+2, Yahoo <spjong07@yahoo.com> wrote:


Dear Dr. Gavrilă,

I herewith send you the revised paper of Lalamentik et al and Rondonuwu et al (enclosed). Those articles have been corrected and edited following the reviewers' comments. I also provide some arguments in Lalamentik et al's paper to clarify the issue after discussing it with the corresponding author, particularly concerning the coral species identification. As far as I know, Lalamentik is one of the best coral taxonomists in Indonesia who focuses on morphological characteristics, and therefore, he has strongly followed Veron's coral description. All corrections are given in yellow. Also, the payment for Rondonuwu et al's publication fee will be done on next Monday. Thank you

Sincerely Yours,

Silvester B. Pratasik

Respons terhadap revisi 1.



Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate

Laurentius T. X. Lalamentik, Rene C. Kepel, Lawrence J. L. Lumingas, Unstain N. W. J. Rembet, Silvester B. Pratasik, Desy M. H. Mantiri

Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia. Corresponding author: L. T. X. Lalamentik, ottylalamentik@yahoo.co.id

Abstract. A study on colonization of Faviidae corals on the agricultural waste-materialized artificial substrate was conducted in Selat Besar, Ratatotok district, southeast Minahasa regency, North Sulawesi. Nine artificial substrates modules made of mixture of cement, sand, padi husk, and bamboo were placed for about 5 years on the sea bottom of Selat Besar waters. All corals of family Faviidae found on the artificial substrate were collected. Results showed that Faviidae corals could live and develop on those substrates. Fifteen species of 5 genera of Faviidae, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennettiae*, *Echinopora gemmacea*, *E. lamellosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. sinensis* were recorded in the present study. Mean number of colonies of Faviidae corals was 3 col mod⁻¹, while mean diameter of the corals attached on the artificial substrate was 7.45 cm long. The distribution pattern of Faviidae corals was clumped. The diversity of Faviidae corals on the artificial substrate was low ($H' = 0.677 < 1$). The dominance index showed no dominant species ($D = 0.23$). In addition, the artificial substrate module in this study could become an alternative technique to rehabilitate the degraded coral reefs.

Key Words: diversity, dominance, distribution pattern, Selat Besar.

Introduction. Corals are animals belonging to phylum Cnidaria (Reid et al 2009), while coral reef is an ecosystem built by calcium carbonate-producing marine biota, especially coral animals, together with other biota living on the sea bottom or water column, such as mollusks, crustaceans, echinoderms, porifera, tunicates, and other biota free-living in surrounding waters including plankton and fishes (Giyanto et al 2017; Lalamentik 1995). According to Barus (2013), coral reef is one of the unique communities that are entirely formed from biological activities. It is one of the main coastal and marine ecosystems that possess the highest productivity and biodiversity so that it is often called as tropical marine forest.

Coral Triangle (CTI) is known as center of world marine biodiversity with the highest coral diversity, 76% of total coral species. The region inside the ecological boundary of CTI covers nearly 73,000 km² of coral reefs (29% of total world coral reef area) distributed in six countries, Indonesia, Malaysia, Philippines, Timor Leste, Papua New Guinea and Solomon Islands (Burke et al 2012). As part of the CTI, Indonesia is an archipelagic country with 16,056 islands, 1,922,570 km² terrestrial area and 3,257,483 km² water in the west Indo-Pacific (BIG 2016). Indonesia that is located in the CTI area is recognized as one of the countries with high biodiversity consisting of more than 80 coral genera and 596 species, particularly in North Sulawesi waters where have been recorded more than 80 genera (Suharsono 2008).

Coral taxonomic and distribution studies in North Sulawesi were carried out by several scientists (Lalamentik 1998; Halidu et al 2016; Nasaru et al 2017; Suleman et al 2017; Lalamentik & Rembet 2018). So far, there is no study on corals of family Faviidae living and growing on the artificial substrate, especially in North Sulawesi waters. Hence, the present study focuses on Faviidae coral colonies on the artificial substrate with the

Commented [indra1]: see the comment on Table 1

Commented [indra2]: same as above

Commented [indra3]: it is missing from the list

Commented [indra4]: to be added to the list

Commented [indra5]: add to list

Commented [indra6]: add to list

objectives of identifying the coral species, analysing the density and the diameter of the coral colony, the ecological index, and assessing the distribution patterns. This finding is expected to be one of the references in coral development studies, concerning Faviidae coral colonization on the artificial substrate. Besides, it could provide alternative artificial substrates in coral reef rehabilitation program.

Material and Method

Study site. This work was accomplished in Selat Besar waters, Ratatotok district, southeast Minahasa regency (Figure 1), North Sulawesi. Selat Besar is flanked by Totok bay and Moluccas Sea. The study site was laid on 0°51'34.945"S and 124°44'14.416" E (Figure 1) using Global Position System (GPS).




Figure 1. Sampling locality.

Artificial substrate construction. In this study, 9 artificial substrate modules were set on the sea bottom, approximately 7-8 m depth. Sheehy (1986) stated that one of the major roles of artificial substrate placement in the water is to create new habitats similar to artificial reefs to be inhabited by target species. The use of artificial substrate as coral new habitat had been conducted in Singaporean waters in order to restore the coral reefs (Ng & Chou 2017). Each modul consists of 10 concrete blocks made of mixture of padi husk, sand, and cement. The block frames were made of bamboo. The use of bamboo in concrete construction has been done before as mentioned in Glenn (1950) and Mehra & Ghosh (1965). One of the properties that would make bamboo a good substitute to steel in reinforced concrete is its strength (Varma 2017). The block had a dimension of 100 cm length, 15 cm width, and 15 cm height. The blocks were set in such a manner to yield a module. Each module is estimated to have an area of 4.5 m². These 9 modules were placed since June 2014. Sample collections were carried out in July 2019 (after 5 years of

Commented [indra7]: it would be great to add some photo or drawing depicting one modul

Commented [indra8]: may be natural?

Commented [indra9]: add to list

deployment). The period of 5 years is determined because the massive coral growth is slow. This duration is considered enough to see whether there is Faviidae coral capable of adhering and growing on the artificial substrate.

Sample collection and handling procedures. All massive coral species found to attach on the module substrate were taken using a chisel and hammer. The samples were then put into labelled bag as the module number. All corals were collected by using SCUBA equipments. All activities were documented by using an underwater camera.

Before species identification, the coral specimens were soaked in bleach-containing freshwater for 24 h in order to kill the coral and to prevent the presence of fungi. The corals were then cleaned under the running water to remove the attached dirt so that the clean coral and the bleaching coral skeleton were obtained. The corals were placed under direct sunlight to get dry to ease the identification.

Species identification. Coral identification was done following Veron (1986, 2000), Suharsono (2008), and Veron et al (1977). The specimens of Faviidae corals were examined through morphological observations under the SZ51 Olympus microscope on the characteristics of paliform lobe, septa cleats, coralite type, number of first septa, second septa, and third septa. The measurements of width of paliform lobe, coralite size, and calix size used a multitoyo digital caliper.

Data analysis. Faviidae coral colonization data were directed to ecological index analyses as follows:

Dominance index. This index was employed to know the dominant species in data sampling boundary. The present study applied Simpson dominance index (Krebs 1989):

$$D = \frac{\sum_{i=1}^s [n_i]^2}{N} \dots\dots\dots (1)$$

Where D = dominance index, n_i = number of i individuals, and N = total number of individuals.

Dominance index ranges between 0 and 1, if D approaches to 0, it indicates no dominance and the community structure is in stable condition, and if D approaches to 1, it indicates the presence of species dominance.

Density. The population density was calculated using Krebs (1989) as follows

$$K = \frac{N_i}{A} \dots\dots\dots (2)$$

Where, K = density (ind module⁻¹), N_i = number of individuals, and A = number of modules where samples were collected.

Diversity. The diversity index of genus *Favia* Oken, 1815 was calculated using Shannon-Wiener equation (Krebs 1989):

$$H = -\sum_{i=1}^s \frac{n_i}{N} \left(\frac{n_i}{N} \right) \dots\dots\dots (3)$$

Where H = Shannon-Wiener diversity index, n_i = number of individuals i , N = total number of individuals, and s = number of genera.

Distribution pattern. The distribution pattern of Faviidae corals was estimated using Morisita index (Krebs 1989):

$$Id = \frac{\sum_{i=1}^q n_i (n_i - 1)}{N (N - 1)} \dots\dots\dots (4)$$

Commented [indra10]: ???

Where I_d = Morisita index, n_i = number individuals in each plot, n = total number of individuals in all plots, and N = number of plots, in which $I_d = 1$ is categorized as random distribution, $I_d > 1$ as clumped, and $I_d < 1$ as uniform.

Results and Discussion

Water conditions of Selat Besar. Water quality is crucial for colonization of macroinvertebrates (Orwa et al 2018). Water temperature is one of the limiting factors for aquatic organisms to live, one of which is coral. Field measurements found that water temperature in the study site ranged from 29 to 30°C. This range is still suitable for coral development. According to Kurniawan (2011), corals, in general, can live at the temperature above 18°C, while growth, they require an ideal water temperature, between 25 and 32°C. Most coral reefs exist where salinity is stable and where average salinity is that of normal seawater 34-36‰ (Sheppard et al 2017). Water salinity in the study site was in the range of 30-31‰. Guntur (2011) found that water salinity for ideal coral growth in Indonesia ranged from 29 to 36‰. Therefore, water salinity in Selat Besar highly supports the coral to grow well. Light intensity does not influence larval settlement, and their response to light intensity may be species-specific (Mundy & Babcock 1998).

Corals of family Faviidae recorded on the artificial substrate. Physical and biological factors are important for recruitment success (Doropoulos et al 2016). As many as 15 coral species of 5 genera were recorded in this study. *Favia* and *Favites* had the highest number of species, each with 4 species, followed by genus *Goniastrea* with 3 species, then *Echinopora* and *Platygyra* with 2 species each. Table 1 demonstrates the coral species of Faviidae encountered on the module substrate with mean number of colonies of 3.00 per module. Even though these numbers are less than those reported by Kilfoyle et al (2008), 13 col mod⁻¹, the colonies comprised several coral families. The coral genera of Faviidae inhabiting the artificial substrate apparently come the coral reef around Selat Besar waters. It is reasonable, since Faviidae is coral family generally found in Selat Besar waters (Lalamentik 1996, 1997, 1998).

Commented [indra11]: we see no "n" in the equation

Commented [indra12]:

Commented [indra13R12]: is this I_d actually?

Commented [indra14]: add to list

Commented [indra15]: this might change if you will take into account the green comment on table 1

Commented [indra16]: it is Table 1, not 02, since this is the first table in the paper

Commented [indra17]: it has no meaning

Table 1

Faviidae coral species encountered living on the artificial substrate modul

No	Species
1	<i>Echinopora gemmacea</i> (Lamarck, 1816)
2	<i>E. lamellosa</i> (Esper, 1795)
3	<i>Favia laxa</i> (Klunzinger, 1879)
4	<i>F. matthaii</i> (Vaughan, 1918)
5	<i>F. pallida</i> (Dana, 1846)
6	<i>F. stelligera</i> (Dana, 1846)
7	* <i>Favites bennettiae</i> Veron, Pichon & Wijsman-Best, 1977
8	<i>F. complanata</i> (Ehrenberg, 1834)
9	<i>F. pentagona</i> (Esper, 1795)
10	<i>F. russelli</i> (Wells, 1954)
11	<i>Goniastrea aspera</i> Verrill, 1866
12	<i>G. favulus</i> (Dana, 1846)
13	<i>G. pectinata</i> (Ehrenberg, 1834)
14	<i>Platygyra daedalea</i> (Elis & Solander, 1786)
15	<i>P. sinensis</i> (Milne Edwards & Haime, 1849)

* The first record for Central Indonesia.

Favites bennettiae recorded in the present study is the first record found in the coral reef of Central Indonesian waters. This species was previously mostly found in the northeast of Australian waters (Veron et al 1977). Moreover, genus *Favia* was found having the

Commented [indra18]: when searching for the validity of the scientific names we discovered that some of the species in *Favia* genus are actually in *Goniastrea* or *Dipsastrea*. Actually, there are mismatches for almost all the five genera in the table. So, check again all the names by using reliable and updated databases (you could use <http://www.marinespecies.org>).

After checking and updating the species and genera names, you need to make all the corresponding updates and corrections both in text and figures/tables

Commented [indra19]: we removed the parentheses since this is the correct form (<http://www.marinespecies.org/aphia.php?p=taxdetails&id=207437>). We did the necessary corrections for all the authors names

highest mean number of colonies, 1.11 col mod⁻¹, followed by *Goniastrea*, 0.67 col mod⁻¹, and the lowest was found in *Platygyra*, 0.22 col mod⁻¹ (Figure 2). Variations in number of Faviidae colonies per module are dependent upon the abundance of larvae that survive and settle on the substrate. Faviidae tends to have good adaptability to occupying the available artificial substrates. Bachtiar et al (2011) found that Faviidae is one of the coral families capable of surviving in the reef ball deployed on the sea bottom.




Figure 2. Mean no. colonies of Faviidae per modul.

Mean diameter of the coral colony on the artificial substrate was 7.45 cm with the largest recorded in *Favia laxa*, 10.5 cm and the smallest in *F. pallida*, 3 cm. In comparison among genera, mean diameter of Faviidae corals attached on the artificial substrate ranged from 5.88 to 8.50 cm. Genus *Platygyra* has the largest mean diameter, 8.50 cm, and *Echinopora* had the smallest one, 5.88 cm (Figure 3). Variations in colony size could result from different sequence of planular settlements. This condition, according to Bachtiar et al (2011), indicates that coral colonization on the artificial substrate sustainably occurs.




Figure 3. Mean diameter of Faviidae corals living on the artificial substrate.

Ecological index. The present study measured dominance index (D) and diversity index (H') of Faviidae corals thriving on the artificial substrate. The data are given in Table 2.

Commented [indra20]: when writing the numbers (both on the left side and on the top of the columns), follow the English language rules. So, instead of 1,20 you should have 1.20; 1,00 - 1.00; 0,80 - 0.80 and so on.

The comment goes for the figure 3 as well.

Then, *Echinopora* should be written as *Echinopora*

Commented [indra21]: exactly the same comments as for figure 2

Table 2

Ecological index of Faviidae corals

No	Ecological index	Value	Remarks
1	Dominance Index (D)	0.23	No dominance
2	Distribution pattern (Id)	1.79	Clumped
3	Diversity index (H')	3.64	high

The dominance index of Faviidae corals is 0.23 (Table 2). It means that no species outnumbers the others. Selat Besar waters is still capable of supporting the coral life, and there is no competition that could make certain species be dominant (Muqsit et al 2016). The diversity index is 3.64. This value is supported by low species dominance index, in which there were 15 Faviidae species found. These numbers are high enough due to small study area cover. This finding is in agreement with Rondo et al (2014) that high diversity will cause low species dominance.

Distribution pattern analysis found that Faviidae in the study site had clumped distribution ($Id = 1.79 > 1$). This pattern could result from the fact that there were four modules not occupied by Faviidae corals. In Indonesia, studies on coral colonization was also done by Razak (2008) using eco-reef module in Manado bay waters, North Sulawesi, and found that the artificial substrate used are, in fact, inhabited by hard corals. Hence, the artificial substrate could be recommended as an alternative in coral reef rehabilitation program.

Conclusions. Fifteen coral species were found settling on the agricultural waste-materialized artificial substrate, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennettiae*, *Echinopora gemmacea*, *E. lamellosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. sinensis*. They had mean $3.00 \text{ col mod}^{-1}$ with mean diameter of 7.45 cm. The ecological index also indicated that the available artificial substrate could support various coral species to grow. In this study, Faviidae corals growing on the available artificial substrate have reflected clumped distribution. Thus, this type of module could be used for coral reef rehabilitation.

References

- Bachtiar I., Prayogo W., Sukri A. 2011 Coral colonization and growth reef ball modules at the Benete Bay, Sumbawa Island, Indonesia. Manado. P. 31-40.
- Doropoulos C., Roff G., Bozec Y. M., Zupan M., Werminghausen J., Mumby P. J., 2016 Characterizing the ecological trade-offs throughout the early ontogeny of coral recruitment. Ecol. Monogr. 86: 20-44.
- Guntur. 2011 [Coral ecology in the artificial reef]. Ghalia Indonesia. Semarang. 139 pp. [in Indonesian]
- Halidu, A., Lalamentik L. T. X., Rembet U. N. W. J. 2016 [Hard coral distribution in the reef flat of the south coast of Putus-Putus island, east Ratatotok, Ratatotok district, southeast Minahasa]. Jurnal Ilmiah Platax 4(1):19-30. [in Indonesian]
- Kilfoyle K., Rangel M. A., Dodge R. E., Spieler R. E. 2008 Coral Reef Restoration: Standardized Module Intervention and monitoring program in Mexico, Preliminary Results. Poster presentation material, 11 Int Coral Reef Sym Florida, Session number XXIV, p. 541
- Krebs C. J. 1989 Ecological Methodology. Harper Collins Publishers. New York. 654 pp.
- Lalamentik L. Th. X. 1995 [Study on coral reef potential in Tombasian district, Minahasa, North Sulawesi] Fakultas Perikanan dan Ilmu Kelautan UNSRAT. Manado. 28 pp. [in Indonesian] Unpublished
- Lalamentik L. T. X. 1996 Coral Reef conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1996). 23 pp. Unpublished.

Commented [indra22]: edit the list following the journal's format

- Lalamentik L. T. X. 1997 Coral Reef Conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1997). 22 pp. Unpublished.
- Lalamentik L. Th. X. 1998 Coral reef Condition Arround the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study for May 1998 Data in Ratatotok and Adjacent Waters, The District of Minahasa North Sulawesi. Fakultas Perikanan dan Ilmu Kelautan USNRAT. Manado. 23 pp. Unpublished.
- Lalamentik, L. Th. X., Rembet, U. N. J. 2018 [Study on the Taxonomy of Genus Favia (Oken, 1815) at the Reef Flats of Kampung Ambong Village in Likupang Timur District, North Minahasa]. JURNAL ILMIAH PLATAX, 6(1), 188-193. [in Indonesian]
- Mehra S. R., Ghosh R. G. 1965 "Bamboo-reinforced soil-cement," Civil Engineering and Public Works Review, Vol. 60, no. 711, October; Vol. 60, no. 712. November 1965.
- Mundy C. N., Babcock R., 1998 Role of light intensity and spectral quality in coral settlement? J. Exp. Mar. Biol. Ecol. 223 (2): 235-255.
- Muqsit A., Purnama D., Ta'alidin Z. 2016 [The community structure of coral reef in Pulau Dua, Enggano district, North Bengkulu regency.] Jurnal Enggano Vol. 1, No. 1 (75-78). Ilmu Kelautan Fakultas Pertanian Universitas Bengkulu. Bengkulu. [in Indonesian]
- Nasaru J. H., Lalamentik L. Th. X., Rembet U. N. W. J. 2017 [Pocillopora verrucosa (Ellis and Solander, 1786) distribution in the reef flat of the south coast of Putus-Putus island, east Ratatotok village, southeast Minahasa regency]. Jurnal Ilmiah Platax 5(1): 96-103 pp. Universitas Sam Ratulangi Manado. [in Indonesian]
- Ng C. S. L., Chou L. M. 2017. Coral reef restoration in Singapore-past, present and future. Sustainability Matters: Environmental Management in the Anthropocene. Singapore: World Scientific, 3-23 pp.
- Orwa P. O., Omondi R., Chemoiwa E. J. 2018 Colonization patterns of benthic macroinvertebrates in fertilized and non-fertilized earthen fish ponds. Int J Aquac Fish Sci 4(3): 022-026.
- Razak T. 2008. The population of hard coral colonies growing on Ecoreef artificial modules on Manado Tua Island, Bunaken National Park, North Sulawesi, Indonesia. Oral presentation material, 11 Int Coral Reef Sym Floria, Session number XXIV, p. 223.
- Reid C., Marshall J., Logan D., Kleine D. 2009 [Coral reef and climate change]. CoralWatch, The University of Queensland. Australia. 256 pp. [in Indonesian]
- Ricardo G. F., Jones R. J., Nordborg M., Andrew P. Negri. 2017. Settlement patterns of the coral Acropora millepora on sediment-laden surfaces. Science of the Total Environment. 609:277-288.
- Rondo M., Tamanampo J. F. W. S. 2014 Principles of Aquatic Ecology. Fakultas Perikanan dan Ilmu Kelautan, Universitas Sam Ratulangi. Manado. 427 pp.
- Sheppard C., Davy S., Pilling G., Graham N. 2017 The biology of coral reefs. Oxford University Press. 365 pp.
- Sheehy D. J. 1986 New approaches in artificial reef design and applications. p. 256-263. In F. M. D. Trie (ed.) Artificial reefs: marine and freshwater applications. Lewis Publishers Inc., Michigan, USA.
- Suharsono. 2008 [Coral species in Indonesia]. Lembaga Ilmu Pengetahuan (LIPI) : COREMAP Program. Jakarta. 116 p. [in Indonesian]
- Suleman Y., Lalamentik L. Th. X., Rembet U. N. W. J. 2017 [Hard coral Favites abdita (Ellis and Solander, 1786) distribution in the reef flat of Malalayan 2, Malalayang district, Manado city]. Jurnal Ilmiah Platax 5(1): 104-111. Universitas Sam Ratulangi. Manado. [Indonesian] pp [Indonesian]
- Varma M. B. 2017 Properties of cement concrete reinforced with bamboo-strip-Mat. IOSR, Journal of Mechanical and Civil Engineering (IOSR-JMCE), eISSN, 2278-1684: 47-59.
- Veron J. E. N. 1986 Corals of Australia and the Indo-Pacific. Angus and Robertson Publishers. Australia. 644 pp.

Commented [indra23]: to be quoted in text or to be removed from the list

Veron J. E. N. 2000 Corals of the World Volume 1. Australian Institute of Marine Science and CRR Qld Pty Ltd. Australia 463 pp.
Veron J. E. N., Pichon M., Wijsman-Best M. 1977 Scleractinia of Eastern Australia. Canberra. Vol. 3 (Part 2). 233 pp.

Received: 18 February 2020. Accepted: 17 March 2020. Published online: xx March 2020.

Authors:

Laurentius T. X. Lalamentik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: ottylalamentik@yahoo.co.id

R. Ch. Kepel, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: renecharleskepel65@gmail.com

Lawrence J. L. Lumingas, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: ljllumingas@yahoo.com

Unstain N. W. J. Rembet, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: onlyrembet@unsrat.ac.id

Silvester B. Pratasik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: spjong07@yahoo.com

Desy M. H. Mantiri, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: dmh_mantiri@unsrat.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Lalamentik L. T. X., Kepel R. C., Lumingas L. J. L., Rembet U. N. W. J., Pratasik S. B., Mantiri D. M. H., 2020 Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate. AACL Bioflux 13(2):xxx-xxx.