

# Relationship of Heme Oxygenase-1 - HO-1 Level with Onset and Severity in Normotensive Pregnancy and Severe Preeclampsia

*by Billy Kepel 2*

---

**Submission date:** 24-Feb-2021 07:41AM (UTC+0700)

**Submission ID:** 1516585384

**File name:** d\_Severity\_in\_Normotensive\_Pregnancy\_and\_Severe\_Preeclampsia.pdf (415.7K)

**Word count:** 3585

**Character count:** 19151



Published by DiscoverSys

## Relationship of Heme Oxygenase-1 (HO-1) level with onset and severity in normotensive pregnancy and severe preeclampsia



CrossMark

John Johannes Wantania,<sup>1\*</sup> Christian Homenta,<sup>2</sup> Billy Johnson Kepel<sup>3</sup>

### ABSTRACT

**Background:** Preeclampsia still becomes a problem in pregnancies. Various evidences showed that heme oxygenase-1 (HO-1) is very important in pregnancy. This study aims to understand the relationship of heme oxygenase-1 level with onset and severity in normotensive pregnancy and severe preeclampsia.

**Methods:** This is a cross sectional analytic comparative study, the subjects consisted of 26 patients with normotensive pregnancies and 26 patients with severe preeclampsia. Blood samples from women with < 34 / ≥ 34 weeks' normotensive pregnancies and women with severe preeclampsia were taken. HO-1 ELISA kit used to quantitate heme oxygenase-1 level in samples.

**Results:** The level of heme oxygenase-1 in normotensive pregnant women < 34 weeks lower than severe preeclampsia pregnant

women < 34 weeks ( $3.28 \pm 0.46$  ng/mL vs  $7.20 \pm 0.64$  ng/mL,  $p=0.003$ , respectively). The median level of heme oxygenase-1 in normotensive pregnant women ≥ 34 weeks was 2.96 (2.41–4.39) ng/mL, while severe preeclampsia pregnant women ≥ 34 weeks was 3.52 (2.88–5.43) ng/mL, ( $p=0.040$ ). The median level of heme oxygenase-1 in normotensive pregnant women was 3.04 (2.41–4.39) ng/mL, while severe preeclampsia pregnant women was 3.68 (2.88–5.67) ng/mL, ( $p=0.001$ ).

**Conclusions:** There is correlation between the incidence of severe preeclampsia with heme oxygenase-1 level in < 34 and ≥ 34 weeks of pregnancy. There is a significant difference between the level of heme oxygenase-1 in pregnant women with severe preeclampsia and women with normotensive pregnancy.

**Key words:** heme oxygenase-1 (HO-1), normotensive pregnancy, severe preeclampsia

**Cite This Article:** Wantania, J., Homenta, C., Kepel, B. 2016. Relationship of Heme Oxygenase-1 (HO-1) Level with Onset and Severity in Normotensive Pregnancy and Severe Preeclampsia. *Bali Medical Journal* 5(1): 105-109. DOI: 10.15562/bmj.v5i1.267

<sup>1</sup>Maternal Fetal Medicine Consultant in Department of Obstetrics and Gynecology, Faculty of Medicine University Sam Ratulangi/Prof. Dr. R. D. Kandou General Hospital, Manado-Indonesia

<sup>2</sup>Obstetrician and Gynecologist in Department of Obstetrics and Gynecology, Faculty of Medicine University Sam Ratulangi/Prof. Dr. R. D. Kandou General Hospital, Manado-Indonesia

<sup>3</sup>Statistician Consultant in Department of Obstetrics and Gynecology, Faculty of Medicine University Sam Ratulangi/Prof. Dr. R. D. Kandou General Hospital, Manado-Indonesia

\*Correspondence to: John Johannes Wantania Maternal Fetal Medicine Consultant in Department of Obstetrics and Gynecology, Faculty of Medicine University Sam Ratulangi/Prof. Dr. R. D. Kandou General Hospital, Manado-Indonesia. john\_w\_md@yahoo.com

### INTRODUCTION

Preeclampsia is still an obstetric problem that cannot be solved completely. Preeclampsia occurs in approximately 1.8 to 16.7% of pregnancies, the figures varied between countries.<sup>1</sup> Based on a systematic review conducted by the WHO, 16% of maternal deaths in developed countries, including the United States, were caused by hypertension in pregnancy and its complication.

This figure surpassed other major causes including bleeding (13%), abortion (8%), and sepsis (2%).<sup>2</sup> In Indonesia, the average incidence of preeclampsia is 3-10%, with maternal mortality rate of approximately 4.91%, ranging from 8.739 to 170.725 deaths.<sup>3</sup>

Several factors have been associated with the incidence of preeclampsia. In general, these risk factors can be classified as maternal factors (e.g. age, parity, history of preeclampsia), medical risk factors (such as chronic hypertension, diabetes mellitus or renal disease) and factors of the placenta (e.g hyperplacentosis on twin pregnancy and gestational trophoblastic diseases). Nevertheless, these factors are almost entirely a predisposing factor whose mechanism is unclear.<sup>4</sup>

In addition to the dominance of several risk factors, a number of theories have been developed to explain the path mechanism of preeclampsia, including, among others, genetic, immunological, oxidative stress, inflammation, hypoxia, angiogenic and hormonal factors. Although most theories tend to agree that the initial problem is the disruption of trophoblastic invasion, to date there are no theories with a satisfactory answer, with each factor being related to the others in a very complicated manner.

In recent years, a variety of important evidence has occurred, showing that heme oxygenase-1 (HO-1) is very important in pregnancy. Therefore, we need to understand the mechanisms underlying the protective effect of heme oxygenase-1 (HO-1).

These mechanisms can vary depending on the reproductive phase in which heme oxygenase-1 (HO-1) is involved. The microenvironment in which heme oxygenase-1 (HO-1) performs its action also greatly affects the mechanisms involved. Research on the mechanisms underlying the emergence of the effects of heme oxygenase-1 (HO-1), which positively affects the process of pregnancy is the release

of CO by heme oxygenase-1 (HO-1). This process does not only occur in the reproductive process,<sup>5,6</sup> but also occurs in several inflammatory processes.<sup>7,8</sup>

## MATERIALS AND METHODS

This is a cross sectional analytic comparative study, conducted at the Department of Obstetrics and Gynecology, Faculty of Medicine, University of Sam Ratulangi/Prof. Dr. R. D. Kandou General Hospital Manado, from March 1 2015 to June 30 2015. Inclusion criteria includes: women with < 34 / ≥ 34 weeks' normotensive pregnancies and women with severe preeclampsia who had had filled informed consent and willing to participate in the study. Exclusion criteria include: pregnant women with diabetes mellitus, kidney disorders, heart conditions, chronic hypertension, premature rupture of membranes, women with clinical signs of infection, women with multiple pregnancy, in utero fetal death, and women who are not willing to participate in this study. HO-1 ELISA kit used to quantitate heme oxygenase-1 (HO-1) level in

samples. Non parametric Mann-Whitney test and T-test variances was used to analyze the relationship between the level of heme oxygenase-1 in normotensive pregnancy and in severe preeclampsia.

## RESULTS

This study was conducted at the Department of Obstetrics and Gynecology, Faculty of Medicine, University of Sam Ratulangi/Prof. Dr. R. D. Kandou General Hospital Manado, from March 1 2015 to June 30 2015. The subjects consisted of 26 patients with normotensive pregnancies and 26 patients with severe preeclampsia.

Based on the data presented in Table 1, most of the subjects are multiparous (65.38% in normotensive pregnancy and 69.23% in severe preeclampsia), with gestational age of ≥ 34 week (69.23% in normotensive pregnancy and 57.69% in severe preeclampsia), with senior high school education (88.46% in normotensive pregnancy and 57.69% in severe preeclampsia), and work as housewives (69.23% in normotensive pregnancy and 65.38% in severe preeclampsia).

Based on the data presented in Table 2, in 8 samples of normotensive pregnancy, the average level of heme oxygenase-1 (HO-1) is at 3.28 ng/mL (95% CI 2.89-3.66), with a standard deviation of 0.46 ng/mL. In 11 samples of severe preeclampsia, the average value of heme-oxygenase-1 (HO-1) level is 4.20 ng/mL (95% CI 3.77-4.63), with a standard deviation of 0.64 ng/mL. T-test variances by 2 means onset of heme oxygenase-1 (HO-1) level showed that p value: 0.003, meaning that there are significant differences of heme oxygenase-1 (HO-1) level between the normotensive group and the severe preeclampsia group in < 34 weeks of pregnancy. Based on the data presented in Table 3, in 18 samples of normotensive pregnancy, the median level of heme oxygenase-1 (HO-1) is at 2.96 (min-max 2.41-4.39) ng/mL. In 15 samples of severe preeclampsia, the median value of heme-oxygenase-1 (HO-1) level is 3.52 (min-max 2.88-5.43) ng/mL. Mann-Whitney test showed that p value: 0.040, meaning that there are significant differences of heme oxygenase-1 (HO-1) level between the normotensive group and the severe preeclampsia group in ≥ 34 weeks of pregnancy.

Based on the data presented in Figure 1, statistically significant correlation between the incidence of severe preeclampsia with heme oxygenase-1 (HO-1) level in < 34 and ≥ 34 weeks of pregnancy.

Based on the data presented in Table 4, in 26 samples of normotensive pregnancy, the median level of heme oxygenase-1 (HO-1) is at 3.04 (min-max 2.41-4.39) ng/mL. In 26 samples of severe preeclampsia, the median value of heme-oxygenase-1 (HO-1) level is 3.68 (min-max

**Table 1** Characteristics of subjects

Characteristics N%	Normotensive		Severe Preeclampsia	
	N	%	N	%
Parity				
Primiparous	9	34.61	8	30.76
Multiparous	17	65.38	18	69.23
Gestational Age				
< 34 weeks	8	30.76	11	42.30
≥ 34 weeks	18	69.23	15	57.69
Education				
Elementary school	1	3.84	4	15.38
Junior high school	2	7.69	3	11.53
Senior high school	23	88.46	15	57.69
University	0	0	4	15.38
Occupation				
Housewife	18	69.23	17	65.38
Student	3	11.53	1	3.84
Private company employee	5	19.23	4	15.38
Government employee	0	0	4	15.38

**Table 2** Descriptive analysis of the difference of heme oxygenase-1 (HO-1) level in < 34 weeks of pregnancy


Variables	Normotensive, mean ± SD	Severe preeclampsia, mean ± SD	p
heme oxygenase-1 (HO-1), ng/mL	3.28 ± 0.46	4.20 ± 0.64	0.003

**Table 3** Descriptive analysis of the difference of heme oxygenase-1 (HO-1) level in  $\geq 34$  weeks of pregnancy


Variables	Normotensive, median (min-max)	Severe preeclampsia, median (min-max)	p
heme oxygenase-1 (HO-1), ng/mL	2.96 (2.41-4.39)	3.52 (2.88-5.43)	0.040

**Table 4** Descriptive analysis of the difference of heme oxygenase-1 (HO-1) level in normotensive pregnancy and severe preeclampsia

Variables	Normotensive, median (min-max)	Severe preeclampsia, median (min-max)	p
heme oxygenase-1 (HO-1), ng/mL	3.04 (2.41-4.39)	3.68 (2.88-5.67)	0.001



**Figure 1** Box and whisker plots analysis of the difference of heme oxygenase-1 (HO-1) level in  $< 34$  and  $\geq 34$  weeks of pregnancy



**Figure 2** Box and whisker plots analysis of the difference of heme oxygenase-1 (HO-1) level in normotensive pregnancy and severe preeclampsia

2.88-5.67) ng/mL. Mann-Whitney test showed that p value: 0.001, meaning that there are significant

differences in the levels of heme oxygenase-1 (HO-1) in normotensive patients and patients with severe preeclampsia.

Based on the data presented in Figure 2, statistically significant relationship was found between the levels of heme oxygenase-1 (HO-1) with the incidence of severe preeclampsia.

## DISCUSSION

This study shows that the level of heme oxygenase-1 (HO-1) in maternal serum during pregnancy were significantly higher in women with severe preeclampsia compared with women comprising the control group of normotensive pregnancy. Furthermore, this study shows positive correlation between the levels of serum heme oxygenase-1 (HO-1) in women with severe preeclampsia compared with women comprising the control group of normotensive pregnancy.

HO is an important enzyme system in the human body. There are three isoforms of HO, inducible HO-1, constitutive HO-2, and HO-3 whose function is still unknown.<sup>9</sup> The importance of this enzyme and its catalytic product in the maintenance of normal pregnancy to term has recently been disclosed. HO catalyzes the oxidation of heme into carbon monoxide (CO), biliverdin and iron, as well as having a key role in tissue protection against oxidative stress.<sup>10</sup> Today, it is understood clearly that the HO-CO-biliverdin system is involved in the formation of a normal placenta and in the hemodynamic control of the placenta and fetal tissues. HO system regulation in the placenta is a complex process and partly dependent on the local concentration of glucose and oxygen.<sup>11,12</sup>

Most researchers found a decrease in the expression and / or activity of HO-1 in human placenta in pregnancy or in individuals with hypertensive disorders, while other researchers reported a decrease only in the levels of HO-2 and not on HO-1.<sup>13,14</sup>

Several other studies have shown that there is no difference of HO-1 protein levels between pregnancies with mild preeclampsia and uncomplicated pregnancy.<sup>10</sup> However, all these studies were conducted on the levels of HO in the placenta, and only few data are available regarding the changes in maternal serum HO-1 level in normal pregnancies and pregnancies complicated by preeclampsia. Our results show that HO-1 level is increased in plasma of women with severe preeclampsia compared with women with normal pregnancies. To our knowledge, there are only two studies, namely the research conducted Eide et al and Vitoratos et al, whose findings are in line with our findings, i.e. increased

level of HO-1 in serum and decidual tissue, accompanied by changes in the expression of the **22** *cidua*, supporting the role of these substances in **oxidative stress and excessive inflammatory responses in the pathogenesis of maternal preeclampsia.**<sup>15,16</sup>

Reactive oxygen species (ROS) will be sequestered by antioxidants, which may have non-structural **13** *teins*, such as vitamin E, C and A, as well as by **metabolites such as glutathione, ubiquinone and uric acid.** Protein based antioxidants **include** catalase, HO, glutathione peroxidase and thioredoxin peroxidase.<sup>17</sup> Normal pregnancy is characterized by a transient increase in the production of ROS partially neutralized by the induction **13** antioxidant defense mechanisms.<sup>18</sup> Preeclampsia is **associated with increases 21** oxidative stress not in the placenta but scattered in the maternal circulation, and is thought to be part of a systemic inflammatory response.<sup>19,20</sup> Increased oxidative stress occurs as a result of excessive ROS production or due to interference in antioxidant capacity,<sup>17</sup> but furthermore, **oxidative stress is closely related to the severity of preeclampsia clinically.**<sup>21</sup> **28** -1 enzyme experiences rapid upregulation by **oxidative stress and the induction of HO-1** may protect the cells by **26** -ing metaloporphyrin catalyze pro-oxidants, **such as heme, bile pigments (biliverdin, bilirubin)** that functions as **17** radicals.<sup>22</sup> We found an increased serum HO-1 in **severe preeclampsia** compared to **normal pregnancy.** Previous report **showed** that the level of oxygen radical absorbance based on the direct neutralization of free radicals is unchanged in women with mild preeclampsia.<sup>23</sup>

In contrast, free irons and particularly carbon monoxide are produced from HO-1 mediated heme catabolism.<sup>24</sup> Previous report observed an increase in serum iron concentration and carboxy-hemoglobin in preeclampsia, which reflects an increased heme and erythrocyte turnover and demonstrate that this endogenous production can change the maternal and fetal oxygenation.<sup>25</sup> Thus, there may be a potential link between HO-1 and the severity of preeclampsia. Indeed, based on our **8** *gts*, the serum levels of HO-1 seem to be related to **the severity of the disease in women with severe preeclampsia,** which also showed a positive **27** relation between the levels of HO-1 enzyme. Our findings are consistent with that of Eide and Vitoratos *et al.*, considering that there is a similar parallel relationship between serum levels of HO-1 in the **40** *ases* of preeclampsia.<sup>15,16</sup> Eide *et al.* reported that **HO-1 levels were significantly higher** among cases compared to controls (3.1±1.3 versus 1.9±0.5 ng/ml, p=0.008) and Vitorat **41** *et al.* reported the severe preeclampsia group **had significantly higher serum HO-1 levels** antepartum compared to the mild preeclampsia and normotensive groups

(5.50 ± 1.54 vs. 3.04 ± 0.72 ng/ml, p=0.0003, and 5.50 ± 1.54 vs. 3.12 ± 1.57 ng/ml, p=0.002). Although in those study the serum levels of HO-1 were positively correlated with both the overall study population and in women with preeclampsia, we cannot show such a correlation in our control group. In addition, Vitoratos *et al.* study have shown for the first time that the increased activity of HO-1 serum in severe preeclampsia will persist long after childbirth, showing the key role of persistent oxidative stress, increased vascular resistance and the chronic excessive maternal inflammatory response in the pathophysiology of preeclampsia.<sup>16</sup> However, further clinical trial with larger sample and repeated measurement du **19** *pregnancy* is required to properly evaluate the **exact role of HO-1** both in **normal pregnancy and in preeclampsia.**

There are several limitations to our study. The limited number of patients can reduce the strength of this study and increase the chances of error. As a result, a significant correlation may be missed. Measurement of HO-1 level was only performed once during pregnancy. Repeated measurements of maternal serum HO-1 level must be performed every week to allow a better assessment of the temporal changes and may explain the interaction further.

## CONCLUSION

In conclusion, this study examined **17** *the levels of maternal serum HO-1 in severe preeclampsia* as well as in normal pregnancy. Increased levels of HO-1 were found in severe preeclampsia **37** *mpared with normal pregnancy,* reflecting the **significant correlation between HO-1 levels with the severity of preeclampsia.** Statistic **31** *alysis* found a significant correlation between **heme oxygenase-1 (HO-1) level and the incidence of severe preeclampsia in < 34 and ≥ 34 weeks of pregnancy.**

## REFERENCES

1. Osungbade KO, Ige OK. Public health perspectives of preeclampsia in developing countries: implication for health system strengthening. *J Pregnancy.* 2011;1-6.
2. Khan KS, Wojdyla D, Say L, *et al.* WHO analysis of causes of maternal death: A systematic review. *Lancet.* 2006;367:1066.
3. Roeshadi HR. Upaya menurunkan angka kesakitan dan angka kematian ibu pada penderita preeklampsia dan eklampsia. Disampaikan pada pengukuhan Jabatan Guru Besar tetap dalam Bidang Ilmu Kebidanan dan Penyakit Kandungan. Medan. 2006.
4. Ros HS, Cnattingius S, Lipworth L. Comparison of risk factors for preeclampsia and gestational hypertension in a population-based cohort study. *Am J Epidemiol.* 1998; 147(11):1062-70.
5. El-Mousleh T., Casalis P.A., Wollenberg L, *et al.* Exploring the potential of low doses carbon monoxide as therapy in pregnancy complications. *Med Gas Res.* 2012;2(1):4.

6. Linzke N, Schumacher A, Woidacki K., et al. Carbon monoxide promotes proliferation of uterine natural killer cells and remodeling of spiral arteries in pregnant hypertensive hemoxygenase-1 mutant mice. *Hypertension*. 2014;63:580-8.
7. Gozzelino R, Jeney V, Soares MP. Mechanisms of cell protection by hemoxygenase-1. *Annu.Rev.Pharmacol. Toxicol*. 2010;50: 323-54.
8. Wegiel B, Hedblom A, Li M, et al. Hemoxygenase-1 derived carbon monoxide permits maturation of myeloid cells. *Cell Death Dis*. 2014a 5:e1139.
9. Bainbridge SA, Smith GN. HO in pregnancy. *Free Radic Biol Med*. 2005;38:979-88.
10. Lash GE, McLaughlin BE, MacDonald-Good fellow SK, Smith GN, Brien JF, Marks GS, Nakatsu K., Graham CH. Relationship between tissue damage and heme oxygenase expression in chorionic villi of term human placenta. *Am J Physiol Heart Circ Physiol*. 2003;284:160-7.
11. Appleton SD, Marks GS, Nakatsu K, et al. Heme oxygenase activity in placenta: direct dependence on oxygen availability. *Am J Physiol Heart Circ Physiol*. 2002;282:2055-9.
12. Appleton SD, Lash GE, Marks GS, et al. Effect of glucose and oxygen deprivation on heme oxygenase expression in human chorionic villi explants and immortalized trophoblast cells. *Am J Physiol Regul Integr Comp Physiol*. 2003;285:1453-60.
13. Barber A, Robson SC, Myatt L, et al. Heme oxygenase expression in human placenta and placental bed: reduced expression of placenta endothelial HO-2 in pre-eclampsia and fetal growth restriction. *FASEB J*. 2001;15:1158-68.
14. Zenclussen AC, Lim E, Knoeller S, et al. Heme oxygenases in pregnancy II: HO-2 is down-regulated in human pathologic pregnancies. *Am J Reprod Immunol*. 2003;50:66-76.
15. Eide IP, Isaksen CV, Salvesen KA, et al. Decidual expression and maternal serum levels of heme oxygenase 1 are increased in preeclampsia. *Acta Obstet Gynecol Scan*. 2008;87:272-9.
16. Vitoratos N, Papakonstantinou K, Deliveliotou A, et al. Antepartum and postpartum serum heme oxygenase-1 levels in preeclamptic and normotensive pregnant women. *In Vivo*. 2011;25:445- 50.
17. Vanderlelie J, Venardos K, Clifton VL, et al. Increased biological oxidation and reduced antioxidant enzyme activity in pre-eclamptic placentae. *Placenta*. 2005;26:53-8.
18. Chappell LC, Seed PT, Briley A, et al. A longitudinal study of biochemical variables in women at risk of pre-eclampsia. *Am J Obstet Gynecol*. 2002;187:127-136.
19. Jauniaux E, Poston L, Burton GJ. Placental-related diseases of pregnancy: Involvement of oxidative stress and implications in human evolution. *Hum Reprod Update*. 2006;12:747-55.
20. Lyall F, Belfort M. Ed Pre-eclampsia etiology and clinical practice. Cambridge, University Press, New York, 2007: 110.
21. Hung JH. Oxidative stress and antioxidants in pre-eclampsia. *J Chin Med Assoc*. 2007;70:430-2.
22. Stocker R, Yamamoto Y, McDonagh AF, et al. Bilirubin is an antioxidant of possible physiological importance. *Science*. 1987;235(4792):1043-6.
23. Zusterzeel PL, Steegers-Theunissen RP, Harren FJ, et al. Ethene and other biomarkers of oxidative stress in hypertensive disorders of pregnancy. *Hypertens Pregnancy*. 2002;21:39-49.
24. Schipper HM, Chertkow H, Mehindate K, et al. Evaluation of heme oxygenase-1 as a systemic biological marker of sporadic AD. *Neurol*. 2000;54:1297- 1304.
25. Entman SS, Kambam JR, Bradley CA, et al. Increased levels of carboxyhemoglobin and serum iron as an indicator of increased red cell turnover in preeclampsia. *Am J Obstet Gynecol*. 1987;156:1169-73.



This work is licensed under a Creative Commons Attribution

# Relationship of Heme Oxygenase-1 - HO-1 Level with Onset and Severity in Normotensive Pregnancy and Severe Preeclampsia

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

16%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.jcdr.net">www.jcdr.net</a> Internet Source	1%
2	Aleksandra Nikolic, Velibor Cabarkapa, Aleksandra Novakov Mikic, Ana Jakovljević, Zoran Stosic. "Ceruloplasmin and antioxidative enzymes in pre-eclampsia", The Journal of Maternal-Fetal & Neonatal Medicine, 2015 Publication	1%
3	<a href="http://www.perinataljournal.com">www.perinataljournal.com</a> Internet Source	1%
4	Shannon A. Bainbridge, Graeme N. Smith. "HO in pregnancy", Free Radical Biology and Medicine, 2005 Publication	1%
5	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	1%
6	<a href="http://medicalgasresearch.biomedcentral.com">medicalgasresearch.biomedcentral.com</a> Internet Source	1%

---

7	Valéria C. Sandrim, Mayara Caldeira-Dias, Heloisa Bettiol, Marco Antonio Barbieri et al. "Circulating Heme Oxygenase-1: Not a Predictor of Preeclampsia but Highly Expressed in Pregnant Women Who Subsequently Develop Severe Preeclampsia", Oxidative Medicine and Cellular Longevity, 2018 Publication	1 %
8	epdf.pub Internet Source	1 %
9	repository.unp.ac.id Internet Source	1 %
10	Submitted to Universitas Diponegoro Student Paper	1 %
11	journal.frontiersin.org Internet Source	1 %
12	synapse.koreamed.org Internet Source	1 %
13	Clifton, V.L.. "Increased anti-oxidant enzyme activity and biological oxidation in placentae of pregnancies complicated by maternal asthma", Placenta, 200511 Publication	1 %
14	www.thieme-connect.de Internet Source	1 %

---



15 S. Magnusson, J. Ekström, E. Elmér, M. Kanje, L. Ny, P. Alm. "Heme oxygenase-1, heme oxygenase-2 and biliverdin reductase in peripheral ganglia from rat, expression and plasticity", Neuroscience, 1999

Publication

1 %

16 Breno Fernando Martins de Almeida, Kathlenn Liezbeth Oliveira Silva, Vanessa Marim Chiku, Aline Aparecida Correa Leal et al. "The effects of increased heme oxygenase-1 on the lymphoproliferative response in dogs with visceral leishmaniasis", Immunobiology, 2017

Publication

<1 %

17 Julian Dewantiningrum, Alini Hafiz. "The role of glutathione peroxidase maternal serum level in late onset of severe preeclampsia", Hypertension in Pregnancy, 2016

Publication

<1 %

18 [journals.lww.com](http://journals.lww.com)

Internet Source

<1 %

19 Eric M George, Joey P Granger. "Recent insights into the pathophysiology of preeclampsia", Expert Review of Obstetrics & Gynecology, 2014

Publication

<1 %

20 Luis M. Gomez, Lauren Anton, Shindu K. Srinivas, Michal A. Elovitz, Samuel Parry. "Low-Dose Aspirin May Prevent Trophoblast

<1 %

# Dysfunction in Women With Chlamydia Pneumoniae Infection", Reproductive Sciences, 2018

Publication

---

21

E. M. George. "Induction of heme oxygenase-1 shifts the balance from pro-injury to pro-survival in the placentas of pregnant rats with reduced uterine perfusion pressure", AJP Regulatory Integrative and Comparative Physiology, 01/11/2012

Publication

---

<1 %

22

S. Kharb, J. Bala, S. Nanda. "Markers of obesity and growth in preeclamptic and normotensive pregnant women", Journal of Obstetrics and Gynaecology, 2017

Publication

---

<1 %

23

Angela Maria Rizzo, Patrizia Berselli, Stefania Zava, Gigliola Montorfano, Manuela Negroni, Paola Corsetto, Bruno Berra. "Chapter 5 Endogenous Antioxidants and Radical Scavengers", Springer Science and Business Media LLC, 2010

Publication

---

<1 %

24

Eishi Hara, Kazuhiro Takahashi, Kazuhisa Takeda, Masaharu Nakayama et al. "Induction of heme oxygenase-1 as a response in sensing the signals evoked by distinct nitric

<1 %

oxide donors", Biochemical Pharmacology,  
1999

Publication

---

25

[pt.scribd.com](http://pt.scribd.com)

Internet Source

<1 %

---

26

Heme Oxygenase in Biology and Medicine,  
2002.

Publication

<1 %

---

27

Tevfik NOYAN. "Serum advanced oxidation  
protein products, myeloperoxidase and  
ascorbic acid in pre-eclampsia and  
eclampsia", The Australian and New Zealand  
Journal of Obstetrics and Gynaecology,  
12/2006

Publication

<1 %

---

28

Vivian B. Neis, Priscila B. Rosa, Morgana  
Moretti, Ana Lucia S. Rodrigues. "Involvement  
of Heme Oxygenase-1 in Neuropsychiatric  
and Neurodegenerative Diseases", Current  
Pharmaceutical Design, 2018

Publication

<1 %

---

29

[hyper.ahajournals.org](http://hyper.ahajournals.org)

Internet Source

<1 %

---

30

[journals.plos.org](http://journals.plos.org)

Internet Source

<1 %

---

31

[repositorio.uam.es](http://repositorio.uam.es)

Internet Source

<1 %

---

32

[repository.ubn.ru.nl](http://repository.ubn.ru.nl)

Internet Source

&lt;1 %

33

[www.alliedacademies.org](http://www.alliedacademies.org)

Internet Source

&lt;1 %

34

[www.physiology.org](http://www.physiology.org)

Internet Source

&lt;1 %

35

G. E. Lash, B. E. McLaughlin, S. K. MacDonald-Goodfellow, G. N. Smith, J. F. Brien, G. S. Marks, K. Nakatsu, C. H. Graham.

"Relationship between tissue damage and heme oxygenase expression in chorionic villi of term human placenta", American Journal of Physiology-Heart and Circulatory Physiology, 2003

Publication

&lt;1 %

36

James R. Reed, George F. Cawley, Wayne L. Backes. "Inhibition of Cytochrome P450 1A2-Mediated Metabolism and Production of Reactive Oxygen Species by Heme Oxygenase-1 in Rat Liver Microsomes", Drug Metabolism Letters, 2011

Publication

&lt;1 %

37

Jeffrey E. Schmidt. "Heme oxygenase-1 messenger RNA expression is induced in peripheral blood mononuclear cells of pediatric cancer patients with systemic

&lt;1 %

inflammatory response syndrome", *Pediatric Critical Care Medicine*, 11/2004

Publication

---

38

Song Kwon Choi, Jong Yun Hwang, Jiyeon Lee, Sung Hun Na, Joong Gyu Ha, Hyang Ah Lee, Dong Hun Lee. " Gene expression of and on trophoblast of preeclampsia ", *Korean Journal of Obstetrics*, 2011

Publication

---

39

Vittorio Calabrese. "Redox Regulation of Heat Shock Protein Expression by Signaling Involving Nitric Oxide and Carbon Monoxide: Relevance to Brain Aging, Neurodegenerative Disorders, and Longevity", *Antioxidants and Redox Signaling*, 03/2006

Publication

---

40

Michiko Miya, Aikou Okamoto, Takashi Nikaido, Rie Tachimoto-Kawaguchi, Tadao Tanaka. "Immunohistochemical localization of heme oxygenase-1 and bilirubin/biopyrrin of heme metabolites as antioxidants in human placenta with preeclampsia", *Hypertension in Pregnancy*, 2019

Publication

---

41

Yohei Kirino. *Arthritis Research & Therapy*, 2005

Publication

---

<1 %

<1 %

<1 %

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On