

Isolation and Identification of Arsenic-Resistant Bacteria for Possible Application in Arsenic Bioremediation

by Billy Kepel 6

Submission date: 24-Feb-2021 07:42AM (UTC+0700)

Submission ID: 1516585434

File name: Bacteria_for_Possible_Application_in_Arsenic_Bioremediation.pdf (567.73K)

Word count: 3293

Character count: 18849

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

**Pakistan
Journal of Biological Sciences**

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan



Research Article

Isolation and Identification of Arsenic-resistant Bacteria for Possible Application in Arsenic Bioremediation

¹Billy Kepel, ¹Widdhi Bodhi, ²Fatimawali and ³Trina Ekawati Tallei

¹Department of Chemistry, Faculty of Medicine, Sam Ratulangi University, Kampus Unsrat Kleak, 95115 Manado, Indonesia

²Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Sam Ratulangi University, Kampus Unsrat Kleak, 95115 Manado, Indonesia

³Department of Biology, Faculty of Mathematics and Natural Sciences, Sam Ratulangi University, Kampus Unsrat Kleak, 95115 Manado, Indonesia

Abstract

Background and Objective: North Sulawesi is rich in minerals, among them gold is also present. The gold mining in the Buyat area produces heavy metal waste which can pollute the environment, among others is arsenic. Arsenic is a heavy metal that is very toxic to humans, so an agent is needed for the remediation process. The aim of this study is to isolate and identify arsenic-resistant bacteria from the Buyat estuary and beach to analyze the isolates' ability to detoxify arsenic. **Materials and Methods:** Soil sediment samples were obtained from Buyat estuary and beach in North Sulawesi. Isolation of arsenic-resistant bacteria was carried out by growing the samples in LB broth media containing 100, 500 and 1000 ppm arsenite. Identification of arsenic-resistant bacteria was carried out by microbiological, biochemical and biomolecular analysis. The ability to detoxify arsenite was analyzed by CVAFS. **Results:** The study showed that there were 4 isolates of arsenic-resistant bacteria isolated from the soil samples. All isolates are rod-shaped, Gram-negative and non-motile bacteria. BLAST results showed that isolate A was *Stenotrophomonas* sp., isolate B was *Stenotrophomonas maltophilia*, isolate C was *Pseudomonas* sp. and isolate D was *Pseudomonas putida*. All isolates reduced the levels of arsenic in media by almost 100% within 72 h. **Conclusion:** The study suggested that *Stenotrophomonas* sp., *S. maltophilia*, *Pseudomonas* sp. and *P. putida* had the potentials to be used in the bioremediation of arsenic.

Key words: Arsenic remediation, arsenic-resistant bacteria, arsenic detoxification, Buyat Bay, Buyat estuary

Citation: Billy Kepel, Widdhi Bodhi, Fatimawali and Trina Ekawati Tallei, 2020. Isolation and identification of arsenic-resistant bacteria for possible application in arsenic bioremediation. Pak. J. Biol. Sci., 23: 63-67.

Corresponding Author: Trina Ekawati Tallei, Department of Biology, Faculty of Mathematics and Natural Sciences, Sam Ratulangi University, Kampus Unsrat Kleak, 95115 Manado, Indonesia Tel: +628179255866

Copyright: © 2020 Billy Kepel *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Arsenic, a natural element that is widespread in the earth's crust, is chemically classified as metalloid, having both metal and non-metal properties. It exists in the form of arsine/arsane, elemental arsenic (0), arsenite (+3) and arsenate (+5). According to the Agency for Toxic Substances and Disease Registry¹, arsenic which is usually found in a combination in the environment with other elements such as oxygen, chlorine and sulfur is called inorganic arsenic, while when combined with carbon and hydrogen is called organic arsenic. Inorganic arsenic is naturally found in soils and in various types of rock, especially in minerals and ores that contain copper or lead. When this ore is heated in a smelter, most of the arsenic builds up and enters the air as fine dust. Smelters can collect this dust and form arsenic compounds called arsenic trioxide (As₂O₃)².

In rock or soil, arsenic is distributed as a mineral. The highest arsenic level is in the form of arsenide from copper, lead, silver and sulfide forms from gold. Human involvement in causing or exacerbating arsenic pollution can be traced to mining related activities^{3,4}. Sediment samples from Buyat and Ratatotok bays were studied in 2002 and 2004 for the analysis of minerals and trace elements⁵. The sediment samples at this location contained heavy metal wastes generated by gold mining activities³. Their study found high concentrations of arsenic (590-660 ppm) in shallow marine sediments both from Buyat Bay and Ratatotok. Arsenic concentration in the interior of Buyat Bay ranged from 0-25 ppm. Furthermore, a study on characterization of the metal composition of river mine tailings and marine sediments in Buyat and Ratatotok was conducted⁵. The study showed that sediments affected by tailings had an arsenic concentration 30 times than uncontaminated sediments. This was confirmed again in a report conducted by an integrated team of the Ministry of Environment in 2004, which stated that illegal mining resulting from the amalgamation process waste had been dumped into a tributary upstream of the Ratatotok river⁶. Research conducted in 2010 showed that the highest arsenic levels at 6 sample points in Buyat and Ratatotok bays were found in the Ratatotok coastal area (127.8 ppm). The highest arsenic level at this location was associated with the accumulation of arsenic waste originating from the Ratatotok river estuary which had arsenic levels of 4-130 ppm. This shows that the disposal of waste through rivers into the sea caused by illegal mining has contributed to polluting the Ratatotok bay^{7,8}.

Arsenic is a pollutant for the environment, the search for arsenic remediation agents is urgently needed. Arsenic

resistant-bacteria are the target of many studies in order to use them as bioremediation agents. Arsenic resistance in bacteria is mediated by the presence of the ars operon gene. The ars operon consists of arsC operon which encodes arsenic reductase, arsB and arsA which encodes the arsenic efflux pump, arsR which encodes the transcription regulator and arsD which encodes for the transfer of arsenite from the glutathione complex to the arsA subunit of the arsAB complex and activates it (Satyapal *et al.*⁹ and Kruger *et al.*¹⁰).

Various studies have identified arsenic-resistant bacteria from water and soil. In addition, research on livestock meat was also carried out. The types of bacteria found in water and in the soil included *Klebsiella pneumoniae*, *Enterobacter* sp., *Bacillus* sp., *Aneurinibacillus aneurinilyticus*, *Enterobacter asburiae* and *E. cloacae*¹¹⁻¹³. While *Campylobacter* spp. was found in the meat of cattle¹⁴. These studies showed that the bacterial colonies that exist in a community can adapt to pollutants such as arsenic, through resistance mechanisms or what is known as pollution-induced community tolerance (PICT)¹⁵.

Research on the use of arsenic-resistant bacteria for bioremediation is still very limited, therefore the purpose of this research was to isolate and identify arsenic resistant-bacteria that were resistant to high concentration of arsenite obtained from sediment soil of Buyat estuary and beach which reduced arsenite levels in the media.

MATERIALS AND METHODS

Sampling sites: Sediment samples were obtained from Buyat estuary and beach which are located on the southern coast of the Minahasa Peninsula, North Sulawesi, Indonesia. The study was conducted from March-August, 2019. In each location, three sampling sites were chosen. In each sampling site, the pH was measured and the sediment samples were put into sterile polyethylene plastics and brought to the laboratory using a cool box for further analysis.

Isolation of arsenic-resistant bacteria: Each of soil sediment was added with sterile 10 mL of 0.9% NaCl and stirred evenly. Into each of 10 mL of sterile Luria Bethani (LB) medium, 100 µL of NaCl solution containing sediment samples and mixed evenly. From each sample, 100 µL was poured onto Nutrient Agar (NA) media, spread evenly and grown at 37 °C for 2 days. Several single colonies with different morphological appearance were picked from the agar and purified by the streak method to obtain pure single colonies. Each pure colony was grown on Nutrient Broth (NB) supplemented with

500 ppm and 1000 ppm of arsenic (As_2O_3) and incubated for 24 h at 37°C. Cultures grown on arsenic media were transferred onto NA containing 500 and 1000 ppm arsenic. Each different colony was grown by the streak method to obtain pure colony. Colonies with the different morphological appearance which survived on arsenic NA media were identified. The identification procedure is described in the section 16S rRNA-based identification.

Bioremediation simulation test for arsenic-resistant bacterial isolates: The bacterial isolates that were survived in 500 and 100 ppm of arsenite were inoculated in LB broth containing 100 ppm arsenite and incubated at 30°C. After incubating for 72 h, the solution was centrifuged at 10,000 rpm for 10 min to separate bacteria from the media, then the concentration of arsenic in the media was measured using the Cold Vapor Atomic Absorption Spectroscopy (CVAAS).

Morphological-, physiological- and biochemical-based identification of arsenic-resistance bacterial isolates: Morphological- and physiological-based identification were carried out using Gram staining, bacterial cell shape and motility tests. Biochemical-based identification was performed based on IMViC (Indol, Methyl red, Voges-Proskauer and Citrate) tests.

16S rRNA-based identification: The identification of the isolates based on 16S rRNA followed the previous protocol¹⁶. The extraction and purification of bacterial DNA were performed using Genomic DNA Mini Kit (Geneaid). The pure DNA was stored at 2-8°C for further analysis. DNA amplification using universal BactF1 and UniB1 primer pairs were done in PCR combi block machine (Whatman Biometra Germany). The composition of the PCR reaction is ddH₂O 20 µL, UniB1 primer 30 pmol µL⁻¹ 1.5 µL, BactFI primer 30 pmol µL⁻¹ 1.5 µL, 2x MyTaq HS Red Mix 25 µL and DNA template 2 µL. The PCR reaction condition is as follow: An initial denaturation 95°C for 180 sec, followed by 30 cycles of denaturation at 95°C for 30 sec, annealing at 72°C for 30 sec and extension at 72°C for 90 sec. The final extension was done at 72°C for 60 sec. The PCR products were sent to 1 Base Malaysia for sequencing.

Analysis of sequencing result of 16S rRNA fragments: Sequencing analysis followed the protocol provided by Suresh *et al.*¹⁷. Briefly, chromatograms were edited using Geneious R10 software. The sequences were trimmed

approximately 50 nucleotides at both ends. The edited sequences were subjected to Basic Local Alignment Search Tool (BLAST) (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) to query for highly similar sequences. The highest identity results determined the expected species or genera.

RESULTS AND DISCUSSION

In this study, 4 isolates were able to be isolated and purified based on different morphological appearance on LB agar media with each concentration of arsenic. Isolates A and B were purified from agar containing 500 ppm arsenic while isolates C and D were purified from agar containing 1000 ppm arsenic. All purified isolates were subsequently inoculated on the agar slant media for further identification. The identification results based on morphological and physiological tests are presented in Table 1. All isolates were rod-shaped, Gram negative and nonmotile bacteria. Based on the test, isolates A and B gave similar results, while C and D gave different results.

The molecular identification based on 16S rRNA gene is presented in Table 2. Isolate A had 93% identity with *Stenotrophomonas maltophilia* strain C-1, isolate B had 99.8% with *S. maltophilia* strain C-1, isolate C had 94.63% identity with *Pseudomonas entomophila* strain AS90 and isolate D had 99.73% identity with *P. putida* strain ZJB-LLJ. The lower identity results for isolates A and C raise doubts about the results of identification to state that they are the species in question. Some researchers have proposed different thresholds for bacterial species delineation using 16S rRNA gene including 18-20 98.65, 95-96 and 97%. This indicates that the cut off is set to minimum 95% for 16S rRNA. Using this threshold, it implies that isolate A is not *Stenotrophomonas maltophilia* but *Stenotrophomonas* sp., even though using the morphological, physiological and biochemical tests showed the similar morphology and characteristics. Using the same principle, then isolate C is not *Pseudomonas entomophila* but *Pseudomonas* sp.

Several bacteria belonging to the genera *Acidithiobacillus*, *Bacillus*, *Deinococcus*, *Desulfitobacterium* and *Pseudomonas* have been reported to be resistant to arsenic^{21,22}. Resistance to arsenic species in both Gram-positive and Gram-negative bacteria is resulted from energy-dependent efflux of either arsenate or arsenite from the cell mediated via the ars operon²³. In *E. coli*, an arsA-arsB complex functions as a primary arsenite pump²⁴. In *S. aureus*, arsB alone is sufficient to act as a chemiosmotic secondary transport system for arsenite resistance without the presence of an arsA ATPase²⁵.

Table 1: Identification results based on morphological and physiological tests

Tests	Code of isolates			
	A	B	C	D
Morphology	Rod-shaped	Rod-shaped	Rod-shaped	Rod-shaped
Gram staining	-	-	-	-
Motility	-	-	-	-
Glucose	+	+	+	+
Indole	-	-	+	+
Citrate	+	+	+	+
Lysine	-	-	-	+
H ₂ S	-	-	-	-
H ₂ O ₂	+	+	+	+

+: Present, -: Absent

Table 2: Results of BLAST analysis of 16S rRNA sequence

Isolates	Species description	Max. identity (%)	Query cover (%)
A	<i>Stenotrophomonas maltophilia</i> strain C-1	93.00	99
B	<i>Stenotrophomonas maltophilia</i> strain C-1	99.80	99
C	<i>Pseudomonas entomophila</i> strain AS90	94.63	99
D	<i>Pseudomonas putida</i> strain ZJB-LLJ	99.73	99

⁴ *Stenotrophomonas maltophilia* is a commensal and an emerging pathogen earlier noted in broad-spectrum life threatening infections among the vulnerable, but more recently as a pathogen in immune-competent individual. This bacterium is also an environmental bacteria occurring in water, rhizospheres, as part of the animals' microflora, in foods and several other microbiota²⁶. *Pseudomonas putida* and *Pseudomonas mendosina*, ¹³ found in arsenic-contaminated environments in Bulgaria, which were of particular interest, because they appeared to gain metabolic energy during the arsenic transformation, were characterised²⁷. Bacteria adapt to arsenic toxicity mainly by the development of resistance mechanisms conferred by chromosomal or plasmid-encoded arsenical resistance (ars) operons. Once the trivalent form of the metalloid accumulates in the cell, resistance is produced by their removal from the cytosol²⁸. The results of the arsenite absorption of isolates that ⁵ analyzed by CVAFS showed that all isolates can reduce the levels of arsenic in media by almost 100% within 72 h. This indicates that all isolates have the ability to be further investigated as bioremediation agents in the arsenic-contaminated sites.

CONCLUSION

The present study revealed that *Stenotrophomonas maltophilia*, *Stenotrophomonas* sp., *Pseudomonas* sp. and *P. putida* isolated from Buyat estuary and beach were able to reduce the arsenite concentration quickly. These bacteria can be investigated further for their potential to remediate the environment with arsenic contamination.

SIGNIFICANCE STATEMENT

This study discovered that *Stenotrophomonas maltophilia*, *Stenotrophomonas* sp., *Pseudomonas* sp. and *Pseudomonas putida* showed the high ability to detoxify arsenite and can be used in arsenic-contamination site for bioremediation of arsenic. However, it is advisable to develop a safe method of arsenic detoxification by using enzymatic methods which can be carried out in further studies.

ACKNOWLEDGMENT

This study was financed by DIPA Sam Ratulangi University Ministry of Research, Technology and Higher Education, Republic of Indonesia, through Sam Ratulangi University Basic Excellence Research Schemes (RDUU) fiscal year 2019 (Grant Number: SP DIPA-042.01.2.400959/2019).

REFERENCES

1. Agency for Toxic Substances and Disease Registry, 2007. Toxicological profile of arsenic. U.S. Department of Health and Human Services, Atlanta.
2. Geiger, A. and J. Cooper, 2010. Overview of airborne metals regulations, exposure limits, health effects and contemporary research. <http://cooperenvironmental.com/wp-content/uploads/2014/09/Airborne-Metals-Health-Effects1.pdf>
3. Garelick, H., H. Jones, A. Dybowska and E. Valsami-Jones, 2008. Arsenic pollution sources. Rev. Environ. Contam. Toxicol., 197: 17-60.
4. Ilahude, D. and E. Herawati, 2010. Heavy metal contents in marine sediments and seawater at Totok bay area, North Sulawesi. Bull. Mar. Geol., 25: 39-51.

5. Edinger, E.N., P.R. Siregar and G.M. Blackwood, 2007. Heavy metal concentrations in shallow marine sediments affected by submarine tailings disposal and artisanal gold mining, Buyat-Ratototok district, North Sulawesi, Indonesia. *Environ. Geol.*, 52: 701-714.
6. Kementerian Lingkungan Hidup, 2004. Laporan penelitian penanganan dugaan kasus pencemaran dan/atau perusakan lingkungan hidup di Desa Buyat Pantai dan Desa Ratatotok Kecamatan Ratatotok Timur Kabupaten Minahasa Selatan Propinsi Sulawesi Utara. Kementerian Lingkungan Hidup Dan Kehutanan, Jakarta.
7. Lahar, H., 2004. Evaluasi sumber daya dan cadangan bahan galian untuk pertambangan skala kecil daerah Lembar Manado, Provinsi Sulawesi Utara. Kolokium Hasil Lapangan DIM., 45: 1-12.
8. Rumampuk, N.D.C. and V. Warouw, 2015. Bioakumulasi total merkuri, arsen, kromium, cadmium, timbal di teluk Totok dan teluk Buyat, Sulawesi Utara. *J. LPPM Sains Teknol.*, 2: 49-59.
9. Satyapal, G.K., S. Rani, M. Kumar and N. Kumar, 2016. Potential role of arsenic resistant bacteria in bioremediation: Current status and future prospects. *J. Microb. Biochem. Technol.*, 8: 256-258.
10. Kruger, M.C., P.N. Bertin, H.J. Heipieper and F. Arsène-Ploetze, 2013. Bacterial metabolism of environmental arsenic-mechanisms and biotechnological applications. *Applied Microbiol. Biotechnol.*, 97: 3827-3841.
11. Abbas, S.Z., M. Riaz, N. Ramzan, M.T. Zahid, F.R. Shakoori and M. Rafatullah, 2014. Isolation and characterization of arsenic resistant bacteria from wastewater. *Braz. J. Microb.*, 45: 1309-1315.
12. Dey, U., S. Chatterjee and N.K. Mondal, 2016. Isolation and characterization of arsenic-resistant bacteria and possible application in bioremediation. *Biotechnol. Rep.*, 10: 1-7.
13. Selvi, M.S., S. Sasikumar, S. Gomathi, P. Rajkumar, P. Sasikumar and S. Govindan, 2014. Isolation and characterization of arsenic resistant bacteria from agricultural soil and their potential for arsenic bioremediation. *Int. J. Agric. Policy Res.*, 2: 393-405.
14. Sapkota, A.R., L.B. Price, E.K. Silbergeld and K.J. Schwab, 2006. Arsenic resistance in *Campylobacter* spp. isolated from retail poultry products. *Applied Environ. Microbiol.*, 72: 3069-3071.
15. Tlili, A., A. Berard, H. Blanck, A. Bouchez and F. Cássio *et al.*, 2016. Pollution-Induced Community Tolerance (PICT): Towards an ecologically relevant risk assessment of chemicals in aquatic systems. *Freshwater Biol.*, 61: 2141-2151.
16. Oremland, R.S., J.F. Stolz and J.T. Hollibaugh, 2004. The microbial arsenic cycle in Mono Lake, California. *FEMS Microbiol. Ecol.*, 48: 15-27.
17. Suresh, K., G.S.N. Reddy, S. Sengupta and S. Shivaji, 2004. *Deinococcus indicus* sp. nov., an arsenic-resistant bacterium from an aquifer in West Bengal, India. *Int. J. Syst. Evol. Microbiol.*, 54: 457-461.
18. Tseng, C.H., 2007. Arsenic methylation, urinary arsenic metabolites and human diseases: Current perspective. *J. Environ. Sci. Health Part C*, 25: 1-22.
19. Yoon, K.P., 2005. Stabilities of artificially transconjugated plasmids for the bioremediation of cocontaminated sites. *J. Microbiol.*, 43: 196-203.
20. Wolfe-Simon, F., J.S. Blum, T.R. Kulp, G.W. Gordon and S.E. Hoefft *et al.*, 2011. A bacterium that can grow by using arsenic instead of phosphorus. *Science*, 332: 1163-1166.
21. Adegoke, A.A., T.A. Stenström and A.I. Okoh, 2017. *Stenotrophomonas maltophilia* as an emerging ubiquitous pathogen: Looking beyond contemporary antibiotic therapy. *Front. Microbiol.*, Vol. 8. 10.3389/fmicb.2017.02276.
22. Krumova, K., M. Nikolovska and V. Groudeva, 2008. Characterization of arsenic-transforming bacteria from arsenic contaminated sites in Bulgaria. *Biotech. Biotech. Equip.*, 22: 729-735.
23. Ramos, J.L. and A. Filloux, 2010. *Pseudomonas*: Volume 6: Molecular Microbiology, Infection and Biodiversity. Springer, Heidelberg, Germany, Pages: 267.
24. Fatimawali, B. Kepel and T.E. Tallei, 2019. Potential of organic mercury-resistant bacteria isolated from mercury contaminated sites for organic mercury remediation. *Pak. J. Biol. Sci.*, 22: 45-50.
25. Tallei, T.E. and B.J. Kolondam, 2015. DNA barcoding of Sangihe Nutmeg (*Myristica fragrans*) using *matK* gene. *HAYATI J. Biosci.*, 22: 41-47.
26. Kim, M. and J. Chun, 2014. 16S rRNA gene-based identification of *Bacteria* and *Archaea* using the EzTaxon server. *Methods Microbiol.*, 41: 61-74.
27. Kim, M., H.S. Oh, S.C. Park and J. Chun, 2014. Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. *Int. J. Syst. Evol. Microbiol.*, 64: 346-351.
28. Mysara, M., P. Vandamme, R. Props, F.M. Kerckhof and N. Leys *et al.*, 2017. Reconciliation between operational taxonomic units and species boundaries. *FEMS Microbiol. Ecol.*, Vol. 93, No. 4. 10.1093/femsec/fix029.

Isolation and Identification of Arsenic-Resistant Bacteria for Possible Application in Arsenic Bioremediation

ORIGINALITY REPORT

20%

SIMILARITY INDEX

11%

INTERNET SOURCES

16%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to Universitas Sam Ratulangi

Student Paper

1%

2

link.springer.com

Internet Source

1%

3

Muh. I. Hidayat, Aaltje Manampiring, Billy J. Kepel. "Isolasi dan Identifikasi Bakteri Resisten Arsen pada Sedimen Tanah di Pesisir Pantai Ratatotok", Jurnal e-Biomedik, 2018

Publication

1%

4

www.labome.org

Internet Source

1%

5

Sana Irshad, Zuoming Xie, Sajid Mehmood, Asad Nawaz, Allah Ditta, Qaisar Mahmood. "Insights into conventional and recent technologies for arsenic bioremediation: A systematic review", Environmental Science and Pollution Research, 2021

Publication

1%

6

pubmed.ncbi.nlm.nih.gov

Internet Source

1%

7	www.atsdr.cdc.gov Internet Source	1 %
8	Guangya Zheng, Jupei Xia, Hailang Liu, Zhengjie Chen. "Arsenic removal from acid extraction solutions of copper smelting flue dust", <i>Journal of Cleaner Production</i> , 2021 Publication	1 %
9	M. Pepi. "Arsenic-resistant bacteria isolated from contaminated sediments of the Orbetello Lagoon, Italy, and their characterization", <i>Journal of Applied Microbiology</i> , 7/5/2007 Publication	1 %
10	Sahar Alipour Kafi, Saeid Arabhosseini, Ebrahim Karimi, Parisa Koobaz, Ali Mohammadi, Akram Sadeghi. "Pseudomonas putida P3-57 induces cucumber (<i>Cucumis sativus</i> L.) defense responses and improves fruit quality characteristics under commercial greenhouse conditions", <i>Scientia Horticulturae</i> , 2021 Publication	1 %
11	Shaheen B. Humayoun, Lari M. Hiott, Sushim K. Gupta, John B. Barrett et al. "An assay for determining the susceptibility of <i>Salmonella</i> isolates to commercial and household biocides", <i>PLOS ONE</i> , 2018 Publication	1 %

12

Varenyam Achal, Xiangliang Pan, Qinglong Fu, Daoyong Zhang. "Biom mineralization based remediation of As(III) contaminated soil by *Sporosarcina ginsengisoli*", *Journal of Hazardous Materials*, 2012

Publication

1 %

13

Submitted to Lovely Professional University

Student Paper

1 %

14

Uttiya Dey, Soumendranath Chatterjee, Naba Kumar Mondal. "Isolation and characterization of arsenic-resistant bacteria and possible application in bioremediation", *Biotechnology Reports*, 2016

Publication

1 %

15

Uttiya Dey, Soumendranath Chatterjee, Naba Kumar Mondal. "Investigation of Bioremediation of Arsenic by Bacteria Isolated from an Arsenic Contaminated Area", *Environmental Processes*, 2016

Publication

1 %

16

doaj.org

Internet Source

1 %

17

Sánchez-Virosta, P., S. Espín, A.J. García-Fernández, and T. Eeva. "A review on exposure and effects of arsenic in passerine birds", *The Science of The Total Environment*, 2015.

Publication

1 %

18	Submitted to Walden University Student Paper	1 %
19	Submitted to Bethesda-Chevy Chase High School Student Paper	<1 %
20	www.ijsrp.org Internet Source	<1 %
21	www.nature.com Internet Source	<1 %
22	K. Krumova, M. Nikolovska, V. Groudeva. "Isolation and Identification of Arsenic-Transforming Bacteria from Arsenic Contaminated Sites in Bulgaria", Biotechnology & Biotechnological Equipment, 2014 Publication	<1 %
23	Olubisi E. Adeyi, David O. Babayemi, Babajide O. Ajayi, Akindele O. Adeyi et al. "Co-administration of sodium selenite and sodium arsenite exacerbates hepatic, renal, pulmonary and splenic inflammation in rats", Scientific African, 2021 Publication	<1 %
24	www.nrcresearchpress.com Internet Source	<1 %
25	www.tnrc.org Internet Source	<1 %

26 Fengcai Ye, Dafei Gong, Caoping Pang, Jieling Luo, Xiaomei Zeng, Changhua Shang. "Analysis of Fungal Composition in Mine-Contaminated Soils in Hechi City", *Current Microbiology*, 2020

Publication

27 Jean, . "Arsenic in terrestrial ecosystems and biota", *Arsenic in Geosphere and Human Diseases Arsenic 2010 Proceedings of the Third International Congress on Arsenic in the Environment (As-2010)*, 2010.

Publication

28 Jebeli, Mohammad Ahmadi, Afshin Maleki, Mohammad Ali Amoozegar, Enayatollah Kalantar, Hassan Izanloo, and Fardin Gharibi. "Bacillus flexus strain As-12, a new arsenic transformer bacterium isolated from contaminated water resources", *Chemosphere*, 2017.

Publication

29 Payal Mazumder, Subhash Kumar Sharma, Kaling Taki, Ajay S. Kalamdhad, Manish Kumar. "Microbes involved in arsenic mobilization and respiration: a review on isolation, identification, isolates and implications", *Environmental Geochemistry and Health*, 2020

Publication

30 Ravindranath Shashidhar, Jayant R. Bandekar. " sp. nov., a radiation-resistant

pleomorphic bacterium isolated from
Mumbai, India ", FEMS Microbiology Letters,
2006

Publication

31

"Microbial Biotechnology", Springer Science
and Business Media LLC, 2017

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On