

# Deep sea squid in Sulawesi Sea, North Sulawesi Province, Indonesia

*by* Silvester Pratasik 36

---

**Submission date:** 09-May-2022 09:42AM (UTC+0700)

**Submission ID:** 1831584611

**File name:** Deep-sea\_Squid-Biodiversitas.pdf (515.16K)

**Word count:** 4759

**Character count:** 25193

## Deep sea squid in Sulawesi Sea, North Sulawesi Province, Indonesia

SILVESTER BENNY PRATASIK\*, LAURENTIUS TH.X. LALAMENTIK, LEFRAND MANOPPO,  
JOHNNY BUDIMAN

25  
Faculty of Fisheries and Marine Science, Universitas Sam Ratulangi, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia.  
Tel/fax: +62-431-863886, \*email: spjong07@yahoo.com

Manuscript received: 15 May 2021. Revision accepted: 15 March 2022.

**Abstract.** *Pratik SB, Lalamentik LTHX, Manoppo L, Budiman J. 2022. Deep sea squid in Sulawesi Sea, North Sulawesi Province, Indonesia. Biodiversitas 23: 1774-1779.* This study aimed to identify one of the deep sea squids caught in artisanal fisheries by traditional fishermen in Sulawesi Sea, North Sulawesi Province, Indonesia. Samples were collected using the vertical hand line of traditional fishermen. The specimen was identified using Cytochrome c oxidase subunit I (COI). For morphometric characteristics, all body parts were measured, and each part was compared with mantle size. Results showed that the specimen was identified as purpleback flying squid *Sthenoteuthis oualaniensis* and had 99.35% similarity to *S. oualaniensis* (COI) from China (NCBI - MN101944) with sufficient genetic diversity. Based on the body size, species recorded in the present study belong to the dwarf form, the smallest form of *S. oualaniensis* groups. The species has the following morphometric characteristics: Head length is 0.28 of mantle length, tentacle length is 1.158 of mantle length, and fin area is 0.5 of mantle length. The fourth arm is the shortest and the second arm is the longest. *S. oualaniensis* recorded in the present study belonged to dwarf form living in equatorial waters. This finding may contribute to the list of marine resources of fisheries in Indonesian waters.

**Keywords:** COI gene, hand line, morphometric, species identification, traditional fishermen

### INTRODUCTION

Squids are cephalopods that live in the marine environment. This group is an ecological opportunist adapted to exploit favorable environmental conditions (Rodhouse 2013). It is a favorite food type due to its high nutritional content with high-quality protein for human consumption and nearly all body parts are edible (Triharyuni and Puspasari 2012). Cephalopod production from fishing has continued to grow (Doubleday et al. 2016), with total commercial annual catches varying between 3.5 and 4.9 million tonnes in 2008-2017 (FAO 2019) and averagely supports about 15 and 20% of marine fishery landings and landed values, respectively (FAO 2019). Demand for this commodity has increased in both fresh and processed forms (Baskoro and Mustaruddin 2019). High market demand for squids as an export commodity of Indonesia has made squids become one of the major catches besides fish and lobster (Wulandari 2018). This trend will urge fishermen to conduct intensive fishing while the squid production is still dependent upon the stock in the wild (Tresnati et al. 2012). Squids also grow fast with high feeding rates and conversion efficiencies (Arkhipkin et al. 2015). These biological features make them be ecological opportunists in which they can quickly exploit favorable environmental conditions, but their abundance responds rapidly to poor conditions so that recruitment and abundance may be highly variable on annual time scales (Rodhouse et al. 2014). Therefore, high dependence on wild stock can lead to stock overfishing.

In Indonesia, squid production is far below the world market demand despite its increasing squid production (Triharyuni and Puspasari 2012; Hulalata et al. 2013). This condition is caused by an uneven distribution of squid fisheries across the country. Only several regions do the squid fisheries, and the others do on a very low scale for local consumption.

Information on this group is, so far, mostly related to commercial promotion for fisheries production. This group has been introduced together with other cephalopod groups as an export commodity. It is evident that there is a major artisanal and small-scale inshore element to the world squid fishing fleet and that large volumes of loliginid squids caught in tropical and subtropical regions have high species diversity. Thus, a taxonomic study on this group in Indonesian waters is needed. The taxonomy of the squid fauna generally is poorly understood. Correct species identification is the basis of ecological studies (Veijalainen 2011).

There are 290 species of squids, and about 30-40 species have substantial commercial importance (Arkhipkin et al. 2015) belonging to the family Ommastrephidae, Loliginidae, Onychoteuthidae, and Gonatidae. Triharyuni and Puspasari (2012) have grouped family Loliginidae from Indonesian waters into several genera, *Afrololigo*, *Alloteuthis*, *Doryteuthis*, *Heterololigo*, *Loliolus*, *Lollinguncula*, *Pickfordiateuthis*, *Sepioteuthis*, and *Uroteuthis*, but there are much more species to be described. In Rembang waters, Central Java, four species were found, *Loligo chinensis*, *L. singhalensis*, *L. edulis*, and *L. duvauceli*, in which *L. duvauceli* is the most often caught species (Triharyuni and Puspasari 2012).

In North Sulawesi, Indonesia, there are no squid fisheries and the squid catches are bycatch of other fisheries, but squid production of different species is available every month. Squid catches are obtained at a certain moon period, especially during an early new moon, when the large squid schools come near the surface. The local fishermen catch them using small mesh-sized seine, lift net or using scoop net. A small group of fishermen along Manado Bay, North Sulawesi, Indonesia, have benefitted deep sea squid as an income source and food by relying on hand line fishing or jig fishing as practiced by international squid fisheries (Sundaram and Sawant 2014). Nevertheless, very few fishermen do squid fishing, the amount of catches is very low. This group has become a very important resource so that its sustainability needs to be maintained, and thus, basic information on this resource needs to be provided. This study is intended to identify the deep-sea squid caught by local fishermen of Manado Bay in Sulawesi waters. The findings may enrich the inventory of economically valuable squid species and develop squid fisheries in this area for future regional economic growth.

#### 40 MATERIALS AND METHODS

Squid samples were obtained from fishermen's catch in the Sulawesi Sea, North Sulawesi, Indonesia, about 5 miles from the shore of Manado Bay. The fishing operations were carried out at the geographic position covered by the area formed in the east border (1°30'45"N and 124°42'49"E to 1°32'47"N and 124°42'59"E) and the west one (1°29'30"N and 124°40'48"E to 1°33'45"N and 124°40'50"E) (Figure 1).

Ten fishing trips were done in May to June 2020 using a traditional outboard-motored outrigger boat and fishing activities relied on hand-line facilitated with flashlight artificial bait that was lowered down to 20-25 m depth to get the squid to bite.

44 A total of 600 squid samples were collected and measured in order to describe the species physical characteristics through comparison of mantle size and other morphometric characteristics, such as head length, tentacle length, arm length, fin length, and fin width. For DNA identification, a piece of the squid arm was taken and preserved in 95% ethanol solution before extraction to wash the 35 sample from saltwater and draw water from the cell. All samples were preserved in 95% ethanol and stored at room temperature before DNA extraction.

#### Extraction, PCR, and sequencing

Genome DNA extraction of all samples used Innu PREP DNA Micro Kit (Analytic Jena). The COI gene was amplified applying universal primer pairs LCO1490: 5'-ggtaacaacaatcataaagatattg-3' and HCO2198: 5'taacctcagggtgaccaaaaaatca-3' (Folmer et al. 1994). Polymerase chain reaction (PCR) was carried out in 35 cycles at 95°C (30 sec), 50°C (30 sec), 72°C (50 sec). The PCR product was visualized in 1% (b/v) agarose gel electrophoresis. Bi-directional sequencing was done by First Base CO (Malaysia) using Big Dye® terminator chemistry (PerkinElmer).

#### Data analysis

2 The specimen chromatogram was edited using Mega X v10.1 software (Kumar et al. 2018). The sequences were then compared with Gen Bank data using BLAST (BasicLocal Alignment Search Tools) method (Altschul et al. 1997) and BOLD Systems (Ratnasingham and Hebert 2007). In this study, the sample sequence was compared with that of 7 specimens from South China Sea, one from Mid-Pacific, and one from Eastern Pacific. Besides, another specimen of the same genus *S. pteropus* from Spain, was also used as an 37 outgroup in order to strengthen the comparison. The phylogenetic tree was built using Neighbor-Joining Method of MEGA X and the heterogeneity index was also estimated (Kumar et al. 2018).




Figure 1. Sampling location. Red lines point out the sampling area in Manado Bay, North Sulawesi, Indonesia

## RESULTS AND DISCUSSION

### DNA characteristics

Based on molecular identification using the Internal Transcribed Spacer (ITS) region, the DNA bands obtained were around 500-750bp, the success of PCR was detected by the presence of a single DNA band around 680 bp, the PCR results can be seen in Figure 2.

The sample was identified as purpleback flying squid *Sthenoteuthis oualaniensis*, as shown in the BLAST results in the Gen Bank of NCBI. This species, according to the WoRMs database, has synonymized names as *Loligo brevitentaculata* Quoy & Gaimard, 1832 (synonym), *L. oualaniensis* Lesson, 1830 (original combination), *L. vanikoriensis* Quoy & Gaimard, 1832 (synonym), and *Symplectoteuthis oualaniensis* (Lesson, 1830). This species belongs to class Cephalopod, subclass Coleoidea, suborder Decapodiformes, order Oegopsida, superfamily Ommastrephidae, family Ommastrephidae, subfamily Ommastrephinae, and genus *Sthenoteuthis* (Jerep and Roper 2010). This finding is the first report on this species from the Sulawesi Sea. To confirm the species status, the sample sequence was then compared to 9 *Sthenoteuthis oualaniensis* specimens from the South China Sea, Eastern Pacific, and Central Pacific. The present study shows that *S. oualaniensis* specimen from the Sulawesi Sea has a similarity rate of 99.3% to NCBI record (acc. no. MN101944.1) from South China and 96.4-96.8% to 8 other NCBI records from the South China Sea, Eastern Pacific, and Mid Pacific (Table 1). Specimen of the same genus *Sthenoteuthis pteropus* from Spain used as outgroup also clearly indicates great difference from the specimen and supports the kinship status of the specimen on the study.

The difference is also demonstrated in specimen grouping (Figure 3) as well in which there are 3 different groups formed to indicate the kinship. Group 1 comprises 5 specimens from the South China Sea, one from Mid Pacific, and one from Eastern Pacific. Group 2 comprises specimen from the Sulawesi Sea and 1 NCBI China specimen indicating the species on the study has the closest similarity to one of the South China Sea specimens, while

group 3 is an outgroup that is taken from different species of Spain specimen, *Sthenoteuthis pteropus*. These grouping reflects the kinship of compared *S. oualaniensis* specimens from different areas.

This evidence is supported by the closest kinship to the deep sea squid *S. oualaniensis* specimens from South China waters of NCBI record and they could originate from the same ancestor. Besides, this comparison also indicates that there is intraspecific genetic diversity in *S. oualaniensis* (Table 1 and Figure 3). Genetic diversity could result from population size, in which the larger the population size within species is, the higher genetic diversity will be. This condition is also supported by Hague and Routmant (2016). It could also be caused by different environmental conditions with localities that restrict the organism distribution due to its tolerance limit to the environment.




Figure 1. PCR product of the sample specimen

Table 1. Similarity rate of *Sthenoteuthis oualaniensis* from Sulawesi Sea and those from the Gen Bank DNA sequence database

Sequence	1	2	3	4	5	6	7	8	9	10
MN101944.1_ <i>S. oualaniensis</i> (South_China Sea)										
MW542205.1_ <i>S. oualaniensis</i> (South China Sea)	0.029									
EU660577.1_ <i>S. oualaniensis</i> (Eastern_Pacific)	0.029	0.000								
MF411106.1_ <i>S. oualaniensis</i> (South_China_Sea)	0.031	0.005	0.005							
MF411105.1_ <i>S. oualaniensis</i> (South China Sea)	0.031	0.005	0.005	0.003						
MF411100.1_ <i>S. oualaniensis</i> (South China Sea)	0.031	0.005	0.005	0.003	0.003					
MF411094.1_ <i>S. oualaniensis</i> (South China Sea)	0.031	0.007	0.007	0.005	0.005	0.005				
MF411108.1_ <i>S. oualaniensis</i> (South China Sea)	0.032	0.007	0.007	0.005	0.005	0.005	0.007			
DQ885836.1_ <i>S. oualaniensis</i> (Mid Pacific)	0.029	0.003	0.003	0.002	0.002	0.002	0.003	0.003		
BP_ <i>S. oualaniensis</i> (Indonesia- North_Sulawesi)	0.007	0.032	0.032	0.034	0.034	0.034	0.034	0.036	0.032	
MF980608.1_ <i>Sthenoteuthis pteropus</i> (Spain)	0.119	0.114	0.114	0.116	0.116	0.112	0.116	0.114	0.114	0.116



Figure 3. Phylogenetic tree of *Sthenoteuthis oualaniensis*




Figure 4. Tentacular club (left) and Squid *Sthenoteuthis oualaniensis*

#### Morphological description

*Sthenoteuthis oualaniensis* specimen from Sulawesi Sea has the following morphometric characteristics: Head length is 0.28 of mantle length (ML), tentacle length is 1.158 ML, fin width is 0.3 ML, and fin length is 0.4 ML.

The fourth arm is the shortest and the second arm is the longest among the squid arms (Figure 4). These morphometric characteristics could become typical characteristics in *S. oualaniensis* from the Sulawesi 38 a. Hence, the present study is in agreement with Staaf et al. (2010) and Xua et al. (2020), who found the existence of a morphologically distinct form that is highly genetically divergent in *S. oualaniensis* that separates the dwarf and medium-sized species and exhibits a distinct biogeographic break at equatorial waters of Eastern Pacific Oceans.

*Sthenoteuthis oualaniensis* is known as purpleback flying squid, a tropical Indo-Pacific species that occurs in the Pacific from southern Japan to southern Queensland and from just south of Baja California to northern C 30. *S. oualaniensis* belongs to family Ommastrephidae widely distributed in the tropical and subtropical areas of the Pacific Ocean and the Indian Oce 50 and this species is important for resource exploitation in the South China Sea as one of the major target species of large-scale light falling-net fishing (Yu et al. 2019; Zhao et al. 2021). This species is known to do diurnal vertical migration between the surface at night to the deeper layer during the day, and thus, the species is caught by local fishermen of Manado Bay using the artificial flashlight bait near the surface water column. Fishing operations used a traditional outboard-motored outrigger boat and fishing activities relied on hand-line facilitated with flashlight artificial bait that was lowered down to 20-25 m depth to get the squid to bite.

Cephalopods are sensitive to water temperature (Li et al. 2020) that becomes one of the major environmental factors affecting squid activities, including aggregation, breeding, and emigration (Klemas 2013). Deep sea squids *S. oualaniensis* are distributed from 0-4500 M depth with a peak at 0-500 M depth, water temperature of 25-32°C, and salinity of 34-35 PSU (Jersep and Roper 2010). These seawater temperature ranges cover that of Sulawesi Sea, which enables to support of the occurrence of this squid species based on 10 years of Sulawesi Sea water temperature data (Seawater temperature Sulawesi today Indonesia (seawater.info)). Besides sea surface temperature (SST), sea surface height anomaly (SSHA) at -0.05-0.05 m and chlorophyll-a concentration higher than 0.18 µg/L are required to gain higher catch (Zhao et al. 2021).

For this, *S. oualaniensis* can move at high speeds through the water, easily manoeuvre and quickly respond to their environmental changes. Under distress of external factors, such as predation, this species can reach a high speed and glide above the surface over ten meters. They also found that the cruising speed of an adult squid is between 3 to 10 km per hour, their burst speed achieves greater speeds of up to 35 km per hour, and it can be important for sudden changes in movement and escape behavior. They can occur into shoals from 2 individuals to 800 individuals, and when geographical distributions overlap, they can exist with other members of similar-sized family Ommastrephidae schools, *Dosidicus gigas* and *Ommastrephes bartramii* that are distributed from the Indo-Pacific to Indian Ocean as well (Liu et al. 2016). Ommastrephidae squids are known as voracious and adaptable predators of a broad range of prey, including small crustaceans and fishes at early life stages and shift to micronekton, larger fishes, and cephalopods (including cannibalism) (Gong et al. 2020). These species are highly migratory undertake diel vertical migrations of several hundred meters and seasonal migrations between the shelf and open ocean (Stewart et al. 2013). Thus, they can work as important linkages between both neritic and oceanic food webs (Arkhipkin 2013; Alegre et al. 2014).

In North Sulawesi, this squid species fishing is done by artisanal fishermen using jigging with artificial flashing bait, and the fishing operation has been done in the Sulawesi Sea. In the present study, the squids *S. oualaniensis* caught in the Sulawesi Sea had a size range of 68 mm to 119 mm ML, with a mean mantle length of 89.833±0.9000 mm, reflecting that the squid catches are dominated by large individuals. According to the classification of Nesis (1993), this size range belongs to dwarf form with body size of 90-100 mm ML for mature males and 90-120 mm ML in mature females, and occurs in the equatorial waters and lacks of dorsal mantle photophore patch. The occurrence of *S. oualaniensis* in Sulawesi waters could result from the influence of environmental factors. Similar finding is also reported that changes in marine environments affect cephalopod fishery resources at different spatial and temporal scales (Zhang et al. 2012). Furthermore, the distribution of purpleback flying squid *S. oualaniensis* in this area could not be separated from the impact of the Kuroshio from the Pacific Northern

Equatorial current toward the east coast of Luzon, Philippines that forms the southward-flowing Mindanao Current (Qiu and Lukas 1996) and transfers the upper ocean waters from the Pacific to the Indian Ocean through the Indonesian Seas (Taufiqurrahman et al. 2020). As the water mass enters the Indonesian Seas, the warm and relatively salty characteristics of the Pacific water tend to disappear due to strong vertical tidal mixing. The mixing is believed to affect the carbon, oxygen, and nutrient (nitrate, phosphates) concentrations in the eastern Indonesian Seas and subsequently in the Indian Ocean (Ayers et al. 2014). This event makes Indonesian marine waters be fertile to support the high occurrence of marine animals in the area. The Indonesian Throughflow is the only ocean connector pathway in the equator (Sprintall et al. 2014), and it has an important purpose in the transport of mass and heat from the Pacific into the Indian Ocean (Feng et al. 2018). The throughflow brings the eggs and larvae, along with the rich detritus of the sea that is swept up from the deep offshore basins. It is in agreement with Dell et al. (2011) that the biophysical environment plays an essential role in controlling the distribution and abundance of pelagic predators in the ocean. Cheng et al. (2018), who studied the sword tip squid *Uroteuthis edulis* found that complex oceanographic conditions might affect their population in the Southern East China Sea due to seasonal changes in the Kuroshio Current and Mainland China Coastal Current during the northeasterly monsoon and southwesterly monsoon seasons. This finding has reconfirmed the distribution of *S. oualaniensis* reported in previous studies as tropical deep sea squid.

The present study has provided information on the occurrence of one of the important economic squid resources in Indonesian waters, especially the Sulawesi Sea, that can be used for national economic development. As a center of biodiversity, we strongly believe that there are more squid species living in these waters that need to be described so that there are more taxonomic works that need to be done on species richness of squids in Sulawesi Sea waters as well for future development. Besides, other biological studies of the species need to be done as well in order to have a better understanding of this species that future management and conservation efforts could be well prepared.

#### ACKNOWLEDGEMENTS

We would greatly appreciate Sam Ratulangi University, Manado, Indonesia, for providing a small research grant and laboratory facilities. Our high appreciation is also addressed to Isroja Paransa, who has been involved in this study but passed away in the middle of study.

#### REFERENCES

- Alegre, J. P., Ménéard F., Tafur R., Espinoza P., Argüelles J., Machara V., Flores S., Simier M., Bertrand A. 2014. Comprehensive model of jumbo squid *Dosidicus gigas* trophic ecology in the northern Humboldt

- current system. *Plos One* 9: e0085919. DOI: 10.1371/journal.pone.0085919.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search program. *Nucleic Acids Res* 25: 3389-3402. DOI: 10.1093/nar/25.17.3389.
- Arkipkin AI. 2013. Squid as sentinels linking Southwest Atlantic marine ecosystems. *Deep Res Part II Top Stud Oceanogr* 95: 7-20. DOI: 10.1016/j.dsr2.2012.07.003.
- Arkipkin AI, Rodhouse PGK, Pierce GJ, Sauer W, Sakai M, Allcock L, Arguelles J, Bower JR, Castillo G, Ceriola L. 2015. World squid fisheries. *Rev Fish Sci Aquac* 23: 92-252. DOI: 10.1080/23308249.2015.1026226.
- Ayers JM, Strutton PG, Coles VJ, Hood RR, Matear RJ. 2014. Indonesian throughflow nutrient fluxes and their potential impact on Indian Ocean productivity. *Geophys Res Lett* 41 (14): 5060-5067. DOI: 10.1002/2014GL060593.
- Baskoro MS, Mustaruddin. 2019. The integrated development strategic of capture fisheries based on local superior resources: Case Study of Squid Fisheries in South Bangka Regency. *Jurnal Ilmu dan Teknologi Kelautan Tropis* 11 (3): 541-553. DOI: 10.29244/jitkt.v11i3.24978. [Indonesia]
- Cheng HL, Kuo WL, Hsin YH, Kae YW, Yan LW. 2018. Variation in the catch rate and distribution of swordtip squid *Uroteuthis edulis* associated with factors of the oceanic environment in the Southern East China Sea. *Mar Coast Fish Dyn Manag Ecosyst Sci* 10: 452-464. DOI: 10.1002/mcf2.10039.
- Dell J, Wilcox C, Hobda AJ. 2011. Estimation of yellowfin tuna (*Thunnus albacares*) habitat in waters adjacent to Australia's east coast: Making the most of commercial catch data. *Fish Oceanogr* 20: 383-396. DOI: 10.1111/j.1365-2419.2011.00591.x.
- Doubleday ZA, Prowse TAA, Arkipkin A, Pierce GJ, Semmens J, Steer M, Leporati SC, Lourenço S, Quetglas A, Sauer W, Gillanders BM. 2015. Global proliferation of cephalopods. *Curr Biol* 26: 387-407. DOI: 10.1016/j.cub.2016.04.002.
- FAO. 2019. *FAO Yearbook: Fishery and Aquaculture Statistics 2017*. Food and Agriculture Organization of the United Nations, Rome.
- Feng M, Zhang N, Liu Q, Wijffels S. 2018. The Indonesian throughflow, its variability and centennial change. *Geosci Lett* 5 (1): 3. DOI: 10.1186/s40562-018-0102-2.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* 3 (5): 294-299.
- Gong Y, Li Y, Chen X, Yu W. 2020. Trophic niche and diversity of a pelagic squid (*Dosidicus gigas*): A comparative study using stable isotope, fatty acid, and feeding apparatuses morphology. *Front Mar Sci* 7: 642. DOI: 10.3389/fmars.2020.00642.
- Hague MTJ, Routman EJ. 2016. Does population size affect genetic diversity? A test with sympatric lizard species. *Heredity* 116: 92-98. DOI: 10.1038/hdy.2015.76.
- Hulalata A, Makapedua DM, Papararang RW. 2013. Study on dry salted squid (*Loligo* sp.) processing at different water content and consumer's preference level. *Jurnal Media Teknologi Hasil Perikanan* 11 (1): 26-33. [Indonesia]
- Jerpe P, Roper CF. 2010. *Cephalopods of the World. An Annotated and Illustrated Catalogue of Cephalopod Species Known to date, Vol. 2. Myopsid and Oegopsid*. FAO, Rome.
- Klemas V. 2013. Fisheries applications of remote sensing: An overview. *Fish Res* 148: 124-136. DOI: 10.1016/j.fishres.2012.02.027.
- Kumar S, Stecher G, Li M, Nknyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Mol Biol Evol* 35: 1547-1549. DOI: 10.1093/molbev/msy096.
- Li JJ, Wang JT, Chen XJ, Lei L, Guan CT. 2020. Spatio-temporal variation of *Ommastrephes bartramii* resources (winter & spring groups) in Northwest Pacific under different climate modes. *South China Fish Sci* 16: 62-70. DOI: 10.12131/20190175.
- Liu BL, Chen XJ, Li JH, Chen Y. 2016. Age, growth and maturation of *Sthenoteuthis oualaniensis* in the eastern tropical Pacific Ocean by statolith analysis. *Mar Freshw Res* 67 (12): 1973-1981. DOI: 10.1071/MF14427.
- Nesis KN. 1993. Population structure of oceanic ommastrephids, with particular reference to *Sthenoteuthis oualaniensis*: A review. In: Okutani T, O'Dor RK, Kubodera T (eds). *Recent Advances in Fisheries Biology*. Tokai Univ Press, Tokyo.
- Qiu B, Lukas R. 1996. Seasonal and interannual variability of the North Equatorial Current, the Mindanao Current, and the Kuroshio along the Pacific western boundary. *J Geophys Res Oceans* 101 (C5): 12315-12330. DOI: 10.1029/95JC01515.
- Ratasingham S, Hebert PDN. 2007. BOLD: The barcode of life data system. *Mol Ecol Notes* 7: 355-364. DOI: 10.1111/j.1471-8286.2007.01678.x.
- Rodhouse PGK. 2013. Role of squid in the Southern Ocean pelagic ecosystem and the possible consequences of climate change. *Deep Res Part II: Top Stud Oceanogr* 95: 129-138. DOI: 10.1016/j.dsr2.2012.07.001.
- Rodhouse PGK, Pierce GJ, Nichols S, Sauer WHH, Arkipkin AI, Laptikhovsky VV, Lipinski MR, Ramos J, Gras M, Kidokoro H, Sadayasu K, Pereira J, Lefkaditou E, Pita C, Gasalla M, Haimovici S, Sakai M, Downey N. 2014. Environmental effects on cephalopod population dynamics: Applications for management of fisheries. *Adv Mar Biol* 67: 99-223. DOI: 10.1016/B978-0-12-800287-2.00002-0.
- Sprattall J, Gordon AL, Koch-Larrouy A, Lee T, Potemra JT, Pujiana K, Wijffels SE. 2014. The Indonesian seas and their role in the coupled ocean-climate system. *Nat Geosci* 7 (7): 487-492. DOI: 10.1038/ngeo2188.
- Stauf DJ, Ruiz-Cooley RI, Elliger C, Lebaric Z, Campos B, Markaida U, Gilly WF. 2010. Ommastrephid squids *Sthenoteuthis oualaniensis* and *Dosidicus gigas* in the eastern Pacific show convergent biogeographic breaks but contrasting population structures. *Mar Ecol Prog Ser* 418: 165-178. DOI: 10.3354/meps08829.
- Stewart JS, Gilly WF, Field JC, Payne JC. 2013. Onshore offshore movement of jumbo squid (*Dosidicus gigas*) on the continental shelf. *Deep Res Part II: Top Stud Oceanogr* 95: 193-196. DOI: 10.1016/j.dsr2.2012.08.019.
- Sundaram S, Sawant D. 2014. Large scale exploitation of Indian squid, *Loligo duvauceli* by jigging from nearshore waters of Ratnagiri, Maharashtra. *Mar Fish Infor Sci* 22 (1) & E Ser 2014: 221.
- Taufiqurrahman E, Wahyudi AJ, Masumoto Y. 2020. The Indonesian throughflow and its Impact on Biogeochemistry in the Indonesian Seas. *ASEAN J Sci Technol Dev* 37 (1): 29-35. DOI: 10.29037/ajstd.596.
- Tresnati J, Mallawa A, Nuraeni L, Rapi NL. 2012. Size structure, age groups and growth of squid *Loligo bealei* in the waters of Barru Regency, South Sulawesi. *The Proceedings of 2nd Annual International Conference Syiah Kuala University 2012 & 8th IMT-GT Uninet Biosciences Conference*. Banda Aceh, 22-24 November 2012.
- Triharyuni S, Puspasari R. 2012. Production and fishery season of squids (*Loligo* spp.) in Rembang waters, Central Java. *Jurnal Penelitian dan Perikanan Indonesia* 18 (2): 77-83. DOI: 10.15578/jppi.18.2.2012.77-83.
- Veijalainen A, Broad GR, Wahlberg N, Longino JT, Sääksjärvi. 2011. DNA barcoding and morphology reveal two common species in one *Pimpla molesta* stat. rev. separated from *P. croceipes* (Hymenoptera, Ichneumonidae). *ZooKeys* 124: 59-70. DOI: 10.3897/zookeys.124.1780.
- Wulandari DA. 2018. Morphology, classification, distribution of family Loliginidae squids. *Oseana* 43 (2): 48-65. [Indonesia]
- Xua L, Liu P, Wanga X, van Damme K, Dua F. 2020. Phylogenetic relationships and cryptic species in the genus *Sthenoteuthis* (Cephalopoda: Ommastrephidae) in the South China. *Mol Phylogenet Evol* 149: 1-10. DOI: 10.1016/j.ympev.2020.106846.
- Yu J, Hu Q, Tang D, Zhao H, Chen P. 2019. Response of *Sthenoteuthis oualaniensis* to marine environmental changes in the northern South China Sea based on satellite and in situ observations. *Plos One* 14 (1): e0211474. DOI: 10.1371/journal.pone.0211474.
- Zhang YW, Wang KY, Lu HJ, Chang KJ. 2012. A study on moon phase effect to the hatching of swordtip squid (*Uroteuthis edulis*). *J Fish Soc Taiwan* 39: 209-222. DOI: 10.29822/JFS.41.212.0001.
- Zhao C, Shen C, Bakun A, Yan Y, Kang B. 2021. Purpleback flying squid *Sthenoteuthis oualaniensis* in the South China Sea: Growth, resources and association with the environment. *Water* 13: 65. DOI: 10.3390/w13010065.

# Deep sea squid in Sulawesi Sea, North Sulawesi Province, Indonesia

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

17%

INTERNET SOURCES

15%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Nan Li, Zhou Fang, Xinjun Chen, Zhiping Feng. "Preliminary study on the migration characteristics of swordtip squid (*Uroteuthis edulis*) based on the trace elements of statolith in the East China Sea", *Regional Studies in Marine Science*, 2021  
Publication 1%
- 2 [nrl.northumbria.ac.uk](http://nrl.northumbria.ac.uk)  
Internet Source 1%
- 3 [123dok.co](http://123dok.co)  
Internet Source 1%
- 4 Rongzhen Shi, Manhong Huang, Jing Wang, Chuhan He, Xiaoguo Ying, Xiaohui Xiong, Xiong Xiong. "Molecular identification of dried squid products sold in China using DNA barcoding and SYBR green real time PCR", *Food Additives & Contaminants: Part A*, 2020  
Publication 1%
- 5 [www.fao.org](http://www.fao.org)  
Internet Source 1%



6	<a href="https://www.geoscienceletters.springeropen.com">geoscienceletters.springeropen.com</a> Internet Source	1 %
7	<a href="http://www.ijlsci.in">www.ijlsci.in</a> Internet Source	1 %
8	<a href="http://par.nsf.gov">par.nsf.gov</a> Internet Source	1 %
9	<a href="https://zenodo.org">zenodo.org</a> Internet Source	1 %
10	Tsan-Yu Chiu, Tai-Sheng Chiu, Chih-Shin Chen. "Movement patterns determine the availability of Argentine shortfin squid <i>Illex argentinus</i> to fisheries", Fisheries Research, 2017 Publication	1 %
11	Yumeng Pang, Chih-Shin Chen, Tomohiko Kawamura, Yoko Iwata. "Environmental Influence on Life-History Traits in Male Squid <i>Uroteuthis Edulis</i> With Alternative Reproductive Tactics", Research Square Platform LLC, 2021 Publication	1 %
12	Peter J. Mumby, Robert A. B. Mason, Karlo Hock. "Reconnecting reef recovery in a world of coral bleaching", Limnology and Oceanography: Methods, 2021 Publication	<1 %

13	Stefanie Keller, Antoni Quetglas, Patricia Puerta, Isabella Bitetto et al. "Environmentally driven synchronies of Mediterranean cephalopod populations", Progress in Oceanography, 2017 Publication	<1 %
14	<a href="http://www.documentation.ird.fr">www.documentation.ird.fr</a> Internet Source	<1 %
15	Allen Rodrigo. "The perils of plenty: what are we going to do with all these genes?", Philosophical Transactions of The Royal Society B Biological Sciences, 12/27/2008 Publication	<1 %
16	Rohani Ambo-Rappe. "The success of seagrass restoration using <i>Enhalus acoroides</i> seeds is correlated with substrate and hydrodynamic conditions", Journal of Environmental Management, 2022 Publication	<1 %
17	<a href="http://documents.mx">documents.mx</a> Internet Source	<1 %
18	<a href="http://ir.canterbury.ac.nz">ir.canterbury.ac.nz</a> Internet Source	<1 %
19	<a href="http://journals.biologists.com">journals.biologists.com</a> Internet Source	<1 %

20 T G Hou, X B Yang, T M Wang, J H Liang, S W Li, Y B Fan. "Locomotor transition: how squid jet from water to air", Bioinspiration & Biomimetics, 2020  
Publication <1 %

---

21 [onlinelibrary.wiley.com](https://onlinelibrary.wiley.com)  
Internet Source <1 %

---

22 M W Suryadarma, A S Atmadipoera, N M N Natih, A Koch-Larrouy. "Seasonal Ekman upwelling in the Southwest Sumba from INDESO Model", IOP Conference Series: Earth and Environmental Science, 2021  
Publication <1 %

---

23 [www.schinafish.cn](http://www.schinafish.cn)  
Internet Source <1 %

---

24 [journal.unhas.ac.id](http://journal.unhas.ac.id)  
Internet Source <1 %

---

25 [pasca.unsrat.ac.id](http://pasca.unsrat.ac.id)  
Internet Source <1 %

---

26 [www.rug.nl](http://www.rug.nl)  
Internet Source <1 %

---

27 [fjfsdata01prod.blob.core.windows.net](http://fjfsdata01prod.blob.core.windows.net)  
Internet Source <1 %

---

28 [cris.cientifica.edu.pe](http://cris.cientifica.edu.pe)  
Internet Source <1 %

---

29

[www.esp.org](http://www.esp.org)

Internet Source

&lt;1 %

30

[www.scxuebao.cn](http://www.scxuebao.cn)

Internet Source

&lt;1 %

31

R. SNYDER. "ASPECTS OF THE BIOLOGY OF THE GIANT FORM OF STHENOTEUTHIS OUALANIENSIS CEPHALOPODA: OMMASTREPHIDAE) FROM THE ARABIAN SEA", *Journal of Molluscan Studies*, 1998

Publication

&lt;1 %

32

Hussam El-Din Mohamed, Ali Fadl, Omar Anas, Youssef Wageeh, Noha ElMasry, Ayman Nabil, Ayman Atia. "MSR-YOLO: Method to Enhance Fish Detection and Tracking in Fish Farms", *Procedia Computer Science*, 2020

Publication

&lt;1 %

33

Warwick H. H. Sauer, Ian G. Gleadall, Nicola Downey-Breedt, Zöe Doubleday et al. "World Octopus Fisheries", *Reviews in Fisheries Science & Aquaculture*, 2019

Publication

&lt;1 %

34

[ejlss.indexedresearch.org](http://ejlss.indexedresearch.org)

Internet Source

&lt;1 %

35

[pdffox.com](http://pdffox.com)

Internet Source

&lt;1 %

36

[portals.iucn.org](http://portals.iucn.org)

Internet Source

<1 %

37

[www.biorxiv.org](http://www.biorxiv.org)

Internet Source

<1 %

38

Kornel Kasperek, Beata Horecka, Andrzej Jakubczak, Brygida Ślaska et al. "Analysis of Genetic Variability in Farmed and Wild Populations of Raccoon Dog (*Nyctereutes Procyonoides*) Using Microsatellite Sequences", *Annals of Animal Science*, 2015

Publication

<1 %

39

[firms.fao.org](http://firms.fao.org)

Internet Source

<1 %

40

Geetha Sasikumar, Kolliyil Sunilkumar Mohamed, Kalappurakkal Gopalan Mini, Kurichithara Kunjumani Sajikumar. " Effect of tropical monsoon on fishery abundance of Indian squid ( ( ) ) ", *Journal of Natural History*, 2018

Publication

<1 %

41

[journal15.magtechjournal.com](http://journal15.magtechjournal.com)

Internet Source

<1 %

42

María Delia Viñas, Leocadio Blanco-Bercial, Ann Bucklin, Hans Verheye, José Guilherme F. Bersano, Sara Ceballos. "Phylogeography of the copepod *Calanoides carinatus* s.l. (Krøyer) reveals cryptic species and delimits C.

<1 %

carinatus s.s. distribution in SW Atlantic Ocean", Journal of Experimental Marine Biology and Ecology, 2015

Publication

---

43

Ming Feng, Ningning Zhang, Qinyan Liu, Susan Wijffels. "The Indonesian throughflow, its variability and centennial change", Geoscience Letters, 2018

Publication

---

<1 %

44

[nagasaki-u.repo.nii.ac.jp](http://nagasaki-u.repo.nii.ac.jp)

Internet Source

---

<1 %

45

Gutknecht, Elodie, Guillaume Reffray, Marion Gehlen, Iis Triyulianti, Dessy Berlianty, and Philippe Gaspar. "Evaluation of an operational ocean model configuration at 1/12° spatial resolution for the Indonesian seas (NEMO2.3/INDO12) – Part 2: Biogeochemistry", Geoscientific Model Development, 2016.

Publication

---

<1 %

46

Nalan Gökoğlu. "Shellfish Processing and Preservation", Springer Science and Business Media LLC, 2021

Publication

---

<1 %

47

Nandini Gokte-Narkhedkar, Sameera Siddiqui, Dipak T. Nagrale, Shailesh P. Gawande, P. K. Chakrabarty#. "Race profiling and phylogenetic analysis of the root knot

<1 %

nematode *Meloidogyne incognita* presages the possible emergence of virulence towards cotton in Central India", Research Square Platform LLC, 2022

Publication

48

[darchive.mblwhoilibrary.org](https://darchive.mblwhoilibrary.org)

Internet Source

<1 %

49

[www.ipcc.ch](http://www.ipcc.ch)

Internet Source

<1 %

50

Bilin Liu, Jingyuan Lin, Chunlei Feng, Jianhua Li, Hang Su. "Estimation of age, growth and maturation of purpleback flying squid, *Sthenoteuthis oualaniensis*, in Bashi Channel, central Pacific Ocean", Journal of Ocean University of China, 2017

Publication

<1 %

51

G. R. Carvalho, S. Creer, Michael J. Allen, F. O. Costa et al. "Chapter 1 Genomics in the Discovery and Monitoring of Marine Biodiversity", Springer Science and Business Media LLC, 2010

Publication

<1 %

52

Siyu Jiang, Fuminori Hashihama, Yukio Masumoto, Hongbin Liu, Hiroshi Ogawa, Hiroaki Saito. "Phytoplankton dynamics as a response to physical events in the oligotrophic Eastern Indian Ocean", Progress in Oceanography, 2022

<1 %

53

Yit Sen Bull, Christopher, and Erik van Sebille.  
"Sources, fate, and pathways of Leeuwin  
Current water in the Indian Ocean and Great  
Australian Bight: A Lagrangian study in an  
eddy-resolving ocean model", Journal of  
Geophysical Research Oceans, 2016.

<1 %

Publication

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off