

# Extraction of uronic acid from Sargassum crassifolium and its feeding effects on the immunity of Lohman chicken eggs

*by Malcky Telleng 2*

---

**Submission date:** 22-Sep-2022 08:10AM (UTC+0700)

**Submission ID:** 1905809537

**File name:** xtraction\_of\_uronic\_acid\_from\_Sargassum\_crassifolium\_and\_its.pdf (303.87K)

**Word count:** 2904

**Character count:** 15214

PAPER • OPEN ACCESS


## Extraction of uronic acid from *Sargassum crassifolium* and its feeding effects on the immunity of Lohman chicken eggs

To cite this article: V G Kereh <sup>30</sup> *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **788** 012047

View the [article online](#) for updates and enhancements.

### You may also like

- <sup>13</sup> - [Statistical determination of the step size of secular motors](#)  
K C Neuman, O A Saleh, T Lionnet et al.
- <sup>34</sup> - [Action of bacteriophage T4 and T7 single-stranded DNA-binding proteins with DNA](#)  
Leila Shokri, Ioulia Rouzina and Mark C Williams
- <sup>9</sup> - [Estimation of percentage body fat by dual-energy x-ray absorptiometry: evaluation by \*in vivo\* human elemental composition](#)  
Z Mian Wang, Steven B Heymsfield, Zhao Chen et al.



**ECS** The Electrochemical Society  
Advancing solid state & electrochemical science & technology

242nd ECS Meeting  
Oct 9 – 13, 2022 • Atlanta, GA, US

Abstract submission deadline: **April 8, 2022**

Connect. Engage. Champion. Empower. Accelerate.

**MOVE SCIENCE FORWARD**

Submit your abstract

This banner features a teal background with white text. On the right, there are two images: the top one shows a group of people in a professional setting, and the bottom one shows a city skyline with a lake in the foreground.

## 2 Extraction of uronic acid from *Sargassum crassifolium* and its feeding effects on the immunity of Lohman chicken eggs

2 V G Kereh, I M Usttu, M Najoan, T F D Lumi and M M Telleng

29 Faculty of Animal Science, Sam Ratulangi University, Jln Kampus Bahu, Manado, 95115

E-mail: veybekereh@yahoo.com

**Abstract.** The use of antibiotics is currently banned because it can cause resistance to pathogenic bacteria and cause residues in products. This study aimed to see the effect of drinking water containing uronic acid extracted from *Sargassum crassifolium* (*S. crassifolium*) on the immunity of Lohman chicken eggs. One hundred and twenty laying hens were divided into 2 groups: (1) chickens given commercial feed containing antibiotics and (2) feed without additional antibiotics. The chickens were randomly assigned to one of the 5 brown seaweed supplementation treatments in drinking water A1=0.0% *S. crassifolium* (control); A2=2.5% *S. crassifolium*; A3=4.2% *S. crassifolium*; A4=7.5% *S. crassifolium*; A5=10.0% *S. crassifolium*. The study used a completely randomized factorial design of 5 treatments, 2 factors, and 3 replications. Each replication consisted of 6 heads of laying hens. There were differences between treatments on titer antibody but were no difference on infection of *Salmonella sp* the Conclusion. The uronic acid extracted from *S. crassifolium* has been able to increase the immunity of Lohman chicken eggs.

### 1. Introduction

*Sargassum crassifolium* (*S. crassifolium*) belongs to the *Phaeophyceae* (brown algae) class, which has true roots, stems, and leaves [1], varied forms, and mostly brown or blonde coloration (these colors do not change even with drying) [2]. Brown seaweed (*S. crassifolium*) contains major components including sugar, sulfate and uronic acid and has been shown to play an antiviral and antibacterial role [3]. *S. crassifolium* contains polysaccharides [2], polyphenols, and carotenoids [4]. Polysaccharides play a role in reducing blood lipid, and cholesterol levels, they aid in digestion, and they have antithrombotic, anticancer, antioxidant [5], antiproliferative (uncontrolled cell division), anti-inflammatory [6], and anticoagulants properties [7]. In particular, many researchers report the availability of various antioxidants in seaweed, such as polysaccharides, dietary fiber, minerals, proteins, amino acids, vitamins, polyphenols, and carotenoids [8].

Seaweed availability is quite abundant and has not been optimally utilized thus, it can be processed into animal feed ingredients. As stated by Anggadiredja et al. and March et al. Seaweed is a natural source of non-starch polysaccharides that contains many crude fibers and its bioactive factors affect the digestive process resulting in changes in the microflora in the caecum and the efficient use of nutrients by laying hens [9,10].

However, microbes such as viruses or bacteria in the air, food, or water have the potential to harm livestock. Bacteria, including the *Salmonella sp.* group, often contaminate chickens from the



surrounding environment to hatching, to growth and post-harvest, to the consumers' hands[11]. In addition to affecting the health of livestock, these bacteria will also affect the safety aspects of meat or egg products that will be consumed by humans. Several attempts have been made to overcome this, such as vaccination, sanitation, or antibiotic use. This effort is useful but has limitations, for example, some of the antibiotic resistance to strains of bacteria[32]. Antibiotics are used to inhibit the development of pathogens [13,14]. However, currently, the use of antibiotics in feed has been restricted because it tends to increase the resistance of pathogen bacteria [15,16], therefore antibiotic alternatives from natural ingredients are needed in feed formulas [15–20] to produce safe, healthy, and competitive meat and egg products [21,22]. The use of *S. crassifolium* with uronic acid content may become an alternative substitute for antibiotics. The use of *S. crassifolium* as a potential seaweed additive in feed ingredients, especially feed additives are still not reported. Therefore this study aimed to examine the effects of uronic acid extracted from *S. crassifolium* as an antibiotic substitute on the immunity of Lohman chicken eggs.

## 2. Material and method

### 2.1. Research Material

This study used 120 Lohman strains aged 22 weeks and brown seaweed (*S. crassifolium*). The commercial feeds used in the study were feed containing antibiotics and feed without additional antibiotics. Supplementation of brown seaweed (*S. crassifolium*) extract was 0, 2.5, 5.0, 7.5, and 10.0% given in drinking water. The nutrient content of the feed is presented in table 1. The cage used as an individual battery system (size 35x36x42 cm) equipped with a feeding area, drinking water, and lights (16L/8D lighting system). The chickens were adapted to the provisional feed for 1 month and drinking water for 1 week before treatment began. The Chicken maintenance was carried out for 3 (three) months.

**Table 1.** Nutrient content of the feed.

	Nutrient composition
Dry matter (%)	93.02
Ash (%)	10.77
Crude protein (%)	18.12
Ether extract (%)	5.63
Crude fiber (%)	6.16
BETN (%)	52.34
Gross energy (Ccal/kg)	37.34
Calcium (%)	5.85
Phosphor (%)	0.71

2.1.1. *Preparation of Seaweed extract.* Seaweed extract was made by mixing 100 grams of dried seaweed with ethanol (90%) (5:1), stirred for 3 hours, and allowed to stand for 24 hours at room temperature and then concentrated at 50°C.

2.1.2. *Feeding Trial.* One hundred and twenty Lohman strains aged 18 weeks were divided into 2 groups: (1) chickens given commercial feed containing antibiotics and (2) chickens given feed without additional antibiotics. The chickens were randomly assigned to one of the 5 brown seaweed supplementation treatments: (0, 2.5, 5.0, 7.5, or 10.%) in the drinking water. Feed and drinking water were given *ad libitum* at 07.00 am and afternoon 17.00 pm.

2.1.3. *Variables observed.* The variables observed in this study included the following: the immunity of laying hens to *Salmonella sp.* Detected by Coagglutination test and antibody titer detected by serological tests.

2.2. *Trial design and data analysis.* An experiment was performed in a 5x2 factorial arrangement using a completely randomized design with 3 replications. Each replication consisted of six laying hens. The first factor was the level of brown seaweed (*S. crassifolium*) in drinking water (A1 = 0.0% *S. crassifolium* (control); A2 = 2.5% *S. crassifolium*; A3 = 5.0% *S. crassifolium*; A4 = 7.5% *S. crassifolium*; A5 = 10.0% *S. crassifolium*). The second factor was the presence or absence of antibiotics in feed (B1 = feed with additional antibiotics, B2 = feed without additional antibiotics).

The data were analyzed using analysis of variance followed by Duncan's multiple range test and the orthogonal polynomial test using the SPSS® 21.0 statistical software program.

### 3. Results and discussion

#### 3.1. Results

The administration of uronic acid extracted from *S. crassifolium* in drinking water as a substitute for antibiotics to immunity of Lohmann chickens detected by Coagglutination test, and serological tests. Level of chicken immunity to *Salmonella sp.* detected by Coagglutination test. The analysis of chicken blood Coagglutination showed that all research chickens treated with the uronic acid level of *S. crassifolium* with the addition of antibiotics or without antibiotics in their feed showed positive reactions to *Salmonella sp.* These results showed that all chickens treated with uronic acid levels of *S. crassifolium* have immunity against *Salmonella sp.*

The contained antibody titer of poultry blood serum was detected by serological tests. The results of statistical analysis on the antibody titer of Lohmann chickens showed that the treatment of uronic acid levels and interaction type of feed have non-significant difference ( $P > 0.05$ ) at the beginning of the study (3<sup>rd</sup> week) but at the end of the study (34<sup>th</sup> week 34) that have significant difference ( $P < 0.05$ ), where the treatment of feed without antibiotics with a level of 10% *S. crassifolium* (B2A5) gave the highest antibody titer compared to other treatments (table 1).

**Table 2.** Effect of uronic acid level on the chicken antibody to *Salmonella sp.*

Factor	3 <sup>rd</sup> week			34 <sup>th</sup> week		
	B1	B2	Average	B1	B2	Average
A1	2.33	0.33	1.33	2.33 <sup>a</sup>	2.33 <sup>a</sup>	2.33
A2	2.67	2.00	2.33	3.67 <sup>b</sup>	2.33 <sup>a</sup>	3.00
A3	3.00	1.67	2.33	2.67 <sup>a</sup>	3.00 <sup>a</sup>	2.84
A4	2.67	2.67	2.67	3.33 <sup>a</sup>	3.00 <sup>a</sup>	3.17
A4	3.00	3.33	3.17	3.00 <sup>a</sup>	4.67 <sup>b</sup>	3.84
Average	2.73	2.00		3.00	3.07	

A1 = 0% Uronic acid (control), A2 = 2.5% Uronic acid, A3 = 5% Uronic acid, A4 = 7.5% Uronic acid, A5 = 10% Uronic acid, B1 = Feed with antibiotic; B2 = Feed without antibiotic. A different superscript in the same row shows a significant difference ( $P < 0.05$ ).

#### 3.2. Discussion

Generally, feed intake (g/head/d) without antibiotics with a level of uronic acid extracted from *S. crassifolium* in drinking water tended to be higher than feed intake with antibiotics. This finding shows that uronic acid extracted from *S. crassifolium* in drinking water can increase feed intake and cause a smooth digestion process for feed. Zhao *et al.* [5] Stated that *S. crassifolium* can facilitate food digestion. This study shows that alginate from the uronic acid extracted from *S. crassifolium* probably played an important role in increasing the feed intake of Lohman chickens. Brownlee *et al.*[23] stated

that alginate is a soluble fiber that is beneficial in reducing blood glucose levels, reducing toxicity levels of the intestinal lumen, reducing microbial colonies that are not beneficial, absorbing toxins in the colon, and changing intestinal microflora. These conditions cause the feed absorption process, and the rate of digestive tract emptying to be faster and cause increased feed intake.

The use of uronic acid extracted from *S. crassifolium* in drinking water tends to show the same effect. All chickens treated with uronic acid levels of *S. crassifolium* have immunity against *Salmonella sp.* This is possible because the polysaccharides found in brown seaweed have immunomodulatory activity. The Immunomodulators have an increasing character of the body's defenses both specifically and non-specifically and non-specific induction of both cellular and humoral defense mechanisms. These polysaccharides are non-starch polysaccharides that are resistant to digestion and hydrolysis by saliva in the mouth, stomach, and small intestine, so that they arrive in the intestines intact and act as immunostimulators [24].

The uronic acid level of *S. crassifolium* in laying hens tended to increase the antibody titer value at the end of this research (table 1). That shows the uronic acid from *S. crassifolium* can be used to increase the antibody of layer hens. The uronic acid from *S. crassifolium* has the potential to inhibit viral replication by forming antibodies. Han and Marasco [25] stated that antibody-mediated immune responses are very important for the body's defense against viral infections. Antibody inhibits viral replication by binding viral proteins, thus inhibiting the replication function.

#### 4. Conclusion

The administration of uronic acid extracted from *S. crassifolium* in drinking water as a substitute for antibiotics has been able to increase the immunity of Lohman chicken eggs.

#### References

- [1] Yenusi, T.N.B. A S dan I W 2014 No TitlStudi komposisi dan potensi antioksidan dari pigmen rumput laut *Turbinaria conoides* yang berasal dari perairan pantai Hamadi Jayapura Papua.e Seminar Nasional Kimia dan Pendidikan Kimia VI : "Pemantapan Riset Kimia dan Asesmen 14 dalam pembelajaran Berbasis pendekatan Sainifik pp 316–25
- [2] Merdekawati W and Susanto A B 2009 Kandungan dan komposisi pigmen rumput laut serta potensinya untuk kesehatan *Squalen Bull. Mar. Fish. Postharvest Biotechnol.* 4 41–7
- [3] Mandal P, Mateu C G, Chattopadhyay K, Pujol C A, Damonte E B and Ray B 2007 Structural features and antiviral activity of sulphated fucans from the brown seaweed *Cystoseira indica Antivir. Chem. Chemother.* 18 153–62
- [4] Wijesinghe W and Jeon Y-J 2012 Enzyme-assistant extraction (EAE) of bioactive components: a useful approach for recovery of industrially important metabolites from seaweeds: a review *Fitoterapia* 83 6–12
- [5] Xue Z, Changhu X, Yuepiao C, Dongfeng W and Fang Y U 2005 The study of antioxidant activities of fucoidan from *Laminaria japonica High Technol. Lett. (English Ed.* 11 91–4
- [6] Shiratori K, Ohgami K, Ilieva I, Jin X-H, Koyama Y, Miyashita K, Yoshida K, Kase S and Ohno S 2005 Effects of fucoxanthin on lipopolysaccharide-induced inflammation in vitro and in vivo *Exp. Eye Res.* 81 422–8
- [7] Chandia N P and Matsuhiro B 2008 Characterization of a fucoidan from *Lessonia vadosa* (Phaeophyta) and its anticoagulant and elicitor properties *Int. J. Biol. Macromol.* 42 235–40
- [8] Burtin P 2003 Nutritional value of seaweeds *Electron. J. Environ. Agric. Food Chem.* 2 498–503
- [9] Anggadiredja J 1996 Screening of marine algae from Warambadi seashore Sumba island of Indonesia antibacterial activity *Phytomedicine* 3 37
- [10] March, W., N. Hamid, T. Liu J L and W L W 2013 Fucoidan from New Zealand *Undaria Pinnatifida*: monthly variations and determination of antioxidant activities *Carbohydr Polym* 95 606–14
- [11] Gantois I, Ducatelle R, Pasmans F, Haesebrouck F, Gast R, Humphrey T J and Van Immerseel F 2009 Mechanisms of egg contamination by *Salmonella Enteritidis FEMS Microbiol. Rev.* 33



- 8 718–38
- [12] Devegowda G, Aravind B I R and Morton M G 1997 Immunosuppression in poultry caused by aflatoxins and its alleviation by *Saccharomyces cerevisiae* (Yea-Sacc 1026), and mannanoligosaccharides (Micosorb) *Biotechnol. Feed Ind. Nottingham Univ. Press. Nottingham, United Kingdom* 11
- [13] Ahmadi F, Rahimi F and others 2011 Factors affecting quality and quantity of egg production in laying hens: a review. *World Appl. Sci. J.* **12** 372–84
- [14] Tete A, Gbeassor M, Decuypere E, Tona K and others 2016 Effects of *Moringa oleifera* leaf on laying rate, egg quality and blood parameters. *Int. J. Poult. Sci.* **15** 277–82
- [15] Abdulhasan S D 2011 Effect of Digestrom® and Poultry Star® on the body performance and immunity status of broiler chickens *Int. J. Poult. Sci.* **17** 385–91
- [16] Santoso, U. Y F and K 2018 The effect of fermented *Sauropus androgynus* plus bay leaf inclusion on the hematologic and lipid profiles of female broiler chicken. *J. Poult. Sci.* **17** 1–8
- [17] Abaza I M 2007 Effect of using fenugreek, chamomile and radish as feed additives on productive performance and digestibility coefficients of laying hens *Poult. Sci.* **27** 199–218
- [18] Windisch W, Schedle K, Plitzner C and Kroismayr A 2008 Use of phytogenic products as feed additive for swine and poultry *J. Anim. Sci.* **86** E140–E148
- [19] Abbas T E 2013 The use of *Moringa oleifera* in poultry diets *Turkish J. Vet. Anim. Sci.* **37** 492–6
- [20] Al-Aqil A-A and others 2016 Effects of adding dietary fenugreek (*Trigonella foenum graecum* L.) powder on productive performance and egg quality of laying hens. *Int. J. Poult. Sci.* **15** 259–63
- [21] Mahfuz S U, Nahar M J, Mo C, Ganfu Z, Zhongjun L and Hui S 2017 Inclusion of probiotic on chicken performance and immunity: A review *Int. J. Poult. Sci.* **16** 328–35
- [22] Voemesse K, Tete A, Nideou D, N'nanle O, Gbeassor M, Decuypere E, and Tona K 2018 Effect of *Moringa oleifera* leaf meal on growth performance and blood parameters of egg type chicken during juvenile growth *Intl. J. Poult. Sci.* **17** 154-9
- [23] Brownlee I A, Allen A, Pearson J P, Dettmar P W, Havler M E, Atherton M R and Onsøyen E 2005 Alginate as a source of dietary fiber *Crit. Rev. Food Sci. Nutr.* **45** 497–510
- [24] Ale M T, Mikkelsen J D and Meyer A S 2011 Important determinants for fucoidan bioactivity: A critical review of structure-function relations and extraction methods for fucose-containing sulfated polysaccharides from brown seaweeds *Mar. Drugs* **9** 2106–30
- [25] Han T M W 2011 Structural basis of influenza virus neutralization *Ann NY Acad Sci.* **1217** 178–90

# Extraction of uronic acid from Sargassum crassifolium and its feeding effects on the immunity of Lohman chicken eggs

## ORIGINALITY REPORT

**20%**  
SIMILARITY INDEX

**18%**  
INTERNET SOURCES

**15%**  
PUBLICATIONS

**%**  
STUDENT PAPERS

## PRIMARY SOURCES

- 1** Muhittin Zengin, Aslıhan Sur, Ziya İlhan, Mehmet Ali Azman, Hakan Tavşanlı, Selim Esen, Oğuz Koray Bacaksız, Ergün Demir. "Effects of fermented distillers grains with solubles, partially replaced with soybean meal, on performance, blood parameters, meat quality, intestinal flora, and immune response in broiler", Research in Veterinary Science, 2022  
Publication **1%**
- 2** [www.sciencegate.app](http://www.sciencegate.app)  
Internet Source **1%**
- 3** [ebin.pub](http://ebin.pub)  
Internet Source **1%**
- 4** [openaccess.ahievran.edu.tr](http://openaccess.ahievran.edu.tr)  
Internet Source **1%**
- 5** [www.ksast.org](http://www.ksast.org)  
Internet Source **1%**
- 6** [ajouronline.com](http://ajouronline.com)  
Internet Source **1%**



1 %

7

[www.pjbs.org](http://www.pjbs.org)

Internet Source

1 %

8

B. Sundu, U. Hatta, A.S. Chaudhry. "Potential use of beta-mannan from copra meal as a feed additive for broilers", World's Poultry Science Journal, 2019

Publication

1 %

9

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

1 %

10

Geena Mariya Jose, Mahadevan Raghavankutty, G. Muraleedhara Kurup. "Sulfated polysaccharides from Padina tetrastromatica induce apoptosis in HeLa cells through ROS triggered mitochondrial pathway", Process Biochemistry, 2018

Publication

1 %

11

Tang, Shirley Gee Hoon, Chin Chin Sieo, Ramasamy Kalavathy, Wan Zuhainis Saad, Su Ting Yong, Hee Kum Wong, and Yin Wan Ho. "Chemical Compositions of Egg Yolks and Egg Quality of Laying Hens Fed Prebiotic, Probiotic, and Synbiotic Diets : Chemical compositions and egg quality...", Journal of Food Science, 2015.

Publication

1 %

12	<a href="http://medpub.litbang.pertanian.go.id">medpub.litbang.pertanian.go.id</a> Internet Source	1 %
13	<a href="http://www.citeulike.org">www.citeulike.org</a> Internet Source	1 %
14	<a href="http://etd.repository.ugm.ac.id">etd.repository.ugm.ac.id</a> Internet Source	1 %
15	<a href="http://journals.ums.ac.id">journals.ums.ac.id</a> Internet Source	1 %
16	<a href="http://snkpk.fkip.uns.ac.id">snkpk.fkip.uns.ac.id</a> Internet Source	1 %
17	Pádraigín A. Harnedy. "BIOACTIVE PROTEINS, PEPTIDES, AND AMINO ACIDS FROM MACROALGAE1 : MACROALGAE: BIOACTIVE AGENT SOURCE", <i>Journal of Phycology</i> , 04/2011 Publication	1 %
18	<a href="http://link.springer.com">link.springer.com</a> Internet Source	1 %
19	Rangsun Charoensoo, Nithat Wichasit, Thitima Pechrkong, Tossaporn Incharoen, Sonthaya Numthuam. "STAT5B Gene Polymorphisms are Associated with Egg Production and Egg Quality Traits in Laying Hens", <i>Asian Journal of Animal and Veterinary Advances</i> , 2016 Publication	1 %

[china.iopscience.iop.org](http://china.iopscience.iop.org)

20	Internet Source	1 %
21	coek.info Internet Source	1 %
22	garuda.kemdikbud.go.id Internet Source	1 %
23	livrepository.liverpool.ac.uk Internet Source	<1 %
24	onlinelibrary.wiley.com Internet Source	<1 %
25	pdffox.com Internet Source	<1 %
26	M. Jimoh, O. A. Enikuomihin, C. G. Afolabi, V. I. O. Olowe, A. Egbontan. " Achieving improved control of foliar diseases of sesame ( L.) intercropped with maize ( L.) through foliar spray of plant extracts ", Archives of Phytopathology and Plant Protection, 2016 Publication	<1 %
27	autodocbox.com Internet Source	<1 %
28	www.ifa-tulln.boku.ac.at Internet Source	<1 %
29	Yamaguchi, Michitaka, Mitsue Miyazaki, Matthew P. Kodrasov, Henki Rotinsulu, Fitje	<1 %

Losung, Remy E.P. Mangindaan, Nicole J. de Voogd, Hideyoshi Yokosawa, Benjamin Nicholson, and Sachiko Tsukamoto.

"Spongiacidin C, a pyrrole alkaloid from the marine sponge *Stylissa massa*, functions as a USP7 inhibitor", *Bioorganic & Medicinal Chemistry Letters*, 2013.

Publication

---

30

[jptcp.com](http://jptcp.com)

Internet Source

<1 %

---

31

Jalal Bayati Zadeh, Nasroallah Moradi kor, Ali Olfati. " The Effects of Different Levels Aqueous Extract of Fenugreek ( L.) on Performance and Immune Response of Laying Hens ", *Journal of Essential Oil Bearing Plants*, 2015

Publication

<1 %

---

32

[bioone.org](http://bioone.org)

Internet Source

<1 %

---

33

[vetjournal.ankara.edu.tr](http://vetjournal.ankara.edu.tr)

Internet Source

<1 %

---

34

Mark C. Williams, Ioulia Rouzina, Victor A. Bloomfield. "Thermodynamics of DNA Interactions from Single Molecule Stretching Experiments", *Accounts of Chemical Research*, 2002

Publication

---

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off