

PROTEINS PROFIL OF SAUSAGE LAYING CHICKEN MEAT WITH ANGKAK (RED RICE) USED AS NATURAL FOOD MATERIAL

by Tiltje Ransalele 8

Submission date: 26-Aug-2022 10:12AM (UTC+0700)

Submission ID: 1887234971

File name: PROTEINS_PROFIL_OF_SAUSAGE_LAYING_CHICKEN_MEAT_WITH.pdf (816.36K)

Word count: 4809


Character count: 28635

1

SCIENTIFIC PAPERS

SERIES D. ANIMAL SCIENCE

Volume LXII, No. 2, 2019



University of Agronomic Sciences and
Veterinary Medicine of Bucharest Faculty of
Animal Productions Engineering and
Management

1
SCIENTIFIC PAPERS

SERIES D

ANIMAL SCIENCE

Volume LXII, No. 2

2019

Bucharest

3

SCIENTIFIC COMMITTEE

- Stelian ACATINCĂI - Banat University of Agricultural Sciences and Veterinary Medicine „King Michael I of Romania“, from Timișoara, Romania
- Vasco A.P. CADAVEZ - Departamento de Ciência Animal & Centro de Investigação de Montanha (CIMO), Escola Superior Agrária, Instituto Politécnico de Bragança - Portugal
- Mioara COSTACHE - Fish Culture Research and Development Station of Nucet, Romania
- Georgeta DINIȚĂ - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Cătălin DRAGOMIR - National Research and Development Institute for Animal Biology and Nutrition Balotești, Romania
- Nicolae EREMIA - State Agrarian University of Moldova, Chisinau, Republic of Moldova
- Horia GROSU - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Khalid Hamid HASSAN - University of Diyala, Irak
- Armagan HAYIRLI - Ataturk University, Erzurum, Turkey
- Mostafa A.R. IBRAHIM - University of Kafrelsheikh, Egipt
- Ondrej KADLEČÍK - Slovak Agricultural University Nitra, Slovakia
- Yusuf KONCA - Erciyes University, Kayseri, Turkey
- Monica Paula MARIN - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Carmen Georgeta NICOLAE - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Mircea POP - University of Agricultural Sciences and Veterinary Medicine „Ion Ionescu de la Brad“ of Iasi, Romania
- Agatha POPESCU - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Elena POPESCU-MICLOȘANU - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Ștefan POPESCU-VIFOR - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Sri Bandiati Komar PRAJOG - Padjadjaran University Bandung, Indonesia
- Paul Rodian TĂPĂLOAGĂ - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Ilie VAN - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania
- Livia VIDU - University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania

EDITORIAL BOARD

General Editor: Prof. Ph.D. Gheorghe Emil MĂRGINEAN
Executive Editor: Assoc. Prof. Ph.D. Monica Paula MARIN

PUBLISHERS:

University of Agronomic Sciences and Veterinary Medicine of Bucharest, Romania Address: 59 Mărăști Blvd, District 1, 011464, Bucharest, Romania Phone: + 40 213 182 564, Fax: +40 213 182 888, www.zootehnie.ro

CERES Publishing House

Address: 29 Oastei Street, District 1, Bucharest, Romania
Phone: + 40 317 90 23, E-mail: edituraceres@yahoo.com, Webpage: www.editura-ceres.ro

Copyright 2019

To be cited: Scientific Papers. Series D. Animal Science, Volume LXII, No. 2, 2019

*The publishers are not responsible for the opinions published in the Volume.
They represent the authors' point of view.*

ISSN 2285-5750; ISSN CD-ROM 2285-5769; ISSN Online 2393-2260; ISSN-L 2285-5750

International Database Indexing: Web of Science Core Collection (Emerging Sources Citation Index), Index Copernicus, CABI, DOAJ, Ulrich's Periodicals Directory (ProQuest), PBN, Cite Factor (Academic Scientific Journals), Scipio, OCLC (WorldCat), Research Bible, Google Scholar.

SUMMARY

I. SESSION GENETICS AND BREEDING

1. THE COMBINATIVE CAPACITY OF HYBRIDS FROM CROSSBREEDING LINES AT THE SILKWORMS BREED BANEASA 75 (*Bombyx mori* L.) - Georgeta DINIȚĂ, Monica Paula MARIN, Daniel Severus DEZMIREAN, Minodora TUDORACHE, Ioan CUSTURĂ13
2. ASSESSMENT OF THE EXTERIOR OF FIRST-CALF HEIFERS OF HOLSTEIN BREED - Valentin FOKSHA, Alexandra KONSTANDOGLO, Alexander KENDIGELIAN, Igor KIBASH, Vasily KURULYUK19
3. STUDY ON INDICES OF REPRODUCTION AT POPULATION OF THE HOLSTEIN COWS OF DIFFERENT ORIGIN IN THE SOUTH AREA OF REPUBLIC OF MOLDOVA - Vera GRANACI, Valentin FOKSHA, Alexandra KONSTANDOGLO24
4. THE RELATIONSHIP BETWEEN HOLSTEIN COWS EXTERIOR AND DAIRY PRODUCTIVITY BY VARIOUS BREEDING - Alexandra KONSTANDOGLO, Valentin FOKSHA, Vera GRANACI.....29

II. SESSION NUTRITION

3. EFFECT OF USE OF PREBIOTICS^{BLS} (*Bacillus l.*, *Lactobacillus spp.* AND *Saccharomyces s.*) BASED ON RIMP WASTE ON PROTEIN EFFICIENCY RATIO IN INDONESIA LOCAL CHICKEN - Abun ABUN, Denny RUSMANA, Tuti WIDJASTUTI, Kiki HAETAMI37
2. EFFECTIVENESS OF JAPANESE ANTS (*Uromoides dermestoides*) AS ANTI-DIABETIC ON WHITE RATS (*Rattus norvegicus*) - Jeanne Renny AGU, Max TULUNG, Christin SALAKI, Jusuf MANUEKE44
3. THE INFLUENCE OF OREGANO ESSENTIAL OIL ON EGG QUALITY AND EGG SHELL CONTAMINATION OF LAYING HENS KEPT IN FURNISHED CAGES - Muzaffer DENLI, Aydin VURAL, Simten YESILMEN ALP48
4. PROTEOLYTIC POTENTIAL OF *Bacillus sp.* FROM FISH GUT AND NUTRIENT CONTENT OF SUBSTRATE - Kiki HAETAMI, YuniarMULYANI, YeniMULYANI, Abun ABUN, Junianto UNIANTO53
7. PHYTOCHEMICAL POTENCY AND ANTIMICROBIAL ACTIVITY OF ARECA VESTIARIA GISEKE AS A CANDIDATE FEED ADDITIVES IN BROILER - Jola Josephien Mariane Roosje LONDOK, Sumiati SUMIATI, John Ernst Gustaaf ROMPIS, Jet Saartje MANDEY 60
6. BENEFICIAL USES OF CHROMIUM IN LAYING HENS NUTRITION: A REVIEW - Elena Narcisa POGURSCHI, Carmen Georgeta NICOLAE, Corina Aurelia ZUGRAVU, Silian VLAD, Marius MAFTEI, Monica Paula MARIN, Melania Florina MUNTEANU65
5. ANTIBACTERIAL ACTIVITY OF NONI JUICE FRUIT (*Morinda citrifolia* L) ON PERFORMANCE AND HEMATOLOGIC INDICATOR ON SENTUL CHICKEN - Tuti WIDJASTUTI, Iwan SETIAWAN, Abun HASBUNA, Indrawati YUDAASMARA71
8. RESEARCH ON THE INFLUENCE OF SOME PROBIOTICS ON THE PRODUCTION PERFORMANCE OF BROILER CHICKENS - Daniel RIZEA, Horia GROSU77
9. NUTRITION CONSULTANT BASED ON MACHINE LEARNING FOR PREECLAMPSIA COMPLICATIONS - Iuliana MARIN, Nicolae GOGA82

10. ENRICHING THE DIET IN POLYUNSATURATED FATTY ACIDS FOR LAYING HENS USING FLAXSEED MEAL AND RICE BRAN - **Tatiana Dumitra PANAITE, Margareta OLTEANU, Mariana ROPOTA, Cristina SOICA, Petru Alexandru VLAICU, Petruta VISINESCU, Rodica Diana CRISTE**.....88

III. SESSION REPRODUCTION, PHYSIOLOGY, ANATOMY

1. COMPARATIVE STUDY ON HEMATOLOGICAL AND BIOCHEMICAL CHARACTERIZATION BLOOD PROFILE IN BIRDS GROWN IN FARMS OF ROMANIA - **Mircea LAZĂR, Roxana LAZĂR, Cristina SIMEANU, Paul Corneliu BOIȘTEANU**97
2. STUDY REGARDING SOME REPRODUCTIVE PARAMETERS IN SHAGYA ARABIAN MARES FROM RĂDĂUȚI STUD FARM - **Claudia PĂNZARU, Ioan GÎLCĂ, Marius DOLIȘ, Răzvan RADU-RUSU, Roxana Nicoleta RAȚU, Mihaela IVANCIA**103
3. PRESERVATION OF RAMS' SPERM AT +2-+4°C REGION OF MOLDOVA - **Doina ROTARI** .. 109
4. PROTON TRANSVERSE RELAXATION TIMES OF FREE AND BOUND WATER IN RAT LIVER AND RED BLOOD CELLS FED ON CHOLESTEROL REACH DIET .THE EFFECT OF PROCANE AND ASLAVITAL - **Cristian Romeo REVNIC, Flory REVNIC, Silviu VOINEA**114

IV. SESSION TECHNOLOGIES OF ANIMAL HUSBANDRY

1. THE RELATIONSHIP BETWEEN ABUNDANCE, DIVERSITY WITH COWS SKIN DEFECTS ACCORDING TO DIFFERENT ALTITUDE, HUMIDITY AND TEMPERATURE - **Geetruida J.V. ASSA, Wisje L. TOAR, Meis J. NANGOY, Endang PUDJIHASTUTI, Siane RIMBING, Albert J. PODUNG**123
2. EUROPEAN LEGAL FRAMEWORK IN MANUFACTURING AND PROCESSING OF MILK. THE MILK PACKAGE. ELIMINATION OF MILK QUOTA. CASE STUDY: GERMANY - **Ion Tudor COMAN, Livia VIDU, Constantin TRĂIS TARU, Gheorghe Emil MĂRGINEAN** 128
3. RESEARCHES ABOUT INFLUENCE OF PRO-BIOTICS ON BROILER PRODUCTION PERFORMANCES - **Ioan CUSTURĂ, Minodora TUDORACHE, Ilie VAN, Monica Paula MARIN, Andrei MARMANDIU, Emanuel Sergiu PANĂ**135
4. STUDY ON THE INFLUENCE OF THE FEEDING PROGRAM ON THE PREPARATION OF HEIFERS FOR FIRST INSEMINATION - **Oana (MILITARU) DOROBĂȚ, Gheorghe Emil MĂRGINEAN, Ayman HASSAN, Răzvan POPA, Livia VIDU**140
5. RESEARCH ON THE EVOLUTION OF THE ABERDEEN ANGUS BREED IN ROMANIA - **Ioan Teodor GOCIMAN, Gheorghe Emil MĂRGINEAN, Stelian BĂRĂITĂREANU, Carmen Sorgeta NICOLAE, Livia VIDU**145
6. MAKING AND CHARACTERIZATION OF PEGAGAN DRY EXTRACT (*Centella asiatica*) AS FEED ADDITIVES FOR ANIMAL FEEDING - **Jhondri JHONDRI, Abun ABUN, Kiki HAETAMI**150
7. PIG PERFORMANCES FED WITH COCONUT WATER AND PULP - **Mien LAPIAN, Marie NAJOAN, Anny RAWUNG** 154
8. RESEARCHES REGARDING CANNON BONE PERIMETER AVERAGE PERFORMANCE S IN ROMANIAN HUCUL HORSE BREED – PIETROSU BLOODLINE - **Marius MAFTEL, Gheorghe MĂRGINEAN, Iulian VLAD, Carmen NICOLAE, Dana POPA, Elena POGURSCHI, Lucica NISTOR, Daniela IANITCHI, Serban PURDOIU** 160

| | |
|---|-----|
| 9. THE DYNAMICS OF MILK PRODUCTION IN MONTBELIARDE BREED ON A FARM IN SOUTHERN ROMANIA - Robert MIHAI, Costel MIHALASCU, Gheorghe Emil MĂRGINEAN, Monica Paula MARIN, Mirela Aurora CĂRĂTUȘ, Livia VIDU | 165 |
| 10. STUDY OF HOOF TRIMMING IMPORTANCE FOR TRANSITION PERIOD COW - Ion Silver MILITARU, Laura Florentina VLĂȘCEANU, Alexandru MIHAI, Stelian BĂRĂITĂREANU, Livia VIDU, Gheorghe Emil MĂRGINEAN | 170 |
| 11. INFLUENCE OF HEAVY METALS ON COCOON <i>BOMBYX MORIL</i> . - Tsvetelin KOLOVA | 176 |
| 12. SOME CORRELATIONS BETWEEN ENVIRONMENTAL PARAMETERS AND THE FORAGING BEHAVIOUR OF HONEYBEES (<i>APIS MELLIFERA</i>) ON OIL SEED RAPE (<i>BRASSICA NAPUS</i> L.) - Silvia PĂTRUICĂ, Eliza SIMIZ, Ioan PETI, Lavinia ȘTEF | 180 |
| 13. HOMOLOGATION OF SHEEP BREEDS - EUROPEAN AND NATIONAL LEGISLATION - Andrei PUIE, Ion RĂDUCUȚĂ, Alexandra POLIOPOL, Comel FRUJINĂ, Ion CĂLIN | 185 |
| 14. MONTHLY CHANGES OF BEHAVIORAL CHARACTERISTICS IN HOLSTEIN-FRIESIAN, BROWN SWISS AND SIMMENTAL BULLS - Veysel Ali ÜNAL, Atakan KOÇ | 192 |
| 15. RESEARCH ON THE EVOLUTION OF THE GROWTH PROCESS AT THE TZURCANA SHEEP WITH THE PATERN BREEDS VENDEEN AND WHITE OF CENTRAL MASSIF - Comel FRUJINĂ, Iulian VLAD, Ion CĂLIN, Marius MAFTEL, Ion RĂDUCUȚĂ, Daniela IANIȚHI | 199 |

V. SESSION TECHNOLOGIES OF THE AGRO FOOD PRODUCTS PROCESSING


| | |
|--|-----|
| 1. PHYSICO-CHEMICAL PARAMETERS AND SPECTRAL STRUCTURE (FT-IR) OF HONEY FROM IASI COUNTY (NORTH-EASTERN ROMANIA) - Aida ALBU, Simona-Maria CUCUȘAN, Ioan Mircea POP | 209 |
| 2. AN EVALUATION OF GUELDER ROSE (<i>Viburnum opulus</i> L.) AND HAWTHORN (<i>Crataegus monogyna</i>) CONCENTRATES AS ALTERNATIVE ANTIOXIDANT SOURCES TO BHT AND NITRITE IN POULTRY MEAT MODEL SYSTEM - Burcu ÇEMTEKİN, Emine KILINÇ, Lalehan KARABACAK, Tuğçe DAĞTEKİN, Tuğçe TIRYAKI, Ali SOYUÇOK, Azim ŞİMŞEK, Birol KILIÇ | 217 |
| 3. THE TRACEABILITY OF FOOD PRODUCTS IN RELATION WITH FOOD INTEGRITY – A REVIEW - Florentina MARIN HADDAD, Ioana TOMA, Mona Elena POPA, Mariana PIPIRIGEANU | 228 |
| 4. IDENTIFICATION OF AMINOACID PROFILE AND PROXIMATE COMPOSITION OF THREE TROUT BREEDS REARED IN THE NORTH EASTERN REGION OF ROMANIA - Cătălin Emilian NISTOR, Vasile BĂCILĂ, Iulia VĂRZARU MAROȘ, Gabriel Vasile HOHA, Benone ȘĂRIN | 233 |
| 5. STUDY REFERRING TO THE APPEARANCE OF CONTAMINATION WITH DEOXYNIVALENOL IN GRAINS, GRAIN FLOUR AND BAKERY PRODUCTS ON THE ROMANIAN MARKET - Carmen Daniela PETCU, Ioana Mădălina GEORGESCU, Ovidiu Valentin ZVORIȘTEANU, Cătălin Nicolae NEGREANU | 241 |
| 6. STUDY ON NUTRITIONAL QUALITY OF SMOKED AND MARINATED PRODUCTS FROM HERRING (<i>CLUPEA HARENGUS</i>) - Mircea POP, Gabriela FRUNZĂ, Cădălia POP | 246 |
| 7. FISH MEAT CONTAMINATION WITH HEAVY METALS – A REAL CONCERN FOR THE FOOD CONSUMPTION - Alexandru POPESCU, Geanina VLASE, Carmen Georgeta NICOLAE, Cătălin PĂUN, Magda-Ioana NENCIU | 254 |

8. ON THE OCCURRENCE OF POTASSIUM SORBATE (E202) IN CERTAIN FOOD AND BEVERAGE PRODUCTS - **Cristina Gabriela RADU-RUSU, Ioan Mircea POP, Gabriela FRUNZA, Daniel SIMEANU** 259
9. USAGE OF HISTOLOGICAL AND RHEOLOGICAL TECHNIQUES IN ASSESSMENT AND PREDICTION OF MEAT TEXTURAL PROPERTIES - **Razvan Mihail RADU-RUSU, Marius Giorgi USTUROI, François DJIIE KO UATCHO, Gabriel Vasile HOHA, Ana LEAHU, Mircea OROIAN, SorinaRO PCIUC, Roxana Nicoleta RAȚU, Cădă PĂNZARU** 264
10. PROTEINS PROFIL OF SAUSAGE LAYING CHICKEN MEAT WITH ANGKAK (RED RICE) USED AS NATURAL FOOD MATERIAL - **Delly Bertha Johana RUMONDOR, Rita TINANGON, Jantje PAATH, Endang PUJIHASTUTI, Tiltje RANSALELEH** 270
11. EFFECT OF YEAST AND LACTIC ACID BACTERIA IN CULLED LAYING HENS SALAMI AGAINST *ESCHERICHIA COLI*, *STAPHYLOCOCCUS AUREUS* AND *SALMONELLA SP.* - **Sofi Margritje SEMBOR, Balia ROOSTITA, Hendronoto LENGKEY, Lilis SURYANINGSIH** 275
12. CORRELATIVE RESEARCH REGARDING THE TOTAL POLYPHENOLIC CONTENT, ANTIOXIDANT AND ANTIBACTERIAL ACTIVITY OF THREE TYPES OF ROMANIAN HONEY - **Octavia TAMAS-KRUMPE, Otilia BOBIȘ, Rodica MĂRGĂOAN, Flore CHIRILĂ, Călin LAȚIU, LaurențOGNEAN** 282
13. EVALUATION OF RAW MILK QUALITY GATHERED FROM NORTH EAST AREA OF ROMANIA - **Cătălin Emilian NISTOR, Vasile BĂCILĂ, Petrică AVRAM, Alexandru USTUROI, Bogdan-Vlad AVARVAREI** 289
14. THE INFLUENCE OF AUXILIARY MATERIALS ON HARDNESS, HEAT TREATMENT LOSSES AND SENSORY PROPERTIES OF THE MEAT PRODUCTS - **Daniela IANIȚCHI, Cristina DIACONESCU, Iulian VLAD, Lucica NISTOR, Camelia HODOȘAN, Monica MARIN, Marius MAFTEI** 296
15. SPECIFIC GLUTEN-BASED FLOURS RECOMMENDED IN THE GLUTEN-FREE DIET - **Nela DRAGOMIR, Gratiela-Victoria BAHACIU** 302
16. RESEARCH REGARDING THE USAGE INFLUENCE OF SOYA FLOUR AND FOOD ADDITIVES ON BREAD QUALITY - **Monica MARIN, Georgeta DINIȚĂ, Elena POGURSCHI, Carmen Georgeta NICOLAE** 309

VI. SESSION WILD LIFE MANAGEMENT, FISHERY AND AQUACULTURE

1. EFFECT OF SHELL INJURY ON HAEMOCYTE CONCENTRATION AND SHELL REGROWTH OF GIANT AFRICAN LAND SNAIL (*ARCHACHATINA MARGINATA*) - **John ABIONA, Fatimah DUROSINMI, Yemisi AYO-AJASA, Muhammed ONAGBESAN** 317
2. DETECTION OF *AEROMONAS SALMONICIDA* IN FISH SAMPLES FROM LAKE OHRID BY CULTURE AND POLYMERASE CHAIN REACTION METHODS - **Enkeleda BERBERI, Ariola DEVOLLI, Ilir LLOHA, Kevin MICI** 323
3. INDUCTION AND RECOVERY TIMES OF ANESTHESIA IN CYPRINIDS USING VARIOUS DOSES OF CLOVE OIL (*EUGENIA CARYOPHYLLATA*) - **Daniel COCAN, Florentina POPESCU, Călin LAȚIU, Paul UIUIU, Radu CONSTANTINESCU, Cristian MARTONOS, Aurelia COROIAN, Andrada IHUȚ, Camelia RĂDUCU, Mihai BENȚEA, Vioara MIREȘAN** ... 328
4. PRELIMINARY RESULTS REGARDING THE EFFECTS OF DIETARY-PROTEIN LEVELS ON THE GROWTH PERFORMANCE AND FEED EFFICIENCY OF COMMON CARP fry - **Anca Nicoleta Cordeli (Săvescu), Mirela crețu, Lucian Oprea** 336

TECHNOLOGIES OF THE AGRO FOOD PRODUCTS PROCESSING



PROTEINS PROFIL OF SAUSAGE LAYING CHICKEN MEAT WITH ANGKAK (RED RICE) USED AS NATURAL FOOD MATERIAL

Delly Bertha Johana RUMONDOR, Rita TINANGON, Jantje PAATH,
Endang PUJIHASTUTI, Tiltje RANSALELEH

Sam Ratulangi University, Faculty of Animal Husbandry,
Jalan Kampus Bahu, Manado 95115, Indonesia

Corresponding author email: berthaumondor@gmail.com

Abstract

This study examines the potential of Angkak as a binding compound, color improvement in the process of making sausage chicken layered rejects as a curing material. Testing protein profiles related to dissolved protein, texture, water binding capacity to form a compact and soft texture. The design was carried out in a completely randomized design (CRD) 4 x 4, as a treatment using the Angkak level without Angkak (0%), Angkak 0.5%, Angkak 1% and Angkak 1.5% with replications 4 times followed by a test BNJ. The data obtained were analyzed by analysis of variance (Analysis of Variance) which included chemical, color, total protein, and protein profile determination using SDS-PAGE method (Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis). The results of protein profiles with SDS-PAGE (sodium dodecyl sulphate polyacrylamide gel electrophoresis), where R0 is a sausage without the addition of Angkak, R1: sausages that use 0.5% Angkak concentration, R2: sausages that use a concentration of Angkak 1.0% and R3: sausage who use the Angkak concentration of 1.5%. The conclusion of the study was that the addition of Angkak gave a chemical change and protein profile in the sausages of rejected laying hens and the range of molecular weights found in R0, R1, R2 and R3 were the same, namely 12.44 - 47.89 kDa.

Key words: sausage meat, laying chicken, Angkak, curing.

INTRODUCTION

Reed laying chicken meat as well as other livestock products is livestock commodities that need to be developed and improved. In general, the Indonesian people have known laying chicken meat rejects as a source of food that is mainly expected to be eggs, but the obstacles that have existed during this time rejects laying chicken meat is not much in demand by the public, because laying chicken meat apart from clay meat that is not favored by consumers also variations in processing chicken rejected laying into processed products is very limited, so that a processing technology is pursued to utilize and increase the added value of rejected laying chicken meat. One processing technique that can be developed.

According to Tisnadjaja (2006) that the use of nitrite in the process of curing in meat is 125 ppm, but this use has not been strictly monitored on food products, while the use of Angkak as a substitute for saltpeter or nitrite can be reduced by up to 60% without any apparent changes in organoleptic properties. Angkak pigments are used as a partial

substitute for nitrite in the processing of meat cured like ham and sausage beef, both in improving the red color of meat products and in inhibiting the growth of spore-forming bacteria such as *Bacillus cereus* and *Bacillus stearothermophilus*.

According to Astawan (2012), the use of Angkak can reduce the use of nitrite in food. Nitrite is often used as a component of saltpeter, a substance used to maintain the red color of meat, especially in making sausages, smoked meat and comet. Through its antimicrobial properties, the use of Angkak in making sausages is not only a red giver, but also as a safe preservative for health. Another advantage of using Angkak in making sausages is improving the texture and flavor. Furthermore, according to Sheu et al. (2000), states that the pigments produced by *Monascus purpureus* are very stable and do not change the taste of nata de coco. The dosage used for animal food coloring ranges from 2000-4000 ppm *Monascus* extract while for soft drinks, the concentration used can be lighter which is 0.002% - 0.005% (2-5 ppm).

The curing solution formulation can increase the red color stability of the product during storage, in ham and beef sausage products, the use of nitrite in the curing solution can be reduced from 125 ppm to 80 ppm by adding 2.5 g / kg meat of Angkak pigment. (Fardiaz et al., 2008). Farisandi and Pangesthi (2013) examined the combination of administration of Angkak with sodium nitrite to the organoleptic properties of corned beef, the result was that giving 1% Angkak combined with 50 ppm sodium nitrate affected the color of corned beef products but not flavor and aroma. According to Pattanagu et al. (2007), the optimum use of Angkak in meat products is 1.6% (w / w).

Proteins are high molecular weight complex organic compounds which are polymers of amino acid monomers that are connected to each other by peptide bonds. Proteins generally have a high molecular weight, because of the large weight of protein molecules, so proteins tend to form colloids. Protein solubility depends more on its structure and function and not on its molecular weight. Soluble proteins are good buffers and are very important in maintaining equilibrium reactions. Proteins are formed by units of amino acids that make up polymers so they are long compounds. Protein quality depends on the amino acids it contains. The principle of determining protein profiles by electrophoresis is to separate protein molecules with different charges.

MATERIALS AND METHODS

The ingredients used for the manufacture of sausages are 20 - 24 month old reject chicken meat which has been skinned, washed, 2x2 cm in size taken from the chest and thighs, then separated into 4 parts.

Then the meat is ground and then seasoning is added with a formula from a combination of Bhattacharyya, Mita and Biswas (2005) and Pearson and Dutson (1988), namely: 2% salt, 1.67% sugar, 1.5% garlic, 0.5% pepper, ginger 0.75%, nutmeg 0.5%, which is given in powder form, oil 15%, tapioca flour 5.7%, skim milk 3.5%, ice cubes 16.7% and STPP 0.3% by weight meat.

Each part is given Angkak 0%, 0.5%, 1% and 1.5% in the mixture put in a sleeve with a length of 10 cm and a diameter of 2.5 cm. Then

cooked by steaming at 85⁰C for 30 minutes. Then cooled and analyzed.

The research design carried out was descriptive research. The independent variable in this study was the addition of Angkak to the sausage meat of layered laying hens with concentrations of 0%, 0.5%, 1.0% and 1.5%.

Tools used for protein profile analysis are microtube, beaker glass, Erlenmeyer, electrophoresis chamber, power supply, rotator, mortar cup and spectrophotometer. The method used is SDS-PAGE (Rantam, 2003): Supernatant sample: 20 ml (34) (Phosphate Buffered Saline) solution added 0.5 M NaCl at pH 7.2. A 10 gram sausage sample was pounded with mortar then add a 0.01 ml PBS buffer of 3 ml. Then centrifuged at 6000 rpm for 15 minutes at 4⁰C.

The work procedure of SDS-PAGE is as follows:

Prepare samples:

The protein sample is supplemented by 1:1 Reducing Sample Buffer (RSB) in the Eppendorf tube. Then the sample is heated at 100⁰C for 5 minutes. After being cold, if the sample is not directly used, the sample can be stored at -20⁰C

Prepare separating and stacking gel for 2 plates: Gelling plate is arranged (29) a guide. 15% separating gel is made by: 10% SDS - 60 ml, 10% APS - 60 ml, TEMED 10 ml

Enter the sample in the gel well

A plate that already contains gel is inserted into the electrophoresis chamber. Running buffer is poured until the top and bottom of the gel are submerged. If air bubbles form on the base of the gel or between sample wells, they must be removed. A standard 10 µl marker is inserted in one of the wells (can be gargled at the edge or in the middle well).

Samples of 10 -20 µ (with a minimum protein content of 0.1 µ and a maximum of 20-40 µ) are carefully inserted into the bottom of the gel well, using Hamilton syringe. Syringes are rinsed to 3x using water or by running buffer before being used to insert different samples in the next gel well.

Running sampel

To start running the electrophoresis device is connected to the power supply. Running is carried out at a constant current 20 mA for approximately 40-50 minutes or until tracking

dye reaches a distance of 0.5 cm from the bottom of the gel. After completion, the running buffer is poured and the gel is taken from the plate

Coloring of Gel:

For this stage, a staining solution is needed for coloring gel proteins, the coloring used is Comasie Brilliant Blue or Silver Stain depending on usability. Staining is carried out for 30 minutes. Destaining solution to remove color in the gel and clarify the protein bands formed.

RESULTS AND DISCUSSIONS

Water Content

Decreasing the water content of duck sausage is caused by the increased concentration of Angkak used. Angkak can experience oxidation when heated which causes a decrease in the water in the meat is lost, so that the water content drops (Fardiaz and Zakaria, 1996).

According to Zanardi et al. (2002), the addition of the Angkak concentration produced from the

Monascus purpureus mushroom resulted in a positively charged meat protein and binding to the H + charge, consequently there is no H + that is free or which binds to O which produces free water molecules.

The sausage water content according to the Indonesian National Standard (1995) is a maximum of 67.0%, so the sausage water content from the research which ranges from 63.70 - 64.16% still meets SNI standards.

Fat

This decrease in fat levels is caused by lovastatin compounds in Angkak which act as inhibitors of HMG-CoA reductase (an enzyme that plays a role in cholesterol biosynthesis), where lovastatin is hydrophilic and lipophilic but tends to be lipophilic (Dalimartha, 2001).

The fat content of this study ranged from 8.38 - 9.43%, while fat content according to the Indonesian National Standard (1995) was 18 maximum of 25%, so the fat content of the results of this study still met the standards.

Table 1. The average value of the chemical properties of sausage meat in laying hens is rejected

| Parameter | Con sent rat ion Angkak | | | |
|---------------|----------------------------|----------------------------|----------------------------|----------------------------|
| | 0% | 0,5% | 1% | 1,5% |
| Water Content | 64.16 ± 0,168 ^a | 63.26 ± 0,078 ^a | 63.33 ± 0,213 ^b | 63.70 ± 0,082 ^c |
| Fat | 9.43 ± 0,056 ^a | 9.21 ± 0,055 ^b | 8.98 ± 0,057 ^c | 8.38 ± 0,089 ^d |
| Carbohydrate | 7,75 ± 0,090 ^a | 8.14 ± 0,110 ^b | 9.31 ± 0,029 ^c | 10.89 ± 0,028 ^d |
| Protein | 15,48 ± 0,369 ^a | 15,81 ± 0,088 ^a | 16,09 ± 0,118 ^c | 16,83 ± 0,131 ^d |

Remarks: different notations show significant differences between treatments (P<0.05)

Carbohydrate

The results of the analysis of carbohydrate values indicated that the higher the concentration of Angkak added, the higher the carbohydrate level is. While the carbohydrate carbohydrate levels in this study were 10.89% so that the higher the Angkak added, the higher the carbohydrate content of sausage chicken meat sausages.

Protein

The results of the study for protein levels showed that the higher the concentration of Angkak added, the higher the level of sausage protein in the rejected laying chicken meat. The addition of Angkak in this experiment was immediately given together with seasonings in

the process of making 16 sages. *Monascus* mushrooms that produce enzymes such as α -amylase, β -amylase, glucoamylase, lipase, protease, glucosidase and ribonuclease are able to grow in materials containing starch, protein or lipids (Pattanagu et al., 2007), this condition allows an increase in levels of chicken sausage protein with increasing levels of addition of Angkak to treatments R1, R2, and R3.

Angkak proteins undergo decomposition (oxidative degradation) through a transamination reaction (the enzymatic transfer of amino acid groups from one amino acid to another amino acid) that can bind meat proteins, this is supported by Kramlich (1971) which states that meat protein plays a role in increasing destruction meat during cooking to

form a compact product structure. The role of other proteins is the formation of meat emulsions, which are proteins that function as fat emulsifiers.

Protein content of chicken sausages with Angkak colorant added to the results of this study ranged from 15.48 - 18.83% so that the sausage protein levels still meet SNI standards. Proteins generally have a high molecular weight, because of the large weight of protein molecules, so proteins tend to form colloids. Protein solubility depends more on its structure and function and not on its molecular weight. Soluble proteins are good buffers and are very important in maintaining equilibrium reactions.

Profile protein

The results of SDS-PAGE sausages from rejected laying hens have 7 protein bands that appeared in sausages without the addition of Angkak and the addition of Angkak (0.5%, 1%, 1.5%) with protein molecular weight ranging from 12.44 - and 47.89 kDa (Table 2). Calculation of molecular weight (BM) of protein bands contained in the gel by comparing the molecular weight of the marker and Retardation Factor (Rf), then proceed with making a standard curve with the value of Rf as the x axis and the molecular weight logarithm value as the y axis.

Table 2. Weight of sausage protein molecules

| Pita | Rf | Ro | R1 BM | R2 | R3 |
|------|------|-------|----------|-------|-------|
| 2.1 | 0.33 | 47.89 | 47.89 | 47.89 | 47.89 |
| 2.3 | 0.36 | 42.36 | 42.36 | 42.36 | 42.36 |
| 2.5 | 0.39 | 37.48 | 37.48 | 37.48 | 37.48 |
| 2.1 | 0.45 | 29.33 | 29.33 | 29.33 | 29.33 |
| 3.9 | 0.48 | 25.95 | 25.95 | 25.95 | 25.95 |
| 3.6 | 0.56 | 19.10 | 19.10 | 19.10 | 19.10 |
| 4.3 | 0.67 | 12.44 | 12.44 | 12.44 | 12.44 |





Figure 1. SDS-PAGE Electrophoresis Results
 (R0: Sausage + Without Angkak,
 R1: Sausage + Angkak 0.5%, R2: Sausage + Angkak 1%
 and R3: Sausage + Angkak 1.5%, M: Marker)

Persamaan linear



According to Laemmli (1970), protein bands that are close together indicate that the protein has the same number of amino acids, whereas according to Soepamo (2011) that proteins are formed from amino acids which are bound together to form a series. A small difference in the formation of a series will produce a different type of protein. The types of protein bands detected in processed products are closely related to the functional level of protein damage. Amino acids are increasingly showing the low functional damage to proteins. This is evidenced by the type of protein bands found in meat sausages without the addition of Angkak and added ones which have molecular weights of 12.44 kDa, 19.10 kDa, 25.95 kDa, 29.33 kDa, 42.36 kDa, 37.48 kDa, and 47.89 kDa.

CONCLUSIONS

The addition of Angkak to laying hens sausages can provide changes in water, fat, carbohydrate and protein levels. The molecular weight of the treatment is 0%, 0.5%, 1% and 1.5% ranging from 12.44 kDa - 47. kDa.

REFERENCES

Astawan, M. (2012). *Angkak, Tunjukan Kolesterol*. www.alwadeyonline.com/pengobatan-alternatif/12-angkak-turunkan-kolesterol. Diakses tanggal 14 September 2012.

Bhattacharyya, D., Sinhamahapatra, M., Biswas, S. (2005). Preparation of sausage from spent duck-an acceptability study. *J. Food Sci. Technology*, 42, 24 – 49.

Dalimartha, S. (2001). *36 Resep Tumbuhan Obat Untuk Menurunkan Kolesterol Cetakan ke-3*. Penebar Swadaya, Jakarta.

Fardiaz, S.F.D.B, Zakaria, F. (1996). Toksisitas Dan Imunogenitas Pigmen Angkak Yang Diproduksi Dari Kapang *Monascus purpureus* Pada Substrat Limbah Cair Tapioka. *Buletin Teknologi dan Industri Pangan*, 12, 34-38

Fardiaz, S., Jenie, B.S.L., Rahayu, W.P., Nuraidi, Apriyantono, A., Dewanti, R., Hemarianto (2008). *Produksi Pigmen Untuk Bahan Pewarna Makanan Menggunakan Substrat Limbah Industri Pangan*. <http://anggibit ho-ilmupangan.blogspot.com/2010/02/produksi-pigmen-untuk-bahan-pewarna.html>. Diakses tanggal 25 Juli 2012.

Farisandi, D., Pangesthi, L.T. (2013). Pengaruh Natrium Nitrat Dan Angkak Bubuk Terhadap Sifat Organoleptik Komet. *Ejournal boga*, 2(1), 12-8.

Kramlich, W.E. (1971). *Meat Products*, In Price, J. F. and B.S. Weigert ; *The Science of Meat and Meat Product*. W.H. Freeman and Co. San Fransisco.

Laemmli, UK. (1970). Cleavage en structural proteins during the assembly of the head of Bactiophage T4. *Nature*, 227, 680-85.

Pearson, A.M., Dutson, T.R. (1988). Edible Meat By Product. *Meat Research*, 5, 15 – 42.

Pattanagu, P., Pinthong, R., Phianmongchol, A., Leksawadi, N. (2007). Review Of Angkak Production (*Monascus purpureus*). *Chiang Mai J. Sci.*, 34(3), 319 – 328.

Rantan, F.A. (2003). *Metode Immunologi*. Airlangga University Press. Surabaya. 145-155.

Tisnadjaja, D. (2006). *Bebas Kolesterol dan Demam Berdarah Dengan Angkak*. Jakarta: Penebar Swadaya.

Sheu, F., Wang, C.L., Shyu, Y.T. (2000). Fermentation of *Monascus purpureus* on bacterial cellulose-nata and the color stability of *Monascus-nata* complex. *J. Food Science*, 65(2), 576-581.

Soepamo (2011). *Ilmu Nutrisi Dan Gizi Daging*. Yogyakarta, IND: Gadjah Mada University Press.

Zanardi, E., Dazzi, G., Madarena, G., Chizzoloni, R. (2002). Comparative Study on Nitrite and Nitrate Ions Determination. *Ann. Fac. Medic.*, 22, 79 – 86

PROTEINS PROFIL OF SAUSAGE LAYING CHICKEN MEAT WITH ANGKAK (RED RICE) USED AS NATURAL FOOD MATERIAL

ORIGINALITY REPORT

17%

SIMILARITY INDEX

16%

INTERNET SOURCES

10%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|---|----|
| 1 | ibn.idsi.md Internet Source | 1% |
| 2 | Georgiana Gabriela Codină, Silvia Mironeasa, Costel Mironeasa, Ciprian N Popa, Radiana Tamba-Berehoiu. "Wheat flour dough Alveograph characteristics predicted by Mixolab regression models", Journal of the Science of Food and Agriculture, 2012 Publication | 1% |
| 3 | siat.unpad.ac.id Internet Source | 1% |
| 4 | library.douglascollege.ca Internet Source | 1% |
| 5 | opac.elte.hu Internet Source | 1% |
| 6 | www.agricultureforlife.usamv.ro Internet Source | 1% |
| 7 | sinta3.ristekdikti.go.id Internet Source | 1% |

| | | |
|----|---|------|
| 8 | 123dok.com Internet Source | 1 % |
| 9 | asfa.upatras.gr Internet Source | 1 % |
| 10 | biblioteczka-czytelnicza.blogspot.com Internet Source | <1 % |
| 11 | tdl.libra.titech.ac.jp Internet Source | <1 % |
| 12 | Irma Isnafia Ar, Zakiah Wulandari, Erti Sindy Sin, Dea Marsally S. "Application of Purified Bacteriocin from Lactobacillus plantarum IIA-1A5 as a Bio-preservative of Beef Sausage", Pakistan Journal of Nutrition, 2017 Publication | <1 % |
| 13 | Wei Ding, Hideaki Fujita, Masataka Kawai. "The Length of Cooperative Units on the Thin Filament in Rabbit Psoas Muscle Fibres", Experimental Physiology, 2002 Publication | <1 % |
| 14 | citeseerx.ist.psu.edu Internet Source | <1 % |
| 15 | jast.modares.ac.ir Internet Source | <1 % |
| 16 | onlinelibrary.wiley.com Internet Source | <1 % |
| 17 | ouci.dntb.gov.ua Internet Source | <1 % |

<1 %

18 pt.scribd.com
Internet Source

<1 %

19 ojs.uho.ac.id
Internet Source

<1 %

20 repository.unika.ac.id
Internet Source

<1 %

21 www.semanticscholar.org
Internet Source

<1 %

22 Alfi Rumidatul, Endah Sulistyawa, I Nyoman Pugeg Aryantha. "Trypsin Inhibitor Activity and Protein Analysis of Gall Rust from Sengon Plants (*Falcataria moluccana* Miq.) Infected with *Uromycladium tepperianum* Fungus", *Asian Journal of Plant Sciences*, 2018
Publication

<1 %

23 dspace.uasm.md
Internet Source

<1 %

24 funaab.edu.ng
Internet Source

<1 %

25 eprints.umm.ac.id
Internet Source

<1 %

26 es.scribd.com
Internet Source

<1 %

iopscience.iop.org

| | | |
|----|---|------|
| 27 | Internet Source | <1 % |
| 28 | putrapakrt.blogspot.com Internet Source | <1 % |
| 29 | www.jove.com Internet Source | <1 % |
| 30 | www.tandfonline.com Internet Source | <1 % |
| 31 | www.mdpi.com Internet Source | <1 % |
| 32 | e-journal.undikma.ac.id Internet Source | <1 % |
| 33 | repositori.uin-alauddin.ac.id Internet Source | <1 % |
| 34 | www.bio.mtu.edu Internet Source | <1 % |
| 35 | "Quality Attributes of Muscle Foods", Springer Nature, 1999 Publication | <1 % |
| 36 | hdl.handle.net Internet Source | <1 % |

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off