

Journal Title: Fisheries and Aquatic Sciences
Manuscript ID: fas-2021-0102
Degree (Date created): 4th (2022-04-29)
Manuscript Title: Estimation of first maturity size of dolphinfish *Coryphaena hippurus* Linnaeus in the Molucca Sea, North Sulawesi, Indonesia
Running Title: First maturity size of dolphinfish *Coryphaena hippurus*
Urgency: Normal Manuscript
Type: Short Communication
Category: Ecology and Fisheries Resource Management;
Respond to review: Response to Review

The paper has been shortened to 3,588 words. It means that more than 500 words have been taken out. Normally a short communication contains a maximum of 4,000 words.

Fisheries and Aquatic Sciences

Address: Pukyong National University, 45 Yongso-ro, Nam-gu, Busan 48513, Korea

Tel: +82-51-629-7363,

Email: kosfas@kosfas.or.kr

Homepage: <http://www.e-fas.org/>

1 Estimation of first maturity size of dolphinfish *Coryphaena hippurus* Linnaeus in the Molucca Sea, North
2 Sulawesi, Indonesia

3 Silvester Benny Pratasik*, Ferdinand F. Tilaar, Sofie M. Salaki

4 Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus-Bahu, Manado-95115,
5 North Sulawesi, Indonesia.

6 Correspondence: Silvester B. Pratasik; email: spjong07@yahoo.com

7 **Abstract.** This study aims to estimate the smallest size of mature individuals that can be exploited. Fish samples of
8 *Coryphaena hippurus* were collected from Kalinaun fishermen's catches in the Molucca Sea. They were sexed, then
9 the fork length (FL) and maturity stage were recorded. Results showed that *C. hippurus* in the Molucca Sea had a sex
10 ratio of 1: 1.94 ($P < 0.05$). Males had a length range of 499 – 831 mm FL and females were in the length range of 481-
11 813 mm FL. Size at first maturity was estimated as 529 mm FL for males with a range of 475-588 mm FL and 405
12 mm FL for females. This study provided basic information for future management needs of the dolphinfish, especially
13 in the Molucca Sea.

14 **Keywords (3 to 5):**

15 sex, fork length (FL), maturity stage, Kalinaun, fishermen.

16

Introduction

17 Fisheries management must be directed to maintain the fish populations remain sufficiently abundant to
18 minimize extinction risk and sustain intact ecosystems (Freshwater et al., 2020). Fish reproduction is an important
19 aspect in maintaining the equilibrium of the fish stock population in the water since stock recovery is highly dependent
20 upon reproductive success that is closely related to environmental changes particularly temperature, photoperiod, and
21 food supply (Bagenal, 1978). Thus, fecundity, sexual maturity, and spawning habits must be understood to explain
22 the variation of the population level to increase the amount of fish harvest and maintain the recovery rate (Das et al.,
23 1989; Brown et al., 2003).

24 Dolphinfish, *Coryphaena hippurus* (Linnaeus, 1758) (Coryphaenidae), known as mahi-mahi, is a commercially
25 important species in tropical and temperate waters worldwide that generally inhabit open waters (Benyamin & Kurup,
26 2012) in line with the tuna catch decline in the Indian Ocean since 2007 (IOTC 2012). This species has a sufficiently

27 large size, the young one is about 30 cm long and the adults can reach 200 cm long with bodyweight up to 50 kg. The
28 individual weight of fish caught ranges from 7 to 13 kg and rarely reaches 15 kg. The species is caught as bycatch in
29 several types of fishing gears, such as purse seine, longline, and trolling (Chodrijah & Nugroho, 2016).

30 *C. hippurus* is a long-range and fast swimming fish that displace with time and is an opportunistic epipelagic
31 predator and preys on biota associated with a fish aggregating device (FAD) and floating debris, such as fish, squids,
32 and shrimps (Malone et al., 2011; Whitney et al., 2016). *C. hippurus* can stay several days in association with a raft
33 (Taquet et al., 2007). Dolphinfish spend >80% of daytime activity and 40% of nighttime activity near the surface (Lin
34 et al., 2020) and inhabit warmer seawater temperatures of 24°C- 30°C (Palko et al., 1982; Schlenker et al., 2021).
35 When surface sea temperature (SST) rises, dolphinfish use behavioral thermoregulation by moving deeper up to 250
36 m, and the nighttime activity increased with increasing lunar illumination (Schlenker et al., 2021).

37 The IUCN status of dolphinfish is the least concerned (Carlson et al., 2020). This study is aimed at estimating
38 the size at first maturity of Dolphinfish *C. hippurus* caught in the Molucca Sea, North Sulawesi. Size at first maturity
39 is the smallest size of mature legally taken, the size at which 50% of the individuals are sexually mature (Farley et al.,
40 2013). Knowledge of length at maturity and spawning season is important for the proper management and conservation
41 of fish stocks (Nandikeswari, 2016). Size at first maturity is commonly evaluated for wild populations as a point of
42 biological reference to ensure that a sufficient number of juveniles reaches maturity (Roa et al., 1999) because only
43 fishing the individuals which have reached maturity is one of the basic rules that should be followed to ensure
44 sustainability (Ilkyas et al., 2018). It has been utilized in various exploited animals, such as crustaceans (Peixoto et
45 al., 2018), fish populations (Tsfahun, 2018), and mollusks (Galimany et al., 2015). Proper estimation of size at first
46 maturity is very useful for fish stock management (Karna et al., 2011). These data provide basic information on fish
47 biology that is crucial for dolphinfish fisheries management in Indonesian waters and other neighboring countries.

48

49

Materials and Methods

50 Dolphinfish *Coryphaena hippurus* samples were mainly collected from fishermen in the Kalinaun coast, East
51 Likupang district, North Minahasa, North Sulawesi. The fish samples were obtained from May to July 2021, because
52 there was no catch after this period. Fishing activity was conducted near a man-made Fish Aggregating Device (FAD)
53 in the Molucca Sea located in the northeastern part of the village between 125°11'24" E and 125°13'48" E and
54 1°35'24" and 1°35'24" N. Local fishermen usually used live bait-handline. Live baits were obtained in the multi-hooks
55 handline fishing before daybreak. Trolling was also carried out around the FAD to obtain more samples. The fish were

56 sexed on the beach. The fork length and weight were also recorded, then the gonads were removed and brought to the
 57 Laboratory of the Faculty of Fisheries and Marine Sciences, Sam Ratulangi University, Manado, for further
 58 observation. The estimation of sex ratio used a non-parametric comparative test Chi-Square (χ^2 , $\alpha = 0.05$). **Gonadal**
 59 **maturity was observed under a dissecting microscope. The fish maturity stage was identified following Effendie**
 60 (2002) (Table 1).

61
 62 Table 1. Gonad maturity characteristics.

63 The first gonad maturity was estimated by setting the size class intervals, from the smallest to the largest one. Length
 64 distribution analysis followed Sturges (1926) as follows:

$$65 \quad k = 1 + 3.3 \log n$$

66 where k is number of classes and n is number of data. The class interval was estimated as

$$67 \quad C = \frac{X_n - X_1}{k}$$

68 where C is class interval, X_n is the largest data value, X_1 is the smallest data value, and
 69 k is number of classes.

70 Spearman-Kärber equation was applied to estimate the size at first maturity of the fish (Udupa, 1986) as follows:

$$71 \quad m = x_k + \frac{x}{2} - (x \sum p_i)$$

72 where x_k = log last size in which 100% fish are fully mature

$$73 \quad x = \log \text{ size increment} = x_{l+1} - x_l, l = 1, 2, \dots, k-1$$

74 and x_0 = log last size in which no fish are fully mature

$$75 \quad r_i = \text{number of fully mature fish in size group } i$$

$$76 \quad p_i = \text{proportion of fully mature fish in size group } i$$

$$77 \quad p_i = r_i/n_i, \text{ if } n_i \neq n_{i+1} \text{ for } i = 1, 2, \dots, k-i$$

$$78 \quad \text{and } p_i = r_i/n, \text{ if } n = n_i = n_{i+1} \text{ for } i = 1, 2, \dots, k-i$$

79 Size at first maturity was obtained with $\text{antilog}(m) = M$.

$$80 \quad \text{antilog} \left[m \pm 1.96 \sqrt{x^2 \sum_i \left\{ \frac{(p_i - q_i)}{n_i - 1} \right\}} \right]$$

81 Results

82 During the study 50 fish individuals were collected from local fishermen in Kalinaun, East Likupang District,
 83 North Minahasa Regency, North Sulawesi. Males had a size range of 405 mm - 674 mm FL with a weight range of

84 670 – 1,640 g, and females were at a length range of 431 mm - 687 mm FL with a weight range of 725 – 2,650 g.
85 Based on Sturges (1926), the length distribution was divided into 6 size class intervals.

86

87 Figure 1. Length frequency of *C. hippurus* caught in the Molucca Sea.

88

89 Sex ratio, maturity stage, and size at first maturity

90 Sex ratio information is useful to maximize reproduction. The present study found a sex ratio of 1:1.94 ($P < 0.05$)
91 represented by 17 males and 33 females. Gonad maturity of this species shows that more females mature at a smaller
92 size than males (Table 2).

93

94 Table 2. Gonad maturity stage

95

96 Size at first maturity was estimated as 529 mm FL for males with a range of 475 – 588 mm and 405 mm FL
97 for females.

98 Discussion

99 This low number of catches could result from that *C. hippurus* is not a target species. Local fishermen in this
100 area go fishing for yellowfin tuna, marlin, and sharks, whereas *C. hippurus* is optional when the target fish are not
101 found due to the low market value of this species. Field observations also revealed that the occurrence of *C. hippurus*
102 in this region is seasonal. Besides, although the fish are around, they did not bite at all in trolling or live bait fishing.
103 Only a few individuals of *C. hippurus* are caught, usually 1-5 individuals per boat. However, there is still no study
104 on the fishing season of *C. hippurus*, particularly in this area.

105 A previous study on dolphinfish landing in the Bitung Fisheries Port found 4,160 individuals of *C. hippurus* in
106 the size range of 300 mm FL – 1,210 mm FL with a mean length of 598 ± 13.9 mm FL (Chodrijah & Nugroho, 2016)
107 reflecting small size dominance. The fish samples came from catches of many kinds of fishing gears, such as purse
108 seine, longline, and trolling. **The present study found narrower size distribution, and it could result from less number
109 of samples obtained due to high dependence on local artisanal fishermen who rely on hand-line fishing.**

110 The present size range is far below the maximum individual size previously reported reflecting that the mean
111 individual size of *C. hippurus* has been declining. The recovery rate of a population is related to the mortality rate, the
112 closer the mean individual size to the maximum, the lower the mortality rate (ECTF, 2004). The present finding

113 revealed that the dolphinfish population has a high mortality rate. However, so many factors influence fish population
114 availability in the ocean. This condition is supported by Goldstein et al. (2007) that life-history traits are vulnerable to
115 environmental stress and fishing pressure that result in smaller mature fishes as a response for survival. Fish mortality
116 could occur at specific stages and species and the causes may be single or cumulative pressure from a range of sources,
117 such as pollutants, anthropogenic climate change or natural variability (Olsen et al., 2019), and fishing activities.
118 Recruitment patterns with time can influence the population size as well, and therefore, mortality events in the early
119 life stages may have severe and long-lasting effects on the population (Langangen et al., 2017). Climate change is
120 another factor causing changes in fish populations, which can affect the distribution of particular species and the fish
121 susceptibility to particular fishing fleets (Rijnsdorp et al., 2009). This condition could occur because population size
122 has probably fallen below some threshold level of abundance in which the rate of recovery cannot well respond to the
123 fishing rate.

124 This sex deviation is similar to that reported in the western and central Mediterranean (Potoschi et al., 1999;
125 Benseddik et al., 2019) reflecting sex segregation in *C. hippurus* until reaching the mature stage. Mature individuals
126 seem to gather in the same area for spawning and feeding around the rafts so that more females were caught than
127 males. This result also agrees with Perle et al. (2020) and Oxenford (1999) that sex segregation occurs in *C. hippurus*
128 or males are more susceptible to fisheries than females, even though our finding found more females than males. A
129 higher proportion of females from FADs captures could result from greater availability of females, higher natural
130 mortality in males, or differential growth of both sexes (Benseddik et al., 2019). Moreover, males and females show
131 different maturity stages with size class (Table 2). Both sexes show bigger individual sizes than 400 mm FL with
132 more females at mature stages (III and IV). It indicates that males need a bigger size to reach gonad maturity or females
133 reach gonad maturity earlier than males. These data are consistent with Beardsley (1967) that female dolphinfish begin
134 to mature (reach stage II) at about 350 mm FL (about 6-7 months old), 50% are mature at 450 mm FL, and 100% are
135 mature at 550 mm FL, whereas males are mature at a slightly larger size (427 mm FL). Nevertheless, in the Eastern
136 Tunisian Coast, Central Mediterranean, Benseddik et al. (2019) found that the first maturity size of *C. hippurus* occurs
137 at 553 mm FL for females and 605 mm FL for males. In the present study, females above 400 mm FL reached maturity
138 stages III and IV. This difference could result from different environmental conditions in localities. It means that 50%
139 of mature individuals that occurs at this size, particularly in the Molucca Sea population, could be set as the minimum
140 legal size of this species to meet the sustainability criteria and avoid economic loss due to fishing immature individuals.
141 The use of minimum legal size in fisheries is intended to protect juveniles, let them grow into adults, and spawn at

142 least once before being caught. The size range of *C. hippurus* caught in the Molucca Sea reflects mature individuals
143 and has mostly passed the size at first maturity. Nevertheless, since fishing is a major factor in reducing size and age
144 at first maturity (McIntyre & Hutchings, 2003) and a decline in age and size at maturity may negatively affect the fish
145 recovery (Hutchings, 2002), it needs to be controlled. The individual size decline of *C. hippurus* far below the
146 maximum size could have indicated a reduced population size and should not be ignored. Earlier maturity can be
147 associated with reduced longevity, increased post-reproductive mortality, and smaller sizes at reproductive age.
148 Populations composed of small individuals will reduce reproductive potential (Scott et al., 1999), increase variance in
149 offspring survival (Hutchings & Myers, 1993), and eventually negatively affect population growth.

150 Mesh size control and escapement could be an alternative to maintain or increase the individual size range or
151 even increase the longevity, and the reproductive potentiality of dolphinfish. Larger fish have higher fecundity and
152 can produce more eggs. So far, commercial purse seiners (< 30 GT) for small pelagic fish have fished any fish schools
153 encountered in the open sea using small mesh sizes. As a result, small yellowfin tuna, skipjack, and dolphinfish are
154 also caught (field obs.). Also, fishing gear separation should be established for commercial small pelagic and large
155 pelagic fisheries to maintain stock availability and prevent individual size decline. This effort limitation could help
156 reduce the risk of population collapse and become one of the remedies to population recovery. Fish population
157 recovery, therefore, requires institutional structures that either entice fishers to leave the business, through expensive
158 buyout schemes of fishing boats and licenses or force them to reduce fishing activity (Hutching & Reynolds, 2004).

159 The present study has contributed to providing important biological information for future management,
160 especially dolphinfish *C. hippurus* of Molucca Sea. A long-term study on the biology and ecology of this species is
161 required to well describe the population status of *C. hippurus* so that the management policy could be strengthened.
162 The fisheries committee among neighborhood countries that take advantage of the resources should also participate
163 in sustainable resource utilization programs by maintaining the exploitation level and the ecosystem equilibrium.

164 **References (NLM style; Alphabetical Order)**

- 165 Bagenal TB. Aspects of fish fecundity, In Gerking, SD (ed). Methods of Assessment of Ecology of Freshwater Fish
166 Production. Blackwell, London, 1978; 75-101.
- 167 Beardsley GL, Jr. Age, growth, and reproduction of the dolphin, *Coryphaena hippurus*, in the strait of Florida. Copeia
168 1967:441-451.
- 169 Benjamin D, Kurup BM. Stock assessment of Dolphinfish, *Coryphaena hippurus* (Linnaeus, 1758) off the southwest
170 coast of India Journal of the Marine Biological Association of India 2012; 54(1):95-99.

171 Benseddik AB, Besbes R, Missaoui H, Najaï SE, Jarboui O. Reproductive dynamics and fecundity of *Coryphaena*
172 *hippurus* (Linnaeus,1758) in the Eastern Tunisian Coast (Central Mediterranean). *Current Trends in Fisheries*
173 *and Aquaculture* 2019; 1:23 p.

174 Brown P, Sivakumaran KP, Stoessel D, Giles A, Green C, Walker T. Carp population biology in Victoria. Report No.
175 56, Marine and Freshwater Resources Institute, Department of Primary Industries, Snobs Creek, Victoria. 2003;
176 p. 202.

177 Carlson AK, Rubenstein DI, Levin SA Linking multiscale fisheries using meta coupling models. *Front. Mar. Sci.*
178 2020; 7:614; DOI: 10.3389/fmars.2020.00614. 17 p.

179 Chodrijah U, Nugroho D. Size structure and population parameters of dolphinfish (*Coryphaena hippurus* Linnaeus,
180 1758) in the Celebes Sea. *Bawal* 2016; 8(3):147-158 [in Indonesian]

181 Das M, Dewan S, Debnath SC. Studies on the fecundity of *Heteropneustes fossilis* (Bloch) in a mini pond of
182 Bangladesh Agricultural University, Mymensingh. *Bangladesh Journal of Agricultural Sciences* 1989; 16:1-
183 6.

184 ECTF. General Effort Review: Sustainability of permitted species. 2004;
185 https://www.daf.qld.gov.au/__data/assets/pdf_file/0003/76629/StockAssessment-CTrawl-2004-Part9.

186 Effendie MI. *Biologi Perikanan (Fisheries Biology)*. Yayasan Dewi Sri. Bogor. 2002; 163 p. (In Indonesian).

187 Farley J, Davies C, Hillary R, Eveson P. Estimating size/age at of southern bluefin tuna. CCSBTESC/1309/41, 18th
188 Meeting of the Scientific Committee, 2-7 September 2013, Canberra, Australia. 7 p.

189 Freshwater C, Holt KR, Huang A-M, Holt CA. Benefits and limitations of increasing the stock-selectivity of Pacific
190 salmon fisheries. *Fisheries Research* 2020; 226:1-9.

191 Galimany E, Baeta M, Durfort M, Lleonart J, Ramón M. Reproduction and size at first maturity in the Mediterranean
192 exploited *Callista Chione* bivalve bed. *Scientia marina* 2015; 79(20):233-242.

193 Goldstein J, Heppell S, Cooper A, Brault S, Lutcavage M. Reproductive status and body condition of Atlantic bluefin
194 tuna in the Gulf of Maine, 2000–2002 *Mar Biol.* 2007; 151, 2063. DOI 10.1007/s00227-007-0638-8.

195 Hutchings JA. Life histories of fish. in P.J.B. Hart, J.D. Reynolds eds. *Handbook of Fish and Fisheries*, vol. 1. Oxford
196 (United Kingdom): Blackwell 2002; 149–174

197 Hutchings JA, Myers RA. Effect of age on the seasonality of maturation and spawning of Atlantic cod, *Gadus morhua*,
198 in the Northwest Atlantic. *Canadian Journal of Fisheries and Aquatic Sciences* 1993; 50:2468–2474.

199 Hutchings JA, Reynolds JD. Marine fish population collapses: consequences for recovery and extinction risk.
200 Bioscience 2004; 54(4):297-309.

201 Ilkyaz AT, Metin G, Soykan O, Kinacigil HT. Spawning season, first maturity length, and age of 21 fish species from
202 the Central Aegean Sea, Turkey. Turkish Journal of Fish. and Aq. Sci., 2018; 18: 211-216. DOI: 10.4194/1303-
203 2712-v18_1_24

204 IOTC. Review of the statistical data and fishery trends for tropical tunas. Working Party of Tropical Tunas, Mauritius
205 24-29 October 2012; 63 pp.

206 Karna SK, Panda S. Growth estimation and Length at maturity of a commercially important fish species i. e.,
207 *Daysciaena albida* (Boroga) in Chilika Lagoon, India. Euro. J. Exp. Bio., 2011, 1(2):84-91.

208 Langangen Ø, Ohlberger J, Stige LC, Durant JM, Ravagnan E, Stenseth NC, et al. Cascading effects of mass mortality
209 events in Arctic marine communities. Glob. Change Biol. 2017; 23:283–292. DOI: 10.1111/gcb.13344

210 Lin SJ, Chiang WC, Musyl MK, Wang SP, Su NJ, Chang QX, Ho YS, Nakamura I, Tseng CT, Kawabe R. Movements
211 and habitat use of Dolphinfish (*Coryphaena hippurus*) in the East China Sea. Sustainability 2020; 12, 5793:16
212 p.

213 Malone MA, Buck KM, Moreno G., Sancho G. Diet of three large pelagic fishes associated with drifting fish
214 aggregating devices (DFADs) in the Western Equatorial Indian Ocean. Animal Biodiversity and Conservation
215 2011; 34(2):287-294.

216 McIntyre TM., Hutchings JA. Small-scale temporal and spatial variation in Atlantic cod (*Gadus morhua*) life history.
217 Canadian Journal of Fisheries and Aquatic Sciences 2003; 60:1111–1121.

218 Nandikeswari R. Size at first maturity and maturity stages of *Terapon jarbua* (Forsskal, 1775) from Pondicherry Coast,
219 India. Journal of Fisheries 2016; 4(2):385-389.

220 Olsen E, Hansen C, Nilsen I, Perryman H, Vikebø F. Ecological effects and ecosystem shifts caused by mass mortality
221 events on early life stages of fish. Front. Mar. Sci. 2019; <https://doi.org/10.3389/fmars.2019.00669>. 13 p.

222 Oxenford HA. Biology of the dolphinfish (*Coryphaena hippurus*) in the Western Central Atlantic. Scientia Marina
223 1999; 63:277-301.

224 Palko JB, Beardsley GL, Richards WJ. Synopsis of the biological data on dolphinfish, *Coryphaena hippurus* Linnaeus
225 and *Coryphaena equiselis* Linnaeus. NOAA. Tech. Rep. NMFS Circ. 1982; 443. 28 pp.

226 Peixoto S, Calazans N, Silva EF, Nole L, Soares R, Frédou FL. Reproductive cycle and size at first sexual maturity of
227 the white shrimp *Penaeus schmitti* (Burkenroad, 1936) in northeastern Brazil. Lat. Am. J. Aquat. Res. 2018;
228 46(1):1-9.

229 Perle CR, Snyder S, Merten W, Simmons M, Dacey J, Rodriguez Sanchez R, O'Sullivan J, Ortega GS. Dolphin
230 movements in the Eastern Pacific Ocean of Mexico using conventional and electronic tags. Animal
231 Biotelemetry 2020; 8 (30): 15 p.

232 Potoschi A, Renones O, Cannizzaro L. Sexual development, maturity and reproduction of dolphinfish (*Coryphaena*
233 *hippurus*) in the western and central Mediterranean. Sci. Mar. 1999; 63(3-4):367-372.

234 Rijnsdorp AD, Peck MA, Engelhard GH, Mo'llmann C, Pinnegar JK. Resolving the effect of climate change on fish
235 populations. – ICES Journal of Marine Science 2009; 66:1570–1583.

236 Roa R, Ernst B, Tapia F. Estimation of size at sexual maturity: an evaluation of analytical and resampling procedures.
237 Fish. Bull. 1999; 97:570–580.

238 Schlenker LS, Faillettaz R, Stieglitz JD, Lam CH, Hoenig RH, Cox GK, Heuer RM, Pasparakis C, Benetti DD, Paris
239 CB, Grosell M Remote predictions of mahi-mahi (*Coryphaena hippurus*) spawning in the open ocean using
240 summarized accelerometry data. Front. Mar. Sci. 2021; 8: 1-18. <https://doi.org/10.3389/fmars.2021.626082>.

241 Scott B, Marteinsdottir G, Wright P. Potential effects of maternal factors on spawning stock-recruitment relationships
242 under varying fishing pressure. Canadian Journal of Fisheries and Aquatic Sciences 1999; 56:1882–1890.

243 Sturges HA. The choice of a class interval. Journal of the American Statistical Association 1926; 21(153):65-66.

244 Taquet M, Dagorn L, Gaertner J, Girard C, Aumerruddy R, Sancho G, Itano D. Behavior of dolphinfish (*Coryphaena*
245 *hippurus*) around drifting FADs as observed from automated acoustic receivers. Aquatic Living Resources
246 2007; 20(4):323-330.

247 Tesfahun A. Overview of length-weight relationship, condition factor, and size at first maturity of Nile tilapia
248 *Oreochromis niloticus* (L.) in different water bodies of Ethiopia. A Review. Greener Journal of Biological
249 Sciences 2018; 8(3): 021-028.

250 Udupa KS. Statistical method of estimating the size at first maturity in fishes. Univ. Agricult. Sci. College of Fish.,
251 Mangalore, India 1986; 4(2):8-10.

252 Whitney NM, Schwieterman GD, Taquet M, Brill RW, Dagorn L, Holland KN., Girard C. Swimming depth of
253 dolphinfish (*Coryphaena hippurus*) associated and unassociated with fish aggregating devices. Fishery Bulletin
254 - National Oceanic and Atmospheric Administration 2016; 114(4):426-434.


Table 1. Gonad maturity characteristics

Maturity stage	Note	Female	Male
I	Immature	Small ovary up to ½ the length of the body cavity. It is translucent. Oocyte does not appear.	The testis is small up to ½ the length of the body cavity. It is whitish.
II	Maturing	The ovary is about half the length of the body cavity. It is orange and translucent, and the oocyte cannot be seen by the naked eye.	The testis is about ½ the length of the body cavity. It is white and about symmetrical.
III	Ripening	The ovary is about 2/3 the length of the body cavity. Ovary yellow-orange, oocyte appears. Ovary with blood vessels on the surface. No transparent eggs or translucent, eggs are still dark.	The testis is about 2/3 the length of the body cavity.
IV	Ripe	The ovary is about 2/3 up to full of the body cavity. The ovary is orange-pink with blood vessels on the surface, and eggs are apparent.	The testis is about 2/3 up to fulfilling the body cavity. It is white-soft cream.
V	Spent	Ovary shrinks down to ½ the body cavity. Wall is thick. There may be dark and mature eggs in the ovary that disintegrate from absorption, dark or translucent.	Testis shrinks down to ½ the body cavity. Wall is thick. The testis is soft.

Table 2. Gonad maturity stage of *C. hippurus* recorded in this study.

Size class (mm)	Gonad Maturity Stage (N=50)							
	I		II		III		IV	
	Male	Female	Male	Female	Male	Female	Male	Female
400-449	0	0	4	0	3	3	0	0
450-499	1	0	0	0	1	6	0	3
500-549	1	0	3	0	0	7	1	1
550-599	0	0	0	0	0	8	1	1
600-649	0	0	0	0	0	2	0	0
650-699	0	0	0	0	1	2	1	0

Figure 1. Length frequency distribution of dolphinfish *C. hippurus*



Response to Review

The paper has been shortened to 3,588 words. It means that more than 500 words have been taken out. Normally a short communication contains a maximum of 4,000 words.