



1 Estimation of first maturity size of dolphinfish *Coryphaena hippurus* Linnaeus in the Molucca Sea,
2 North Sulawesi, Indonesia

3 Silvester Benny Pratasik*, Ferdinand ~~Frans F.~~ Tilaar, ~~Sofie Meiske M.~~ Salaki
4 Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus-Bahu, Manado_95115,
5 ~~North Sulawesi~~, Indonesia

6 -

7 *Corresponding ~~author~~ ~~ence~~: Silvester ~~Benny B.~~ Pratasik

8 [Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus-Bahu, Manado 95115,](#)
9 [Indonesia](#)

10

11 [Tel: +62-431-6868027](#); [E-mail: spjong07@yahoo.com](#) [spjong07@yahoo.com](#)

12

13 **Abstract.**

14 This study aims to estimate the smallest size of mature individuals that can be exploited. Fish samples of
15 *Coryphaena hippurus* were collected from Kalinaun fishermen's catches in the Molucca Sea. They were sexed,
16 then the fork length (FL) and maturity stage were recorded. Results showed that *C. hippurus* in the Molucca Sea
17 had a sex ratio of 1:-1.94 ($P_2 < 0.05$). Males had a length range of 499—831 mm FL and females were in the
18 length range of 481—813 mm FL. Size at first maturity was estimated as 529 mm FL for males with a range of
19 475—588 mm FL and 405 mm FL for females. This study provided basic information for future management
20 needs of the dolphinfish, especially in the Molucca Sea.

21 **Keywords (3 to 5):**

22 Sex, Fork length (FL), Maturity stage, Kalinaun, Fishermen-

23

24

25

Introduction

메모 포함[GM61]: Please check.
The author's name in ORCID ID is "Salaki MS" not "Salaki SM". Please make sure which name is correct.

메모 포함[GM62]: Telephone number has been added by referring to the information in the submission system. Please check if the number is correct.

26 Fisheries management must be directed to maintain the fish populations remain sufficiently abundant to
27 minimize extinction risk and sustain intact ecosystems (Freshwater et al., 2020). Fish reproduction is an important
28 aspect in maintaining the equilibrium of the fish stock population in the water since stock recovery is highly
29 dependent upon reproductive success that is closely related to environmental changes particularly temperature,
30 photoperiod, and food supply (Bagenal, 1978). Thus, fecundity, sexual maturity, and spawning habits must be
31 understood to explain the variation of the population level to increase the amount of fish harvest and maintain the
32 recovery rate (Das et al., 1989; Brown et al., 2003; Das et al., 1989).

33 Dolphinfish, *Coryphaena hippurus* (Linnaeus, 1758) (Coryphaenidae), known as mahi-mahi, is a
34 commercially important species in tropical and temperate waters worldwide (Benjamin & Kurup, 2012) in line
35 with the tuna catch decline in the Indian Ocean (IOTC, 2012).- This species has a sufficiently large size, the young
36 one is about 30 cm long and the adults can reach 200 cm long with bodyweight up to 50 kg. The individual weight
37 of fish caught ranges from 7 to 13 kg and rarely reaches 15 kg. The species is caught as bycatch in several types
38 of fishing gears, such as purse seine, longline, and trolling (Chodriyah & Nugroho, 2016).

39 *C. hippurus* is a long-range and fast swimming fish that displace with time and is an opportunistic
40 epipelagic predator and preys on biota associated with a fish aggregating device (FAD) and floating debris, such
41 as fish, squids, and shrimps (Malone et al., 2011; Whitney et al., 2016). *C. hippurus* can stay several days in
42 association with a raft (Taquet et al., 2007). Dolphinfish spend > 80% of daytime activity and 40% of nighttime
43 activity near the surface (Lin et al., 2020) and inhabit warmer seawater temperatures of 24°C–30°C (Palko et al.,
44 1982; Schlenker et al., 2021). When surface sea temperature (SST) rises, dolphinfish use behavioral
45 thermoregulation by moving deeper up to 250 m, and the nighttime activity increased with increasing lunar
46 illumination (Schlenker et al., 2021). The IUCN status of dolphinfish is the least concerned (Carlson et al., 2020).

47 This study is aimed at estimating the size at first maturity of ~~Dolphinfish-dolphinfish~~ *C. hippurus* caught
48 in the Molucca Sea, North Sulawesi. Size at first maturity is the smallest size of mature legally taken, the size at
49 which 50% of the individuals are sexually mature (Farley et al., 2013). Knowledge of length at maturity and
50 spawning season is important for the proper management and conservation of fish stocks (Nandikeswari, 2016).
51 Size at first maturity is commonly evaluated for wild populations as a point of biological reference to ensure that
52 a sufficient number of juveniles reaches maturity (Roa et al., 1999) because only fishing the individuals which
53 have reached maturity is one of the basic rules that should be followed to ensure sustainability (Ilkyaş et al.,
54 2018). It has been utilized in various exploited animals, such as crustaceans (Peixoto et al., 2018), fish
55 populations (Tesfahun, 2018), and mollusks (Galimany et al., 2015), to protect juveniles, let them grow into adults,
56 and spawn at least once before being caught. -Proper estimation of size at first maturity is very useful for fish

메모 포함[U3]: In accordance with the regulations, references have been revised to 'By alphabet' order. Please confirm that the correction is correct.

메모 포함[GM64]: This term appears only once in the text, so abbreviation SST has been omitted. Please check.

Please check the full name of the abbreviation. It is recommended that you specify full names instead of abbreviations in tables, figures, and titles. Please check if there is anything wrong. Ex) TBA (full name)

57 stock management (Kama et al. & Panda, 2011). These data provide basic information on fish biology that is
58 crucial for dolphinfish fisheries management in Indonesian waters and other neighboring countries.

60 **Materials and Methods**

61 Dolphinfish *Coryphaena-C. hippurus* samples were mainly collected from fishermen in the Kalinaun coast,
62 East Likupang district, North Minahasa, North Sulawesi. The fish samples were obtained from May to July 2021,
63 because there was no catch after this period. Fishing activity was conducted near a man-made Fish Aggregating
64 Device (FAD) in the Molucca Sea located in the northeastern part of the village between 125°11'24" E and
65 125°13'48" E and 1°35'24" and 1°35'24" N. Local fishermen usually used live bait-handline. Live baits were
66 obtained in the multi-hooks handline fishing before daybreak. Trolling was also carried out around the FAD to
67 obtain more samples. The fish were sexed on the beach. The fork length (FL) and weight were also recorded, then
68 the gonads were removed and brought to the Laboratory of the Faculty of Fisheries and Marine Sciences, Sam
69 Ratulangi University, Manado, for further observation. The estimation of sex ratio used a non-parametric
70 comparative test Chi-Square-square (χ^2 , $\alpha = 0.05$). Gonadal maturation was observed under a dissecting
71 microscope. The fish maturity stage was identified following Effendic (2002) (Table 1).

73
74 **Table 1. Gonad maturity characteristics.**

75 The first gonad maturity was estimated by setting the size class intervals, from the smallest to the largest one.
76 Length distribution analysis followed Sturges (1926) as follows:

$$78 \quad k = 1 + 3.3 \log n$$

79
80 where k is number of classes and n is number of data. The class interval was estimated as

$$81 \quad C = \frac{X_n - X_1}{k}$$

82
83
84 where C is class interval, X_n is the largest data value, X_1 is the smallest data value, and
85 k is number of classes.

86 Spearman-Kärber equation was applied to estimate the size at first maturity of the fish (Udupa, 1986) as follows:

88
$$m = x_k + \frac{x}{2} - (\sum p_i)$$

89
90 where x_k = log last size in which 100% fish are fully mature

91 x = log size increment = $x_{i+1} - x_i$, $i = 1, 2, \dots, k-1$

92 and x_0 = log last size in which no fish are fully mature

93 r_i = number of fully mature fish in size group i

94 p_i = proportion of fully mature fish in size group i

95 $p_i = r_i/n_i$, if $n_i \neq n_{i+1}$ for $i = 1, 2, \dots, k-1$

96 and $p_i = r_i/n$, if $n_i = n_{i+1}$ for $i = 1, 2, \dots, k-1$

97 Size at first maturity was obtained with antilog (m) = M .

98
99
$$\text{antilog} \left[m \pm 1.96 \sqrt{x^2 \sum \left\{ \frac{(p_i - q_i)}{n_i - 1} \right\}} \right] \text{antilog} \left[m + 1.96 \sqrt{x^2 \sum \left\{ \frac{(p_i - q_i)}{n_i - 1} \right\}} \right]$$

메모 포함[GM65]: The square brackets "]" has been added. Please make sure if the formula is correct.
we modified this equation. please check it is right.

Results

102 During the study 50 fish individuals were collected from local fishermen in Kalinaun coast, East Likupang District,
103 North Minahasa Regency, North Sulawesi. Males had a size range of 405–674 mm FL with a weight range
104 of 670–1,640 g, and females were at a length range of 431–687 mm FL with a weight range of 725–2,650
105 g. Based on Sturges (1926), the length distribution was divided into 6 size classes. (Fig. 1).

메모 포함[GM66]: Figure must be cited in the text. So "Fig. 1" has been added here. Please check if it is correct.

108 *Figure 1. Length frequency of C. hippurus caught in the Molucca Sea.*

Sex ratio, maturity stage, and size at first maturity

111 Sex ratio information is useful to maximize reproduction. The present study found a sex ratio of 1:1.94 (P
112 $p < 0.05$) represented by 17 males and 33 females. Gonad maturity of this species shows that more females mature
113 at a smaller size than males (Table 2).

115 *Table 2. Gonad maturity stage*

117 Size at first maturity was estimated as 529 mm FL for males with a range of 475–588 mm and 405 mm
118 FL for females.
119

120 Discussion

121 This low number of catches could result from that *C. hippurus* is not a target species. Local fishermen in
122 this area go fishing for yellowfin tuna, marlin, and sharks, whereas *C. hippurus* is optional when the target fish
123 are not found due to the low local market value of this species. Field observations also revealed that the occurrence
124 of *C. hippurus* in this region is seasonal. Besides, although the fish are around, they did not bite at all in trolling
125 or live bait fishing. Only a few individuals of *C. hippurus* are caught, usually 1–5 individuals per boat. However,
126 there is still no study on the fishing season of *C. hippurus*, particularly in this area.

127 A previous study on dolphinfish landing in the Bitung Fisheries Port found 4,160 individuals of *C. hippurus*
128 in the size range of 300–1,210 mm FL with a mean length of 598 ± 13.9 mm FL (Chodriyah & Nugroho,
129 2016) reflecting small size dominance. The fish samples came from catches of many kinds of fishing gears, such
130 as purse seine, longline, and trolling. The present study found narrower size distribution, and it could result from
131 less number of samples obtained due to high dependence on local artisanal fishermen who rely on hand-line
132 fishing.

133 The present size range is far below the maximum individual size previously reported reflecting that the
134 mean individual size of *C. hippurus* has been declining. The recovery rate of a population is related to the mortality
135 rate, the closer the mean individual size to the maximum, the lower the mortality rate (ECTF, 2004). The present
136 finding revealed that the dolphinfish population has a high mortality rate. However, so many factors influence
137 fish population availability in the ocean. This condition is supported by Goldstein et al. (2007) that life-history
138 traits are vulnerable to environmental stress and fishing pressure that result in smaller mature fishes as a response
139 for survival. Fish mortality could occur at specific stages and species and the causes may be single or cumulative
140 pressure from a range of sources, such as pollutants, anthropogenic climate change or natural variability (Olsen
141 et al., 2019), and fishing activities. Recruitment patterns with time can influence the population size as well, and
142 therefore, mortality events in the early life stages may have severe and long-lasting effects on the population
143 (Langangen et al., 2017). Climate change is another factor causing changes in fish populations, which can affect
144 the distribution of particular species and the fish susceptibility to particular fishing fleets (Rijnsdorp et al., 2009).
145 This condition could occur because population size has probably fallen below some threshold level of abundance
146 in which the rate of recovery cannot well respond to the fishing rate.

147 This sex deviation is similar to that reported in the western and central Mediterranean (Benseddik et al.,
148 2019; Potoschi et al., 1999; Benseddik et al., 2019) reflecting sex segregation in *C. hippurus* until reaching the
149 mature stage. Mature individuals seem to gather in the same area for spawning and feeding around the rafts so
150 that more females were caught than males. This result also agrees with Perle et al. (2020) and Oxenford (1999)
151 that sex segregation occurs in *C. hippurus* or males are more susceptible to fisheries than females, even though
152 our finding found more females than males. A higher proportion of females from FADs captures could result from
153 greater availability of females, higher natural mortality in males, or differential growth of both sexes (Benseddik
154 et al., 2019). Moreover, males and females show different maturity stages with size class (Table 2). Both sexes
155 show bigger individual sizes than 400 mm FL with more females at mature stages (III and IV). It indicates that
156 males need a bigger size to reach gonad maturity or females reach gonad maturity earlier than males. These data
157 are consistent with Beardsley (1967) that female dolphinfish begin to mature (reach stage II) at about 350 mm FL
158 (about 6–7 months old), 50% are mature at 450 mm FL, and 100% are mature at 550 mm FL, whereas males are
159 mature at a slightly larger size (427 mm FL). Nevertheless, in the Eastern Tunisian Coast, Central Mediterranean,
160 Benseddik et al. (2019) found that the first maturity size of *C. hippurus* occurs at 553 mm FL for females and 605
161 mm FL for males. In the present study, females above 400 mm FL reached maturity stages III and IV. This
162 difference could result from different environmental conditions in localities. It means that 50% of mature
163 individuals that occurs at this size, particularly in the Molucca Sea population, could be set as the minimum legal
164 size of this species to meet the sustainability criteria and avoid economic loss due to fishing immature individuals.
165 The size range of *C. hippurus* caught in the Molucca Sea reflects mature individuals and has mostly passed the
166 size at first maturity. Nevertheless, since fishing is a major factor in reducing size and age at first maturity
167 (McIntyre & Hutchings, 2003) and a decline in age and size at maturity may negatively affect the fish recovery
168 (Hutchings, 2002), it needs to be controlled. The individual size decline of *C. hippurus* far below the maximum
169 size could have indicated a reduced population size and should not be ignored. Earlier maturity can be associated
170 with reduced longevity, increased post-reproductive mortality, and smaller sizes at reproductive age. Populations
171 composed of small individuals will reduce reproductive potential (Scott et al., 1999), increase variance in offspring
172 survival (Hutchings & Myers, 1993), and eventually negatively affect population growth.

173 Mesh size control and escapement could be an alternative to maintain or increase the individual size range
174 or even increase the longevity, and the reproductive potentiality of dolphinfish. Larger fish have higher fecundity
175 and can produce more eggs. So far, commercial purse seiners (< 30 GT) for small pelagic fish have fished any
176 fish schools encountered in the open sea using small mesh sizes. As a result, small yellowfin tuna, skipjack, and
177 dolphinfish are also caught (field observation). Fishing gear separation should be established for commercial

메모 포함[GM67]: "obs." Has been corrected to "observation".

178 small pelagic and large pelagic fisheries to maintain stock availability and prevent individual size decline. This
179 effort limitation could help reduce the risk of population collapse and become one of the remedies to population
180 recovery. Fish population recovery, therefore, requires institutional structures that either entice fishers to leave
181 the business, through expensive buyout schemes of fishing boats and licenses or force them to reduce fishing
182 activity (Hutchings & Reynolds, 2004).

183 The present study has contributed to providing important biological information for future management,
184 especially dolphinfish *C. hippurus* of Molucca Sea. A long-term study on the biology and ecology of this species
185 is required to well describe the population status of *C. hippurus* so that the management policy could be
186 strengthened. The fisheries committee among neighborhood countries that take advantage of the resources should
187 also participate in sustainable resource utilization programs by maintaining the exploitation level and the
188 ecosystem equilibrium.

189
190 **Competing interests**

191 No potential conflict of interest relevant to this article was reported.

192
193 **Funding sources**

194 Not applicable.

195
196 **Acknowledgements**

197 Not applicable.

198
199 **Availability of data and materials**

200 Upon reasonable request, the datasets of this study can be available from the corresponding author.

201
202 **Ethics approval and consent to participate**

203 This article does not require IRB/IACUC approval because there are no human and animal participants.

메모 포함[GM68]: Item for Interest, Acknowledgements, Funding is set in FAS as follow. Some information has been filled in by referring to the contents of the submission system.

Please check if the contents of each item are correct.

204

205

ORCID

206

Silvester Benny Pratasik <https://orcid.org/0000-0002-3765-509X>[207](https://orcid.org/0000-0002-</p></div><div data-bbox=)

[3765-509X](https://orcid.org/0000-0002-3765-509X)

208

Ferdinand Frans Tilaar <https://orcid.org/0000-0003-0448-2513>

209

Meiske Sofie Salaki <https://orcid.org/0000-0001-6442-8856>

210

211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240

References (NLM style; Alphabetical Order)

Bagenal TB. Aspects of fish fecundity. In: Gerking SD (editors). *Methods of Assessment of Ecology of Freshwater Fish Production*. Blackwell, London: Blackwell; 1978; p. 75-101.

Beardsley Jr GL Jr Jr. Age, growth, and reproduction of the dolphin, *Coryphaena hippurus*, in the straits of Florida. *Copeia* 1967;1967:441-451.

Benjamin D, Kurup BM. Stock assessment of Dolphinfish, *Coryphaena hippurus* (Linnaeus, 1758) off the southwest coast of India. *Journal of the Marine Biological Association of India*. 2012;54(1):95-99.

Benseddik AB, Besbes R, Missaoui H, NajaïSE, Jarboui O. Reproductive dynamics and fecundity of *Coryphaena hippurus* (Linnaeus, 1758) in the Eastern Tunisian Coast (Central Mediterranean). *Current Trends in Fisheries and Aquaculture*. 2019;1:23 p.

Brown P, Sivakumaran KP, Stoessel D, Giles A, Green C, Walker T. Carp population biology in Victoria. *Victoria: Marine and Freshwater Resources Institute, Department of Primary Industries, Snobs Creek; 2003. Report No. 56. p. 202. Report No. 56, Marine and Freshwater Resources Institute, Department of Primary Industries, Snobs Creek, Victoria. 2003; p. 202.*

Carlson AK, Rubenstein DI, Levin SA. Linking multiscale fisheries using meta-coupling models. *Front. Mar. Sci.* 2020;7:614; DOI: 10.3389/fmars.2020.00614. 17 p.

Chodrijah U, Nugroho D. Size structure and population parameters of dolphinfish (*Coryphaena hippurus* Linnaeus, 1758) in the Celebes Sea. *BAWAL*. 2016;8(3):147-158 [in Indonesian].

Das M, Dewan S, Debnath SC. Studies on the fecundity of *Heteropneustes fossilis* (Bloch) in a mini pond of Bangladesh Agricultural University, Mymensingh. *Bangladesh Journal of Agricultural Sciences*. 1989; 16:1-6.

Effendie MI. *Biologi Perikanan* (Fisheries Biology). Bogor: Yayasan Dewi Sri. Bogor; 2002; p. 163 p. (In Indonesian).

Farley J, Davies C, Hillary R, Eveson P. Estimating size/age at of southern bluefin tuna. In: *Proceedings of the 18th Meeting of the Scientific Committee; 2013; Canberra, Australia. CCSBTESC/1309/41, 18th Meeting of the Scientific Committee, 2-7 September 2013, Canberra, Australia.* 7 p.

Freshwater C, Holt KR, Huang A-M, Holt CA. Benefits and limitations of increasing the stock-selectivity of Pacific salmon fisheries. *Fisheries Research*. 2020;226:10550-9.

메모 포함[U9]: We have not found this reference. So please make sure it is the right reference.

241 Galimany E, Baeta M, Durfort M, Lleonart J, Ramón M. Reproduction and size at first maturity in the
 242 Mediterranean exploited *Callista chione* bivalve bed. *Scientia Marina*. 2015;79(20):233-242.

243 Goldstein J, Heppell S, Cooper A, Brault S, Lutcavage M. Reproductive status and body condition of Atlantic
 244 bluefin tuna in the Gulf of Maine, 2000–2002. *Mar Biol.* 2007;151:2063-75. DOI 10.1007/s00227-007-
 245 0638-8.

246 Hutchings JA. Life histories of fish. In: P.J.B. Hart P, J.D. Reynolds J, editors. Handbook of Fish biology and
 247 Fisheries, vol. 1. Oxford (United Kingdom): Blackwell Science; 2002. p. 149–174.

248 Hutchings JA, Myers RA. Effect of age on the seasonality of maturation and spawning of Atlantic cod, *Gadus*
 249 *morhua*, in the Northwest Atlantic. *Canadian Journal of Fisheries and Aquatic Sciences* 1993;50:2468–
 250 2474.

251 Hutchings JA, Reynolds JD. Marine fish population collapses: consequences for recovery and extinction risk.
 252 *Bioscience*. 2004;54(4):297-309.

253 Ilkyaz AT, Metin G, Soykan O, Kinacigil HT. Spawning season, first maturity length, and age of 21 fish species
 254 from the Central Aegean Sea, Turkey. *Turkish Journal of Fish and Aquat. Sci.*; 2018;18:211-216. DOI:
 255 10.4194/1303-2712-v18-1-24

256 Indian Ocean Tuna Commission [IOTC]. Review of the statistical data and fishery trends for tropical tunas. In:
 257 *Proceedings of the Fifteenth Working Party on Tropical Tunas; 2013; San Sebastian, Spain. Working Party*
 258 *of Tropical Tunas, Mauritius 24-29 October 2012; 63 pp.*

259 Karna SK, Panda S. Growth estimation and length at maturity of a commercially important fish species i.e.,
 260 *Daysciaena albida* (Boroga) in Chilika Lagoon, India. *Euro. J. Exp. Biol.*; 2011;1(2):84-91.

261 Langangen Ø, Ohlberger J, Stige LC, Durant JM, Ravagnan E, Stenseth NC, et al. Cascading effects of mass
 262 mortality events in Arctic marine communities. *Glob. Change Biol.* 2017;23:283–292. DOI:
 263 10.1111/gcb.13344

264 Lin SJ, Chiang WC, Musyl MK, Wang SP, Su NJ, Chang QX, Ho YS, Nakamura I, Tseng CT, Kawabe R et al.
 265 Movements and habitat use of Dolphinfish (*Coryphaena hippurus*) in the East China Sea. *Sustainability*.
 266 2020;12:5793-16 p.

267 Malone MA, Buck KM, Moreno G, Sancho G. Diet of three large pelagic fishes associated with drifting fish
 268 aggregating devices (DFADs) in the Western Equatorial Indian Ocean. *Animal Biodiversity and*
 269 *Conservation*. 2011;34(2):287-294.

270 McIntyre TM, Hutchings JA. Small-scale temporal and spatial variation in Atlantic cod (*Gadus morhua*) life
 271 history. *Canadian Journal of Fisheries and Aquatic Sciences*. 2003;60:1111–1121.

메모 포함[GM610]: 1) Full name of IOTC has been added.
 Please check if the name is correct or not.

272 **Nandikeswari** R. Size at first maturity and maturity stages of *Terapon jarbua* (Forsskal, 1775) from Pondicherry
 273 Coast, India. *Journal of Fisheries*, 2016; 4(2):385-389.

274 **Olsen** E, Hansen C, Nilsen I, Perryman H, Vikebø F. Ecological effects and ecosystem shifts caused by mass
 275 mortality events on early life stages of fish. *Front. Mar. Sci.* 2019;6:669;
 276 <https://doi.org/10.3389/fmars.2019.00669>. 13 p.

277 **Oxenford** HA. Biology of the dolphinfish (*Coryphaena hippurus*) in the Western Central Atlantic: a review.
 278 *Scientia Marina*, 1999; 63:277-301.

279 **Palko** JBJ, Beardsley GL, Richards WJ. Synopsis of the biological data on dolphinfishes, *Coryphaena hippurus*
 280 Linnaeus and *Coryphaena equiselis* Linnaeus. NOAA Tech Rep NMFS Circ. 1982;443:28 pp.

281 **Peixoto** S, Calazans N, Silva EF, Nole L, Soares R, Frédou FL. Reproductive cycle and size at first sexual maturity
 282 of the white shrimp *Penaeus schmitti* (Burkenroad, 1936) in northeastern Brazil. *Lat. Am. J. Aquat. Res.*
 283 2018; 46(1):1-9.

284 **Perle** CR, Snyder S, Merten W, Simmons M, Dacey J, Rodriguez-Sanchez R, ~~O'Sullivan J, Ortega-GS et al.~~
 285 Dolphinfish movements in the Eastern Pacific Ocean of Mexico using conventional and electronic tags.
 286 *Animal Biotelemetry*, 2020; 8(30):1530 p.

287 **Potoschi** A, Reñones O, Cannizzaro L. Sexual development, maturity and reproduction of dolphinfish
 288 (*Coryphaena hippurus*) in the western and central Mediterranean. *Sci. Mar.* 1999; 63(3-4):367-372.

289 **Rijnsdorp** AD, Peck MA, Engelhard GH, Möllmann C, Pinnegar JK. Resolving the effect of climate change on
 290 fish populations. *ICES Journal of Marine Science*, 2009; 66:1570-1583.

291 **Roa** R, Ernst B, Tapia F. Estimation of size at sexual maturity: an evaluation of analytical and resampling
 292 procedures. *Fish. Bull.* 1999; 97:570-580.

293 **Queensland East Coast Trochus Fishery [ECTF]**. General Effort Review: Sustainability of permitted species
 294 [Internet]. 2004 [cited 2021 Aug 4?];
 295 https://www.daf.qld.gov.au/_data/assets/pdf_file/0003/76629/StockAssessment-CTrawl-2004-Part9

296 **Schlenker** LS, Failandt R, Stieglitz JD, Lam CH, Hoening RH, Cox GK, Heuer RM, Pasparakis C, Benetti
 297 DD, Paris CB, Grosell Met al. Remote predictions of mahi-mahi (*Coryphaena hippurus*) spawning in the
 298 open ocean using summarized accelerometry data. *Front. Mar. Sci.* 2021; 8:1-18626082.
 299 <https://doi.org/10.3389/fmars.2021.626082>.

메모 포함[GM611]: 1) Full name of ECTF has been added. Please check if the name is correct.
 2) Please provide cited day : [Year Month Day].
 2) URL is not valid. Please provide accessible URL.

300 **Scott** B, Marteinsdottir G, Wright P.- Potential effects of maternal factors on spawning stock—recruitment
301 relationships under varying fishing pressure. *Canadian Journal of Fisheries and Aquatic Sciences*, 1999;
302 56:1882–1890.

303 **Sturges** HA. The choice of a class interval. *Journal of the American Statistical Association*, 1926;21(153):65-66.

304 **Taquet** M, Dagorn L, Gaertner JC, Girard C, Aumerruddy R, Sancho G, ~~Hano-Det~~ al. Behavior of dolphinfish
305 (*Coryphaena hippurus*) around drifting FADs as observed from automated acoustic receivers. *Aquatic*
306 *Living Resources*, 2007;20(4):323-330.

307 **Tesfahun** A. Overview of length-weight relationship, condition factor, and size at first maturity of Nile tilapia
308 *Oreochromis niloticus* (L.) in different water bodies of Ethiopia: A Review. *Greener Journal of*
309 *Biological Sciences*, 2018; 8(3):021-028.

310 **Udupae** KS. Statistical method of estimating the size at first maturity in fishes. *Univ. Agricult. Sci. College of*
311 *Fish., Mangalore, India Fishbyte*, 1986; 4(2):8-10.

312 **Whitney** NM, ~~Schwieterman GD~~, Taquet M, Brill RW, Girard C, Schwieterman GD, Dagorn L, ~~Holland K~~
313 al, ~~Girard C~~. Swimming depth of dolphinfish (*Coryphaena hippurus*) associated and unassociated with
314 fish aggregating devices. *Fishery Bulletin— National Oceanic and Atmospheric Administration*, 2016;
315 114(4):426-434.

316

317

318

319 **Table 1. Gonad maturity characteristics**

Maturity stage	Note	Female	Male
I	Immature	Small ovary up to $\frac{1}{4}$ $\frac{1}{2}$ the length of the body cavity. It is translucent. Oocyte does not appear.	The testis is small up to $\frac{1}{4}$ $\frac{1}{2}$ the length of the body cavity. It is whitish.
II	Maturing	The ovary is about half the length of the body cavity. It is orange and translucent, and the oocyte cannot be seen by the naked eye.	The testis is about $\frac{1}{4}$ $\frac{1}{2}$ the length of the body cavity. It is white and about symmetrical.
III	Ripening	The ovary is about $\frac{2}{3}$ the length of the body cavity. Ovary yellow-orange, oocyte appears. Ovary with blood vessels on the surface. No transparent eggs or translucent, eggs are still dark.	The testis is about $\frac{2}{3}$ the length of the body cavity.
IV	Ripe	The ovary is about $\frac{2}{3}$ up to full of the body cavity. The ovary is orange-pink with blood vessels on the surface, and eggs are apparent.	The testis is about $\frac{2}{3}$ up to fulfilling the body cavity. It is white-soft cream.
V	Spent	Ovary shrinks down to $\frac{1}{4}$ $\frac{1}{2}$ the body cavity. Wall is thick. There may be dark and mature eggs in the ovary that disintegrate from absorption, dark or translucent.	Testis shrinks down to $\frac{1}{4}$ $\frac{1}{2}$ the body cavity. Wall is thick. The testis is soft.

320

321

322 **Table 2. Gonad maturity stage of *Coryphaena hippurus* recorded in this study**

323 **Table 2. Gonad maturity stage-**


Size class (mm)	Gonad Maturity maturity Stage-stage (N-n = 50)							
	I		II		III		IV	
	Male	Female	Male	Female	Male	Female	Male	Female
400-449	0	0	4	0	3	3	0	0
450-499	1	0	0	0	1	6	0	3
500-549	1	0	3	0	0	7	1	1
550-599	0	0	0	0	0	8	1	1
600-649	0	0	0	0	0	2	0	0
650-699	0	0	0	0	1	2	1	0

메모 포함[GM612]: There are two titles for Table 2 in the text.
Please select one.

324

325

326



327

328 **Figure Fig. 1. Length frequency distribution of dolphinfish *Coryphaena hippurus*.**

329 **Fig. 1. Length frequency of *Coryphaena hippurus* caught in the Molucca Sea.**

메모 포함[GM613]: There are two titles for Fig. 1 in the text.
Please select only one title.

330

331

332

333