

Glasgow Coma Scale Improvement after Lidocaine Infusion in Moderate Traumatic Brain Injury

by Diana Ch Lalenoh 2

Submission date: 08-Jun-2022 01:18PM (UTC+0700)

Submission ID: 1852776760

File name: AFTER_LIDOCAINE_INFUSION_IN_MODERATE_TRAUMATIC_BRAIN_INJURY.pdf (808.15K)

Word count: 6192

Character count: 33072

Glasgow Coma Scale Improvement after Lidocaine Infusion in Moderate Traumatic Brain Injury

Diana Christine Lalenoh^{1*} and Dewi Yulianti Bisri²

¹Anesthesiology Intensive Care Department, Medical Faculty of Sam Ratulangi University, Indonesia

²Anesthesiology and Intensive Care Department, Medical Faculty of Universitas Padjadjaran, Indonesia

Abstract

Since the Glasgow Coma Scale was developed 40 years ago it has been accepted throughout the world as a method for assessing impaired consciousness.

Forty moderate TBI patients were simple randomly divided into 2 groups: group A take normal saline infusion without lidocaine as control group (N=20), and group B take normal saline with lidocaine at rate 1 mg/kg/BW infusion as treated groups (N=20). Serum samples were collected at before induction and 2 hours after giving infusion treatment to measure serum IL-6 and PLA2 level, and lidocaine serum concentration. GCS were measured before induction and after 24 hours surgery. Data were analyzed using analytic statistical with paired t test to measure serum IL-6, PLA2, & lidocaine serum level before induction and after 2 hours. The correlation of serum IL-6, PLA2 and GCS was measure with Spearman correlation. A P value less than 0.05 were considered significant.

The result are both serum IL-6 and PLA2 levels were significantly lower in saline with lidocaine infusion group ($p < 0.005$) with Mann Whitney and Dependent t test statistical analyses. The lower serum IL-6 and PLA2 levels, the higher GCS value. GCS measurement was significantly higher in lidocaine infusion group ($p < 0.05$) with dependent t test. The decrease in serum IL-6 and PLA2 was correlated with the dosage of lidocaine. Medium doses of lidocaine achieved the optimum decreases in the serum IL-6 and PLA2 level. Therefore, inhibition of the composition and secretion of IL-6 and PLA2 levels by Lidocaine might be one of the mechanisms involved in decreasing inflammatory reaction and cellular damage in brain cells, and may be responsible for the neuroprotective effect after TBI. Glasgow Coma Scale was improving after infusion with lidocaine 1 mg/ kg/ hr during traumatic brain injury surgery by decreasing IL-6 and PLA2 plasm.

27

Keywords: Glasgow coma scale; Traumatic brain injury; Lidocaine

Introduction

The Glasgow Coma Scale is an integral part of assessing levels of consciousness. It uses a simple standardized approach. The scale has been revised to make sure it remains an accurate tool. The overall coma score should not be used to convey clinical findings. The scale can be used with children who are over 5 years old part of the care of patients with acute brain injury, head trauma, intracranial haemorrhage and many other causes. The GCS reflects the initial severity of brain dysfunction, while serial assessments demonstrate the evolution of the injury. Each is crucial for decision making. The GCS is also a guide to prognosis and an essential tool for research studies. Four decades after its introduction, the GCS has gained worldwide acceptance. It is now employed in more than 80 countries, has been translated into more than 60 languages and there are more than 18,000 references to its use [1-4].

Head injury or Traumatic Brain Injury (TBI) is defined as a disorder non degenerative (not aging) and non-congenital (not inherited) that occurs in the brain due to the mechanical strength of the outside/trauma, risk of causing disruption temporarily or permanently in the case of noble function (cognitive), physical, or psychosocial functioning, with accompanying reduction in or loss of consciousness [5]. So far have not found a standard pharmacologic therapy for a primary brain injury. Treatments currently available only to limit or reduce the occurrence of secondary brain injury that would enable the various series of the ischemic cascade and release of proinflammatory cytokines and free radicals product that determine the effects of a brain injury later. This is done with the approach of a series of procedures and stages of brain protection [6].

Local anesthetic lidocaine is one of the drugs that are used as adjuvants in neuroanesthesia, which also affects brain protection. Clinical effect is determined by the dose of lidocaine. At the recommended clinical dose (1.5 mg/kg), lidocaine can reduce the damage of ischemic [7]. Intravenous lidocaine has been known to have a protective effect because the brain works by inhibiting sodium channels, improve recovery through deceleration and a reduction in anoxic depolarization/ischemic and reduce the influx of sodium [8-10].

Several studies conducted to evaluate the outcome of TBI choices of the level of consciousness by using a scoring system on the Glasgow Coma Scale (GCS). Rate GCS can describe the effect of the power of brain injury in the form of compression power, cut outs, or secondary injuries including a series of inflammation that occurs in a brain injury [11-13]. Glasgow Coma Scale (GCS) score is measured on three parameters that is the eye, verbal and motor. Where the highest score was 15 and the lowest value is 3. Determination of brain injury in this study was based on the GCS score 9-12 (moderate TBI).

In addition to GCS, monitoring ICP can also be used for prognostic outcome of a TBI [14]. In one study it was found that GCS determine

***Corresponding author:** Diana Christine Lalenoh, Anesthesiology and Intensive Care Department, Medical Faculty of Sam Ratulangi University, Indonesia, Tel: +6281340601366, +6281340242469; E-mail: diana.lalenoh@yahoo.com

Received July 20, 2016; Accepted August 25, 2016; Published August 31, 2016

Citation: Lalenoh DC, Bisri DY (2016) Glasgow Coma Scale Improvement after Lidocaine Infusion in Moderate Traumatic Brain Injury. J Anesth Clin Res 7:659. doi:10.4172/2155-6148.1000659

Copyright: © 2016 Lalenoh DC, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the prognostic value by measuring the Odds Ratio (OR), where the value of GCS more than 8 when the first patient received indicating outcomes 6.58 times better than the value the GCS 8 downward (OR 6.58; 95% CI 1.87 to 23.14), as observed in a guideline for TBI [15].

In the TBI can be increased proinflammatory cytokines IL-6 and PLA2 [16-20]. Increased IL-6 and PLA2 is signified their cerebral ischemia after TBI [21-30]. Cerebral ischemia caused by TBI can cause cognitive impairment [31-35]. Energy deficit due to supply O₂ lower than the need would cause cognitive impairment [36-43]. The more severe the TBI, the lower the score of GCS. TBI more light, the higher the score GCS [44].

The GCS carries valuable information about the neurological status of patients and constitutes an element of surveillance of their evolution. Yet, by no means should it replace a thorough neurological examination. The same is true for a number of other tools to assess level of consciousness that have been brought forward the previous years. Nonetheless, none of these seems able to replace the GCS. Moreover, even though it was designed for the evaluation of severe TBI, the GCS is currently used in assessment of coma due to any etiology. However, full knowledge of this scale's strengths and limitations is essential in order to assure its proper use [1-4].

Subject and Methodology

Subjects are the patients with moderate TBI (GCS 9-12) with diagnoses epidural hematoma (EDH) undergoing craniotomy surgery with general anesthesia in the operating room emergency room RSU. Prof. R.D. Kandou Manado. Criteria for inclusion are age 17-55 years, ASA physical status I and II, new TBI traumatic experience within a maximum of 48 hours first, no pro-coagulant therapy. The exclusion criteria are patients who experience worsening of traumatic TBI (loss of consciousness or signs of cerebral herniation) during observations, during the observation period decreased general condition that requires intervention by certain drugs that can interact with the type of treatment used and the family refused to participate. Criteria Drop Out are patients TBI traumatic experience complications during administration of intervention in the form of hemodynamic instability during the observation and intervention, surgery craniotomy with operating time of more than 3 hours.

Using simple random method, samples divided into 2 groups. First group takes sodium chloride infusion after intubation, and second group takes sodium chloride with lidocaine infusion 1 mg/kg/hour with syringe pump. All samples take 2 hours of infusion during surgery. Anesthesia maintenance with isoflurane (no more than 1.5 MAC or 2 volume %) and propofol titration by syringe pump 1-2 mg/kg. We give muscle relaxant every 20 minutes with rocuronium 10 mg IV boluses.

Procedures for work

Before being given the treatment, all patients were given nasal cannula oxygenation with 3 liters/min, 1 hour before the start of induction. The subjects were divided into two study groups. The first group (control group) received infusion of Normal Saline 0.9% without lidocaine and the second group (treatment group) received intravenous saline with lidocaine. Each sample was induced with fentanyl 2 mg/kg and propofol 2 mg/kg body weight. Facilities rocuronium 10 mg intubation with 0.9 mg/kg body weight. The patients were divided two groups, the first group was given a bolus of lidocaine 1 mg/kg after intubated followed by infusion of lidocaine 1 mg/kg/h (using a syringe pump), the second group was given a bolus of lidocaine 1 mg/kg after intubated followed by infusion of Normal saline 0.9% (using a syringe pump) at the same speed.

Infusion of lidocaine or Normal Saline was continued for 2 hours while the surgical procedure is still running. Maintenance of anesthesia with isoflurane and propofol titration (syringe pump) 1-2 mg/kg body weight/hour. The type of surgery to be included in the study is limited to craniotomy operation with the operating time of no more than 3 hours, which is calculated from the start until the skin incision to start sewing leather. Inhalation anesthetics (isoflurane) is given no more than 1.5 MAC (2 volume %).

Muscle relaxants are given every 20 minutes with rocuronium 10 mg intravenous bolus. GCS score measure in pre induction period and 24 hour postoperative.

Data processing and analysis

Analisis processing and distribution of data when normal population using statistical tests dependent-test for independent variables numerical scale that lidocaine levels in plasma, but to the data with the population distribution is not normally distributed, used Mann Whitney test. For categorical variables such as the assessment scale used GCS chi-squared test.

Ethical issues

This research is performed without violating ethics, because:

The samples were given lidocaine does not endanger patients because the drug is often used both in the field of neurosurgery as well as in neuroanesthesia in general as a protective agent penumpul brain and sympathetic response.

Patients are given anesthesia and analgesia drugs are also routinely used in anesthesia action.

The study was conducted after obtaining approval from the ethics committee.

Result

Characters of demographic and clinical factors

Of the 40 samples that meet the criteria, divided into treatment and control groups at random; each 20 people per group. With the division of random groups expected demographic and clinical characteristics of research subjects divided equally in both groups or in other words occur homogenization demographic and clinical characteristics between the two groups before treatment, so that the demographic characteristics did not affect differences in the expected effect on both group (Table 1).

Table 1 shows that there are strong negative linier significance ($p < 0.05$) correlation between lidocaine with heart rate, systolic blood pressure, diastolic blood pressure, IL-6, and PLA2. The higher lidocaine concentration, the lower heart rate, systolic blood pressure, diastolic blood pressure, IL-6 and PLA2. There is positive linier correlation between lidocaine concentration with GCS and MMSE.

Table 2 shows correlation analysis between IL-6, PLA2, and GCS. There are strong linier significance correlation ($r = 0.763$, $p < 0.05$) between IL-6 and PLA2. The higher IL-6, the higher PLA2. There are moderate negative linier correlation ($r = -0.508$) between PLA2 with GCS. The higher PLA2, the lower GCS; the higher IL-6, the lower GCS.

Figure 1 shows that there is negative correlation between lidocaine concentration with IL-6; the higher lidocaine concentration, the lower IL-6.

Variables	Lidokain concentration	
	Correlation Coefficient (r)	Significance (p)
HR	$r=-0.701$	$p=0.000$
SBP	$r=-0.823$	$p=0.000$
DBP	$r=-0.788$	$p=0.000$
IL-6	$r=-0.677$	$p=0.000$
PLA2	$r=-0.734$	$p=0.000$
GCS	$r=0.356$	$p=0.012$

Table 1: Correlation between lidocaine with IL-6, PLA2, HR, SBP, and DBP 60 minutes after extubation.

Correlation	Spearman Correlation	
	Correlation Coefficient (r)	Significance (p)
IL-6-CS	$r=-0.298$	$p=0.031$
PLA2-CS	$r=-0.508$	$p=0.000$
IL-6-PLA2	$r=0.763$	$p=0.000$

Table 2: The correlation of IL-6, PLA2 with GCS 60 minutes after extubation.




Figure 1: Correlation between lidocaine with IL-6.

Figure 2 shows there is negative correlation between lidocaineplasmconcentration with PLA2 plasm; the higher lidocaineplasm concentration, the lower PLA2 plasm.

Figure 3 shows there is positive correlation between IL-6 plasm with PLA2 plasm concentration; the higher IL-6 plasm concentration, the higher PLA2 plasm concentration.

Table 3 shows that there is no side effect which are bradycardia and hypotension between control and lidocaine group.

Table 3 shows that heart rate in normal limit during surgery until 60 minute after extubation. Therefore, as diastolic blood pressure. The systolic blood pressure approximately in normal limit, just once less than 90 mmHg but the decrease is no more than 20% from baseline. So there is relative no hypotension effect in this research.

Levels of serum lidocaine

The treatment group was the group that received lidocaine infusion in a liquid, NaCl 0.9% and the control group only received 0.9% NaCl infusion without lidocaine. Two hours post-infusion, examined serum lidocaine levels in both groups. Comparison of serum lidocaine levels between the two groups were tested with independent sample test. The results can be seen in Table 4.

Table 4 shows that serum lidocaine levels 2 hours post-infusion was higher in the treatment group (1901.7 ± 169.6 ng/ml) than the control

group (171.9 ± 75.8 ng/ml). Results of independent t test showed very significant differences ($p=0.000$).

Changes GCS

GCS changes in the control and treatment groups were tested by paired t test. GCS comparison between the two groups before surgery, as well as GCS after extubation and the difference of change in GCS before surgery until after extubation between the two groups were tested by the Mann Whitney test. The results can be seen in Table 5.

Table 5 shows that in the control group there was an increase of 0.1 of GCS (10.1 ± 0.8) to (10.2 ± 0.8) but paired t test results showed no significant changes ($p>0.05$), while the treatment group increased by 1.3 of (10.1 ± 0.6) to (11.4 ± 1.3) and significant ($p<0.05$). Mann Whitney test results showed that the GCS between the two groups prior to surgery showed no significant difference ($p=0.083$), but after two hours of infusion was significantly different ($p<0.001$), as well as differences in changes in GCS, significant ($p<0.05$). Thus, changes in GCS were significantly different between the two groups. GCS increased in the treatment group, whereas the control group did not change.

Figure 4 shows that the same GCS (coincident) before induction of anesthesia, and after a 2 hour infusion of GCS levels increased in the treatment group, while the control group remained or increased slightly.




Figure 2: Correlation between lidokain with PLA2.




Figure 3: Correlation between plasm IL-6 with PLA2 plasm concentration.

Time Observation	Minimal-maximal Heart Rate (x/minute)		Minimal-Maximal SBP (mmHg)		Minimal-Maximal DBP (mmHg)	
	Control	Lidocaine	Control	Lidocaine	Control	Lidocaine
Before surgery	78-105	78-102	115-144	112-147	56-100	59-104
Awal infus	78-100	79-102	112-140	115-145	54-98	61-100
30' infus	78-103	71-100	131-154	92-136	67-100	48-78
60' infus	78-102	70-92	128-152	92-136	78-102	50-72
90' infus	78-101	70-98	128-158	85-126	78-102	52-79
120' infus	78-100	68-90	135-150	92-136	44-103	48-78
Saat ekst	78-100	66-93	132-149	92-136	73-100	48-78
30' post ekst	78-102	65-90	132-150	92-136	72-100	48-78
60' post ekst	78-100	63-91	132-150	92-136	80-100	48-78

Table 3: Distribution of range heart rate, SBP, DBP between control group and lidocaine group.

Level Lidocaine Serum (ng/ml)	Group		Independent Sample t test
	Treatment (n=20)	Kontrol (n=20)	
Mean \pm SD	1901.7 \pm 169.6	171.9 \pm 75.8	p=0.000
Minimal/Maximal	1776-2514	68-388	

Table 4: Comparison of serum lidocaine levels between 0.9% NaCl infusion group (control) and Group NaCl+0.9% Lidocaine (treatment).

Group	GCS			Paired t- test
	Preinduction	2 hours post-operative	Perubahan	
Control (n=20)	(10.1 \pm 0.8) ^a	(10.2 \pm 0.8) ^a	Increase (0.1) ^a	p=0.083
Treatment (n=20)	(10.1 \pm 0.6) ^a	(11.4 \pm 1.3) ^a	Increase (1.3) ^a	p=0.001

Superscript are not different in the same column shows the results of Mann Whitney test was not significant ($p > 0.05$), and if different, Mann Whitney test results showed significant ($p < 0.05$).

Table 5: Comparison of GCS changes between the two groups.




Figure 4: Graph GCS change compared to before the induction of anesthesia until after the 2-hour infusion between control and treatment groups.

Pulse rate changes and average artery pressure

To assess the side effects that may occur, observed on cardiovascular responses (heart rate and mean arterial pressure). Analysis results can be seen in Table 6.

Correlation between GCS improvement with IL-6 and PLA2 decreasing

Table 7 show there are strong correlation ($p=0.000$) between IL-6 plasm decreasing with GCS increasing with coefficient correlation $r=-0.530$. The lower IL-6 plasm concentration, the higher GCS. The lower PLA2 plasm concentration, the higher GCS ($r=-0.351$ and $p=0.031$) ($p < 0.05$).

Discussion

General characteristics

General data of patients including age, height, weight, BMI, initial

pressure systolic, pressure early diastolic blood pressure on average, respiratory rate, heart rate, SpO_2 , core temperature, blood sugar levels, and GCS showed no difference significant between the two groups, which means both relatively homogenous groups.

The average patient age was 30 years; mean traumatic effect will be the same as its value is the same GCS. In contrast to when one group over the age of 60 years. Although it weighs the same primary injury but because of different brain tissues, the therapeutic effect will be different.

Lidocaine effect on intracranial component

Perhaps the most widely used of this intravenous lidocaine, administered before rapid sequence intubation. Although the exact mechanism by which lidocaine works is unclear, it is thought to be related to its ability to blunt the increase in pulse rate and blood pressure associated with laryngoscopy and tracheal intubation, which subsequently cause an increase in intracranial pressure. Lidocaine has also been shown to reduce cerebral blood flow and cerebral vascular resistance. Finally, lidocaine as a sodium channel blocker, may decrease cerebral metabolism and stabilize neuronal membranes, which may lead to a decrease in secondary brain injury in traumatic brain injury. Evidence suggests that using intravenous lidocaine 2 to 3 minutes before rapid sequence intubation is safe and may help attenuate the increase in intracranial pressure associated with intubation in the traumatically brain-injured patient, potentially improving long term outcome. An intracranial pressure greater than 20 mmHg and occurring within the first 72 hours of patient care is associated with an increased morbidity and mortality [45-47].

Each channel for ion transport pathways in the cell membrane of neurons can only be traversed by a certain ion. In the resting, state of cells no stimulation to the cell membrane potential in a state of rest. In the circumstances that led to the opening of voltage-gated channels (usually very fast) changes in the membrane potential and this is what

causes the onset of electrical signals called action membrane potential. There are two types of channels in addition to the canals mentioned above, namely that can open and close. Channels that provide an answer to their chemical signals responsible for the formation of an electrical signal, called a ligand-gated channels. This channel is opened when there is stimulation of extracellular molecules. Some are some channels that respond to molecules in the cell, for example of a molecule "second messengers" such as Ca^{2+} or other molecules such as cyclic GMP, and G protein subunits that are activated by cell surface receptors. The action potential is a cycle of depolarization, hyperpolarization and return to a resting state. Specifically, the electrical changes caused by the Na^+ channels and voltage-gated K^+ that open and close in response to changes in membrane potential [45-47].

In the process, the Krebs cycle takes a number of enzymes found in the mitochondrial matrix that will give the final result of CO_2 . During this process the ion H^+ will be released that will be captured by coenzyme nicotinamide adenine dinucleotide (NAD). Electrons are derived from the hydrogen moving through some respiratory enzymes (flavoproteins and cytochromes) that eventually combine with oxygen to form protons from water molecules (H_2O). Of the latter is to be obtained by the energy stored in the bonds of ATP are derived from ADP. Being necessary enzymes bound to the inner mitochondrial membrane wall composed of sections called inner membrane sub-units (unit globular). Each subunit consists of a round building with a diameter of 9 nm which is connected with the handle along the 3.5 to 4.5 nm [48,49].

A chemical signal is delivered from outside the cell through a receptor on the cell surface, and then sent to the cell nucleus. This process is very complex because it involves a variety of chemical and physical properties of a signal protein, which is not another because of changes in the electrical charge of protein molecules [46]. In a tissue, damage will occur inflammatory process characterized by increased proinflammatory cytokines IL-1B, IL-6, TNF α . Lidocaine effects that reduce cell death as a result of barriers to the Na^+ channels will cause reduced the inflammation process.

Magnesium-lidocaine combination administered according to the above regimen is safe and well tolerated after severe head injury. Patient in the magnesium-lidocaine group showed a more favorable outcome and lower mortality rate when compared to the control group. Lidocaine infusion (IV, 3 mg/kg) triggers a 5-minute 10% reduction of cerebral O_2 consumption within 2 to 3 minutes [49]. Lidocaine reduces in a dose dependent manner brain metabolism, suppresses synaptic transmission and stabilizes membranes by blocking pathologic K^+ efflux, Na^+ channels, and demand for Na^+ - K^+ transport [50].

Classification of traumatic brain injury

Traumatic brain injury division based on a few things in between based on the mechanism of occurrence, based on the severity (severity), and based on its morphology. Based on the mechanism of injury, the TBI is divided into: blunt injury and injury penetrans. Based on the severity (severity) TBI divided based on the criteria of the Glasgow Coma Scale (GCS), namely: mild TBI (GCS 14-15); moderate TBI (GCS 9-13); and severe TBI (GCS 3-8) in patients examined [49].

Based on the morphology TBI is divided into: 1. Focal injuries, ie injuries identified suitable areas affected; divided into: Cerebral contusion, extradural hemorrhage, scalp lacerations, fractures calvaria, subarachnoid hemorrhage, subdural Hemorrhage; 2. Diffuse injuries, ie injuries affecting the whole brain tissue, consisting of: Concussion and Brain Injury Diffuse axonal (Table 8) [47,50].

Glasgow coma scale

Glasgow Coma Scale was first introduced by Teasdale and Jennett as the standard method for evaluating health care givers and describe the degree of alteration of consciousness or coma in patients who suffered a head injury. Glasgow Coma Scale to test three different components, namely: open your eyes, verbal response, and motor response. In each category there is a range of ratings to determine the specific components. Each observation is connected with the scores recorded by health care providers as a standard classification of the patient's

Time of Observation	Minimal-Maximal Heart Rate (x/minutes)		Minimal-Maximal MAP (mmHg)	
	Control	Treatment	Control	Treatment
Before surgery	78-105	78-102	96-127	97-130
60' infusion	78-102	70-92	114-130	78-113
120' infusion	78-100	68-90	109-132	77-114

MAP: Mean Arterial/Blood Pressure

Table 6: The range of distribution of heart rate and mean arterial pressure in the control and treatment.

The correlation between variables	Spearman Correlation	
	Coefficient correlation (r)	Significance (p)
IL-6 plasm decreasing with GCS increasing	$r=-0.530$	$p=0.000$
PLA2 plasm decreasing with GCS increasing	$r=-0.351$	$p=0.013$
IL-6 plasm decreasing with PLA2 plasm decreasing	$r=0.453$	$p=0.002$

Table 7: The correlation between IL-6 and PLA2 plasm decreasing with GCS increasing.

	Mild TBI	Moderate TBI	Severe TBI
Beginning GCS value	1-11	9-13	3-8
Presentation	80	10	10
Abnormality of CT Scan (%)	5-15	30-50	60-90
Demand of Neurologic intervention (out of ICP Monitoring)	1-3	5-30	30-50
Mortality (%)	<1	10-15	30-50
Good Functional Outcome/GEO (%)	>90	20-90	<20

Table 8: There is also a division of TBI based on the outcome produced.

level of consciousness. Components eyes open there are 4 scale ranging from no response (score 1) through the eyes open spontaneously (score 4). Motor component No 6 scale, ranging from no response to the movement (score 1) to comply with orders observer movement (score 6). No verbal components 5 components, ranging from no verbal response at all (score 1) to be able to communicate the appropriate / right orientation (score 5). From each category are then summed to be rated as GCS [49].

Traumatic brain injury is a heterogeneous condition in terms of aetiology, severity, and outcome. The most useful classification of severity is based on the level of consciousness as assessed by the Glasgow Coma Scale after resuscitation. The GCS comprises the sum score of the value from three components: eye, motor, and verbal scales. However, factors such as hypoxia, hypotension, and alcohol intoxication can affect GCS, leading to diagnostic confusion. Therefore, the patient should be resuscitated and reversible causes corrected before GCS assessment. The ability to assess eye responses is influenced by sedative agents or tracheal intubation, leading some to suggest the use of the motor score alone.

Score GCS can describe the effect of the power of brain injury in the form of compression power, cutouts, or secondary injuries including a series of inflammation that occurs in a brain injury [11-13]. In addition to GCS, monitoring ICP can also be used for prognostic outcome of TBI [10]. In one study it was found that GCS determine the prognostic value by measuring the Odds Ratio (OR), where the value of GCS more than 8 when the first patient received indicating outcomes 6.58 times better than the value of GCS 8 downward (OR 6.58; 95% CI 1.87 to 23.14), as observed in a guideline for TBI [15].

Neurological injury progresses over hours and days, resulting in a secondary injury. Inflammatory and neurotoxic processes result in vasogenic fluid accumulation within the brain, contributing to raise intracranial pressure, hypoperfusion, and cerebral ischaemia. Much of secondary injury may be amenable to intervention, as almost one-third of patients who die after a TBI will talk of obey commands before their death. Secondary injury also occurs as a result of further physiological insults. Hypoxia, hypotension, hypercapnia or hypocapnia, hyperglycemia or hypoglycaemia has all been shown to increase the risk of secondary brain injury. Overall those processes involve and influence the value of GCS [45-49].

Conclusion

Glasgow Coma Scale was improving after infusion with lidocaine 1 mg/kg/hr during traumatic brain injury surgery by decreasing IL-6 and PLA2 plasm.

References

- Teasdale G, Maas A, Lecky F, Manley G, Stocchetti N, et al. (2014) The Glasgow Coma Scale at 40 years: standing the test of time. *Lancet Neurol* 13: 844-854.
- Middleton PM (2012) Practical use of the Glasgow Coma Scale; a comprehensive narrative review of GCS methodology. *Australas Emerg Nurs J* 15: 170-183.
- Forty Years of Updating the Glasgow Coma Scale. Nursing practice review neurology. *Nursing times.net*.
- Matis G, Birbilis T (2008) The Glasgow Coma Scale—a brief review. Past, present, future. *Acta Neurol Belg* 108: 75-89.
- Donnelly JP, Donnelly K, Grohman KK (2005) A multi-perspective concept mapping study of problems associated with TBI. *Brain Inj* 19: 1077-1085.
- Papadakis M, Buchan AM (2006) Translational vehicles for neuroprotection. *Biochem Soc Trans* 34: 1318-1322.

- Kass IS, Cottrell JE (2001) Pathophysiology of brain injury: In: *Anesthesia and Neurosurgery*. (4th edn), St Louis, Mosby 69-79.
- Kass IS, Cottrell JE, Baiping L (2010) Brain metabolism, the pathophysiology of brain injury, and potential beneficial agents and techniques: In: *Cottrell and Young's Neuroanesthesia*. (5th edn), Philadelphia, Mosby Elsevier, Inc 5-9.
- Mitchell SJ, Merry AF, Frampton C, Danies E, Grieve D, et al. (2009) Cerebral protection by lidocaine during cardiac operations: a follow-up study. *Ann Thorac Surg* 87: 820-825.
- Puljak L, Kojundzic SL, Hogan QH, Sapunar D (2009) Lidocaine injection into the rat dorsal root ganglion causes neuroinflammation. *Anesth Analg* 108: 1021-1026.
- Mohammad MM (2006) Neuroprotection: Concept and Updating Knowledge. Faculty of Medicine Minia University 2006.
- Peck S (2005) Clinical Guideline for The Care and Treatment of Older People with Delirium in a General Hospital Setting. 2nd ed. Isle of Wright Healthcare. NHS trust. Older Persons Mental Health Service.
- Moppett IK (2007) Traumatic brain injury: assessment, resuscitation and early management. *Br J Anaesth* 99: 18-31.
- Smith M (2007) Monitoring intracranial pressure in traumatic brain injury. *International Anaesthesia Research Society* 106: 240-249.
- Palmer S, Badar MK, Qureshi A, Palmer J, Shaver T, et al. (2001) The impact on outcomes in a community hospital setting of using the AANS Traumatic Brain Injury Guidelines. *J Trauma* 50: 657-664.
- Hengenroeder GW, Moore AW, McCoy JP, Sammel L, Ward III NH, et al. (2010) Serum IL-6: A candidate biomarker for intracranial pressure elevation following isolated TBI. *Research. Journal of NeuroInflammation* 19: 1-13.
- Rahmani M, Faris EA, Benabdeljil M, Aidi S (2013) Cognitive disturbances in Sneddon and antiphospholipid syndrome: In: *Neuroscience*. Croatia.
- Dyail SC (2010) Amyloid-beta peptide, oxidative stress, and inflammation in Alzheimer's disease: potential neuroprotective effects of Omega-3 polyunsaturated fatty acids. *Review Articles: In: International Journal of Alzheimer's Disease* 1-10.
- Ortega FJ, Vidal-Taboada JM, Mahy N, Rodriguez MJ (2010) Molecular mechanisms of acute brain injury and ensuing neurodegeneration: In: *Brain Damage. Bridging Between Basic Research and Clinics* 163-186.
- De Jongh RF, Vissers KC, Meert TF, Booij LH, De Deyne CS, et al. (2003) The role of interleukin-6 in nociception and pain. *Anesth Analg* 96: 1096-1103.
- Nitro C, Kamada H, Endo H, Niizuma K, Myer DJ, et al. (2008) Role of the p38 mitogen-activated protein kinase/ cytosolic phospholipase A2 signaling pathway in blood-brain barrier disruption after focal cerebral ischemia and reperfusion. *J Cereb Blood Flow Metab* 28: 1686-1696.
- Cherubini A, Polidori C, Benedetti C, Ercolani S, Senin U, et al. (2013) Institute of Gerontology and Geriatrics. Adibhatla RM, Hatcher JF, Dempsey RJ (2002) Cytokine: neuroprotective mechanisms in cerebral ischemia. *J Neurochem* 80: 12-23.
- Lee JM, Grabb MC, Zipfel GJ, Choi DW (2000) Brain tissue responses to ischemia. Perspective series. *Tissue responses to ischemia. J Clin Invest* 80: 12-23.
- Siebert FE, Traystman RJ, Martin LJ (1997) Delayed neuronal death after global incomplete ischemia in dogs is accompanied by changes in phospholipase C protein expression. *Journal of Cerebral Blood Flow and Metabolism* 17: 527-533.
- Wei EP, Lamb RG, Kontos HA (1982) Increased phospholipase C activity after experimental brain injury. *J Neurosurg* 56: 695-698.
- Rapoport MJ, McCullagh S, Shammi P, Feinstein A (2005) Cognitive impairment associated with major depression following mild and moderate traumatic brain injury. *J Neuropsychiatry Clin Neurosci* 17: 61-65.
- Nito C, Kamada H, Endo H, Niizuma K, Myer DJ, Chan PH, et al. (2008) Role of the p38 mitogen-activated protein kinase/ cytosolic phospholipase A2 signaling pathway in blood-brain barrier disruption after focal cerebral ischemia and reperfusion. *Journal of Cerebral Blood Flow & Metabolism* 28: 1686-1696.
- Lee JM, Grabb MC, Zipfel GJ, Choi DW (2000) Brain tissue responses to ischemia. Perspective Series. *Tissue Responses to Ischemia. The Journal of Clinical Investigation* 80: 723-731.
- Ravetty, Rosso OA, Benetta R, Mocato P (2010) Uncovering molecular biomarkers that correlates cognitive decline with the changes of hippocampus gene expression profiles. *PLOS one* 5: 1015-1021.

30. Mathuranath PS, Chenan JP, Mathew R, George A, Alexander A, et al. (2007) Minute state examination and the Addenbrooke's cognitive examination: effect of education and norms for a multicultural population. *Neurology India* 55: 106-110.
31. Godefroy O, Fickl A, Rovssel M, Auribault C, Bugnicourt JM, et al. (2011) Is the Montreal Cognitive Assessment superior to the Mini Mental State Examination to detect Post Stroke Cognitive Impairment? A study with neuropsychological evaluation. *Stroke* 42: 1712-1716.
32. Firefighter Written Examination Study Guide. New Jersey Civil Service Commission Firefighter Cognitive Test Study Guide. Diakses dari 2013.
33. Lamarre CJ, Patten SB (1994) A Clinical Evaluation of The Neurobehavioural Cognitive States. Examination in a General Psychiatric In Patient Population. Diakses dari.
34. Minimental State Exam (2010) Diakses dari.
35. Adibhatla RM, Hatcher JF (2008) Altered lipid metabolism in brain injury and disorders. *Subcell Biochem* 49: 241-268.
36. Robertson CL, Scafidi S, McKenna MC, Fiskum G (2009) Mitochondrial mechanisms of cell death and neuroprotection in pediatric ischemic and traumatic brain injury. *Exp Neurol* 218: 371-380.
37. Bramlett HM, Dietrich WD (2004) Pathophysiology of cerebral ischemia and brain trauma: similarities and differences. *J Cereb Blood Flow Metab* 24: 133-150.
38. Mangat HS (2012) Severe traumatic brain injury. *Continuum (Minneapolis)* 18: 532-546.
39. TBI Journal Watch RSS (2008) Northeast Center for Special Care. Diakses dari 2013.
40. Kinnunen KM, Greenwood R, Powell JH, Leech R, Hawkins PC, et al. (2011) White matter damage and cognitive impairment after traumatic brain injury. *Brain* 134: 449-463.
41. Till C, Colella B, Verwegen J, Green RE (2008) Postrecovery cognitive decline in adults with traumatic brain injury. *Arch Phys Med Rehabil* 89: S25-34.
42. Steinmon KJ, Gorn Tempini ML, Glidden DV, Kramer JH, Miller SP, et al. (2009) Neonatal watershed brain injury on MRI correlates with verbal IQ at four years. *Pediatrics* 123: 1025-1030.
43. Brown A, Moor PS (2010) TBI Craniotomy-Supratentorial Craniotomy. Part I Case: In: *Case Studies in Neuroanesthesia and Neurocritical Care*. Cambridge.
44. Chen J, Dohis T, Nozauray B (2002) The Inhibitory effect of local anesthetics on bradykinin-induced phospholipase activation in rat pheochromocytoma PC12 cells. *Anesth Analg* 95: 88-97.
45. Subowo (2011) *Biologi sel*. (6th edn). CV. Sagung Seto; 285-287.
46. Curtis K, Chong S, Mitchell R, Newcombe M, Black D, et al. (2011) Outcomes of severely injured adult trauma patients in an Australian health service: does trauma center level make a difference? *World J Surg* 35: 2332-2340.
47. Heegaard W, Biros M (2007) Traumatic brain injury. *Emerg Med Clin North Am* 25: 655-678, viii.
48. McNett M (2007) A review of the predictive ability of Glasgow Coma Scale scores in head-injured patients. *J Neurosci Nurs* 39: 68-75.
49. Lalenoh DC, Bisri T, Yusuf I (2014) Brain Protection Effect of Lidocaine Measured By Interleukin-6 and Phospholipase A2 Concentration in Epidural Haematoma with Moderate Head Injury Patient. *J Anesth & Crit Care* 1-8.
50. Bisri T (2012) *Buku Cidera Otak Traumatik*.

Glasgow Coma Scale Improvement after Lidocaine Infusion in Moderate Traumatic Brain Injury

ORIGINALITY REPORT

18%
SIMILARITY INDEX

7%
INTERNET SOURCES

10%
PUBLICATIONS

9%
STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|-----|
| 1 | Submitted to The Maldives National University
Student Paper | 1 % |
| 2 | BIAN, Xiao-xing, Xue-song YUAN, and Chuan-ping QI. "Effect of Recombinant Human Erythropoietin on Serum S100B Protein and Interleukin-6 Levels After Traumatic Brain Injury in the Rat", Neurologia medico-chirurgica, 2010.
Publication | 1 % |
| 3 | Submitted to University Of Tasmania
Student Paper | 1 % |
| 4 | Submitted to 97324
Student Paper | 1 % |
| 5 | Sergio Canavero, Vincenzo Bonicalzi, Paolo Narcisi. "Safety of magnesium-lidocaine combination for severe head injury: the turin lidomag pilot study", Surgical Neurology, 2003
Publication | 1 % |

6	"Euroanaesthesia 2004: Joint Meeting of the European Society of Anaesthesiologists and European Academy of Anaesthesiology Lisbon, Portugal, 5–8 June 2004", European Journal of Anaesthesiology, 09/14/2005 Publication	1 %
7	Submitted to University of Wales Swansea Student Paper	1 %
8	journals.plos.org Internet Source	1 %
9	Submitted to Higher Education Commission Pakistan Student Paper	1 %
10	Molly McNett. "A Review of the Predictive Ability of Glasgow Coma Scale Scores in Head-Injured Patients", Journal of Neuroscience Nursing, 04/2007 Publication	1 %
11	blognyaperawatcerdas.blogspot.com Internet Source	1 %
12	"Continuous intravenous perioperative lidocaine infusion for postoperative pain and recovery", Cochrane Database of Systematic Reviews, 2015. Publication	1 %
13	Submitted to Oxford Brookes University Student Paper	1 %

14

Y Purnamasari, Tien, Arimaswati, A N K Syarifin, A Ali, N Massi. "Genotype and Serotype Identification of Hepatitis B Virus in Chronic Hepatitis B Patients Treated with Telbivudine", Journal of Physics: Conference Series, 2019

Publication

1 %

15

I.K. Moppett. "Traumatic brain injury: assessment, resuscitation and early management", British Journal of Anaesthesia, 2007

Publication

<1 %

16

www.ncbi.nlm.nih.gov

Internet Source

<1 %

17

Submitted to California Lutheran University

Student Paper

<1 %

18

biology.ejournals.ca

Internet Source

<1 %

19

Submitted to La Trobe University

Student Paper

<1 %

20

Submitted to University of New England

Student Paper

<1 %

21

ELINORE F. McCANCE-KATZ, CARL L. HART, BETH BOYARSKY, THOMAS KOSTEN, PETER JATLOW. "Gender Effects Following Repeated

<1 %

Administration of Cocaine and Alcohol in Humans", Substance Use & Misuse, 2009

Publication

22

Submitted to University of Wales, Bangor

Student Paper

<1 %

23

"Lidocaine versus magnesium sulfate infusion during isoflurane anesthesia for brain tumor resection, effect on minimum alveolar concentration reduction guided by bispectral index: a prospective randomized controlled trial", Signa Vitae, 2021

Publication

<1 %

24

Tjokorda Gde Agung Senapathi, Made Wiryana, I Gusti Ngurah Mahaalit Aribawa, Christopher Ryalino. "Bispectral index value correlates with Glasgow Coma Scale in traumatic brain injury patients", Open Access Emergency Medicine, 2017

Publication

<1 %

25

Submitted to University of Birmingham

Student Paper

<1 %

26

Katja Heikkinen. "Prostatectomy patients' postoperative pain assessment in the recovery room", Journal of Advanced Nursing, 12/2005

Publication

<1 %

- | | | |
|----|---|------|
| 27 | on behalf of the PENTA study group.
"Assessing the level of consciousness in children: A plea for the Glasgow Coma Motor subscore", Resuscitation, 200802
Publication | <1 % |
| 28 | archive.org
Internet Source | <1 % |
| 29 | coek.info
Internet Source | <1 % |
| 30 | www.tandfonline.com
Internet Source | <1 % |
| 31 | Kumiko Okamura. "Glasgow Coma Scale flow chart: a beginner's guide", British Journal of Nursing, 2014
Publication | <1 % |
| 32 | Giscard K. Kaptso, Nicolas Y. Njintang, Marie Goletti M. Nguemtchouin, Athanasius F. Amungwa et al. "Characterization of Morphology and Structural and Thermal Properties of Legume Flours: Cowpea (Vigna unguiculata L. Walp) and Bambara Groundnut (Vigna subterranea L. Verdc.) Varieties", International Journal of Food Engineering, 2016
Publication | <1 % |
| 33 | Roshaslina Rosli, Maw Pin Tan, William Keith Gray, Pathmawathi Subramanian, Ai-Vyrn | <1 % |

Chin. "Cognitive assessment tools in Asia: a systematic review", International Psychogeriatrics, 2015

Publication

34

docksci.com

Internet Source

<1 %

35

docplayer.net

Internet Source

<1 %

36

"Abstracts of Posters Presented at the International Anesthesia Research Society 78th Clinical and Scientific Congress Tampa, FL March 27-31, 2004", Anesthesia & Analgesia, 2004

Publication

<1 %

37

&NA;. "Abstract", Anesthesia & Analgesia, 1989

Publication

<1 %

38

Yoshinari Oyanagi. "Enhanced expression of interleukin-6 in chronic hepatitis C", Liver International, 12/1999

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On