

# SERUM LIPID PROFILE OF BROILER CONSUMING RATION CONTAINING LAURIC ACID AND FEED FIBER

*by Jola Londok 6*

---

**Submission date:** 15-Sep-2022 12:19PM (UTC+0700)

**Submission ID:** 1900252099

**File name:** SERUM\_LIPID\_PROFILE\_OF\_BROILER\_CONSUMING\_RATION.pdf (352.5K)

**Word count:** 3469

**Character count:** 16058

## SERUM LIPID PROFILE OF BROILER CONSUMING RATION CONTAINING LAURIC ACID AND FEED FIBER

Jola Josephien Mariane Roosje LONDOK<sup>1\*</sup>, John Ernst Gustaaf ROMPIS<sup>2</sup>

12

<sup>1</sup>Department of Animal Nutrition, Animal Husbandry Faculty, Sam Ratulangi University, Jl. Kampus Unsrat Manado 95115, North Sulawesi, Indonesia

12

<sup>2</sup>Department of Animal Production, Animal Product Technology, Animal Husbandry Faculty, Sam Ratulangi University, Jl. Kampus Unsrat Manado 95115, North Sulawesi, Indonesia

\*Corresponding author email: jolalondok\_unsrat@yahoo.com

25

### Abstract

The aim of this experiment was to evaluate effect of level lauric acid (LA) combined with feed fiber (FF) in ration on serum lipid profile of broiler. A total of 360 unsex 1-d old broiler chicks were obtain from broiler breeding company. They were kept in brooding cage with temperature 23-33°C on a light/dark cycle until 7-d. After 7-d adjustment period, all bird randomly assigned to 12 treatments combination. Each group having three replicate cages with 30 birds was fed diet according experimental design. The experiment was conducted in a completely randomized design with a 3 x 4 factorial arrangement. The first factor was level of LA consisted of 3 levels i.e. 1.30%; 1.95%; and 2.60%, while the second was level of FF i.e. 5%, 6%, 7%, and 8% in the diet. Each treatment was given diet from 21 to 35 day of age during the experimental period. Effect of treatment on serum lipid consists of total cholesterol, triglyceride, HDL-cholesterol, and LDL-cholesterol was determined at the end of experiment (35-d of age). The conclusion of this study indicated that the used of 1.95% LA and 8% FF level on the diet would have an optimize serum lipid profile of broiler.

**Key words:** feed fiber, lauric acid, serum lipid profile.

### INTRODUCTION

Functional food is defined as a compound that contains physiologically active compounds (bioactive compounds) and is used for the prevention or cure of a disease or for achieving optimal body health. Increasing the economic status of the community, causing consumption of animal products to increase as well. The essential components in the community's diet are meat and processed meat products (Fernandez-Gines et al., 2005).

According to WHO in 2010, 25% of deaths in developing countries were due to cardiovascular disease. Cardiovascular disease and type-2 diabetes are considered as a result of increased consumption of animal products. Much research has been done to dismiss this assumption.

21

The results showed that total fat, saturated fatty acids and monounsaturated fatty acids were not associated with deaths from coronary heart disease, and consumption of saturated fatty acids was not related to the risk of coronary heart disease, stroke and cardiovascular disease.

Pure lauric acid (LA) can be used in feed with the same function as coconut oil as an energy source as well as a source of saturated fatty acids. Medium chain fatty acids (MCFA) in LA contain 95.5% lauric acid (Londok et al., 2018). The superiority of MCFA over long chain fatty acids (LCFA) is its metabolic processes in the body. MCFA has a smaller molecular weight so it does not require high energy and only requires a little enzyme to break down the fat into a form that is ready to be absorbed (Gheorghe et al., 2019; Papamandjaris et al., 1998; DebMandal and Mandal, 2015). Crude fiber is one of the important food substances in poultry rations, because it functions to stimulate the digestive tract peristalsis so that the digestion process of feed substances goes well. Poultry has limitations in digesting coarse fibers because the fermenter organ is located at the end of the absorptive organ. High crude fiber causes poultry to feel full, which can reduce consumption because crude fiber is voluminous. The level of energy in the feed will determine the amount of feed consumed. Broilers tend to increase their consumption if the metabolic energy content in feed is low.

Feed formulation by optimizing the use of fiber-rich feed ingredients can reduce meat fat content so that it becomes a safe/healthy meat product for consumers.

Strengthening the supply of food from animal experiences is a dilemma because on one hand meat consumption per capita is still low but on the other hand there is a tendency for certain consumers to limit consumption of livestock meat because of the negative effect of food on health. This issue is certainly a challenge for animal husbandry experts on how to develop businesses that can produce livestock commodities with carcasses that have high edible meat portions as a source of food that is safe and healthy for consumers. Food security which aims to increase the availability of food based on food security and independence, improving the quality of consumption and improving the quality and food safety.

45

#### MATERIALS AND METHODS

This study was conducted on 360 day-old unsexed Lohman broilers (MB 202-P) with an

average body weight of  $45.76 \pm 1.22$  grams, obtained from the breeding of PT Japfa Comfeed Indonesia Tbk chickens. Poultry Breeding Division Unit 13 Kauditan, Jl. Raya Manado-Bitung, Tumulung Village, North Minahasa Regency.

The chicken was placed in a brooder cage at a temperature of 23-33°C for 7 days. After the random adjustment period the chickens were placed in 12 treatment plots which were repeated 3 times. The study was designed using a completely randomized 3x4 factorial design. As factor A is the level of lauric acid (LA), which is A1 = 1.30%, A2 = 1.95%, and A3 = 2.60%. Factor B is the level of feed fiber (FF), namely B1 = 5%, B2 = 6%, B3 = 7%, and B4 = 8%. Determination of LA level based on the application of coconut oil in feed by 3% which is calculated lauric acid content.

There are 12 treatment combinations. LA used in this study was pure LA (99.5%) FA 1299 rays produced by Sinar Mas. Composition of the feed and its ingredients are presented in Table 1.

Table 1. Composition and ingredient of treatment feed (as fed)

Items	A1	A1	A1	A1	A2	A2	A2	A2	A3	A3	A3	A3
	B1	B2	B3	B4	B1	B2	B3	B4	B1	B2	B3	B4
<b>Ingredients (%)</b>												
Yellow Corn	31.7	22.7	20.7	8.70	26.05	27.05	20.05	8.05	29.4	23.4	19.4	11.4
SBM	22.0	20.0	23.0	7.0	16.0	19.0	23.0	7.0	21.0	20.0	23.0	7.0
Fish meal	5.0	5.0	6.0	9.0	6.0	6.0	6.0	9.0	5.0	5.0	6.0	8.0
Rice bran	14.0	20.0	28.0	28.0	11.0	21.0	28.0	29.0	14.0	20.0	28.0	29.0
MBM	5.0	4.5	4.0	7.0	5.0	4.0	7.0	6.0	5.0	4.5	4.0	7.0
BR-21F	20.0	25.5	16.0	38.0	19.0	16.0	37.0	32.0	22.0	23.5	16.0	34.0
LA	1.30	1.30	1.30	1.30	1.95	1.95	1.95	1.95	2.60	2.60	2.60	2.60
NaCl	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35	0.35
DL-methionine (99%)	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
Premix <sup>1</sup>	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60
Total	100	100	100	100	100	100	100	100	100	100	100	100
<b>Nutrient content</b>												
ME (Kcal/kg)	3125	3157	3159	3125	3287	3169	3148	3205	3207	3200	3171	3219
CP (%)	20.4	20.3	20.1	20.5	20.5	19.9	20.3	20.3	20.2	19.9	20.4	19.5
EE (%)	5.8	5.9	5.9	6.0	6.7	6.6	6.5	8.0	7.1	7.2	7.2	8.0
CF (%)	5.1	6.3	7.2	8.1	5.1	6.2	7.2	8.2	5.1	6.2	7.2	8.1

SBM, soy bean meal; MBM, meat bone meal; BR-21F, commercial feed; NaCl, natrium chloride; ME, metabolizable energy; CP, crude protein; CF, crude fiber. <sup>1</sup>Premix supplied the following per 100 g of diet: Iron, 40 mg; Copper, 26.16 mg; Zinc, 40 mg; Manganese, 44 mg; Selenium, 0.08 mg; Cobalt, 0.08 mg; Iodine, 0.52 mg; Vit A, 12500 IU; Vit D<sub>3</sub>, 35000 IU; Vit E, 25 IU; Vit K<sub>3</sub>, 4 mg; Vit B<sub>1</sub>, 4 mg; Vit B<sub>2</sub>, 8 mg; Vit B<sub>6</sub>, 20 mg; Vit B<sub>12</sub>, 50 mcg; Pantothenic acid, 15 mg; Niacin, 50 mg; Biotin, 125 mcg; Calcium D-pantothenate, 16.30 mg; Folic acid, 1 mg.

The treatment ration was made at a metabolic energy level between 3100-3200 kcal kg<sup>-1</sup> with crude protein level of 20%. The treatment feed and drinking water were applied ad libitum on days 21 to 35 days to the experimental chicken, so that each treatment combination was applied to 30 chicks. An interaction was shown by variables of total cholesterol, triglycerides, HDL-cholesterol, and LDL-cholesterol.

Measurement of serum fat done through laboratory analysis of total cholesterol, triglycerides, HDL-cholesterol, and LDL-cholesterol. Blood sampling for parameter determination was performed at the end of the study (day 35), in chickens that were fasted for ± 8 hours.

Overall data were analysed using the general linear model in Minitab (version 19). The difference in the mean value of the treatment was tested by Tukey simultaneous test. The difference was evaluated at the P level, 0.05.

## RESULTS AND DISCUSSIONS

The average of total cholesterol, triglycerides (TG), HDL-cholesterol, LDL-cholesterol and the ratio of HDL to LDL in chicken serum of this study were presented in Table 2. Total cholesterol, TG, HDL-cholesterol, LDL-cholesterol and HDL to serum LDL ratio were statistically significant influenced by the level of lauric acid, the level of feed fiber and the interaction of both. The different effect of blood lipids was thought to be caused by differences in fatty acids in the feed, because the transport of MCFA (lauric acid) is not in the form of lipoproteins. In the body, cholesterol is transported through the bloodstream by a type of fat called lipoprotein. There were several types of lipoproteins that were known, namely: very low density lipoprotein (VLDL) and LDL. These types of lipoproteins all contain a lot of TG and cholesterol. VLDL had more TG (60%), whereas LDL and HDL had more cholesterol (44%) (Piliang and Djojosoebagio, 2006). VLDL-cholesterol is usually released into the bloodstream continuously by the liver. This condition ensures a steady supply of TG to other cells of the body as a source of energy for metabolic processes. This increases

lipoproteins that are rich in cholesterol particles, in the form of LDL. MCFA does not reduce cholesterol through the transport of LDL cholesterol to the liver but by other mechanisms. Apolipoprotein B (Apo-B) is the main apolipoprotein from chylomicrons and LDL-cholesterol which is responsible for bringing cholesterol to the tissues. Apo-B in MCFA is lower than LCFA (Wang et al. 2015). The interaction between level of LA and the level of crude fiber ( $P < 0.05$ ) influences total cholesterol levels, TG, HDL-cholesterol, LDL-cholesterol and the ratio of HDL to LDL serum. The highest total serum cholesterol level was shown by the chicken consuming 1.95% LA and 6% FF ( $109.67 \pm 0.28$  mg.dL<sup>-1</sup>), while the lowest concentration was shown by the fed chicken supplemented with 2.6% LA with 7% (103.59 ± 0.28 mg.dL<sup>-1</sup>). Saturated fats decrease LDL receptor (LDLr) activity and increase LDL production (Merchant et al., 2008). LDLr is the key to regulating cholesterol homeostasis feedback (Musa et al., 2007). The absorption of LDL cholesterol from peripheral tissues is mediated by LDLr. LDLr is negatively correlated with serum LDL and abdominal fat percentage (Murwani et al., 2011). When LDLr is high, the serum LDL and abdominal fat are low. Research on adding palm oil and coconut oil to broiler feed was carried out by Piliang et al. (1995). It was reported that a decrease in total cholesterol, HDL and serum LDL. Dong et al. (2003), using coconut oil as a source of MCFA in pregnant mice. The use of coconut oil decreases TG, increasing serum HDL-cholesterol at the age of 3 weeks. Fatimah and Rindengan (2011) found the same thing using pure coconut oil in rat. Londok et al. (2014) supplemented virgin coconut oil as a source of lauric acid up to 3% in high fiber-based feed in broiler feed, it was able to increase serum HDL, reduce serum LDL and reduce meat cholesterol. Furthermore Londok (2018) supplemented 3% coconut oil in natural antioxidant *Areca vestiaria* Giseke at a dose of 1250 mg.kg<sup>-1</sup> showing a marked increase in HDL and a decrease in LDL. Wang et al. (2015) uses coconut oil and soybean oil as a source of MCFA in broiler feed. Linearly, total cholesterol and LDL decreased, but TG increased while HDL did not differ significantly.

Table 2. Average of lipid profile in broiler serum consumed different level of LA and FF (mg dL<sup>-1</sup>)

Parameters	LA Level (%)	FF Level (%)				Average
		5	6	7	8	
Total cholesterol (mg dL <sup>-1</sup> )	1.30	103.91±1.07 <sup>AB</sup>	109.67±0.28 <sup>A</sup>	103.59±0.28 <sup>AB</sup>	101.01±1.16 <sup>BC</sup>	104.54±3.65
	1.95	94.75±1.93 <sup>CDE</sup>	104.96±3.89 <sup>AB</sup>	107.46±3.04 <sup>AB</sup>	103.26±2.27 <sup>AB</sup>	102.61±5.51
	2.60	95.58±2.76 <sup>CD</sup>	91.44±2.49 <sup>DE</sup>	88.12±1.93 <sup>E</sup>	89.08±4.29 <sup>DE</sup>	91.05±3.32
	Average	98.08±5.06	102.02±9.46	99.72±10.23	97.78±7.62	
Triglycerides (mg dL <sup>-1</sup> )	1.30	47.64±2.37 <sup>ABC</sup>	45.87±2.03 <sup>ABCD</sup>	47.04±3.19 <sup>ABC</sup>	46.87±2.37 <sup>ABC</sup>	46.85±0.78
	1.95	41.59±2.25 <sup>BCD</sup>	36.59±4.28 <sup>D</sup>	43.68±2.14 <sup>ABCD</sup>	40.77±2.31 <sup>BCD</sup>	40.66±2.98
	2.60	48.85±4.23 <sup>AB</sup>	40.39±5.00 <sup>BCD</sup>	38.85±5.00 <sup>CD</sup>	52.23±1.47 <sup>A</sup>	45.08±6.49
	Average	46.03±3.89	40.95±4.67	43.19±4.12	46.62±5.74	
HDL-cholesterol (mg dL <sup>-1</sup> )	1.30	52.28±1.11 <sup>DE</sup>	55.74±1.61 <sup>DE</sup>	59.28±2.43 <sup>BCD</sup>	49.31±3.58 <sup>E</sup>	54.15±4.31
	1.95	66.63±2.82 <sup>AB</sup>	67.79±3.15 <sup>AB</sup>	68.22±5.44 <sup>AB</sup>	56.60±3.96 <sup>CDE</sup>	64.81±5.51
	2.60	70.45±1.48 <sup>A</sup>	65.87±2.60 <sup>ABC</sup>	60.19±3.83 <sup>BCD</sup>	56.92±3.15 <sup>CDE</sup>	63.36±6.00
	Average	63.12±9.58	63.13±6.48	62.56±4.92	54.28±4.30	
LDL-cholesterol (mg dL <sup>-1</sup> )	1.30	42.10±1.71 <sup>A</sup>	44.76±1.74 <sup>A</sup>	34.91±3.34 <sup>BC</sup>	42.33±2.93 <sup>A</sup>	41.02±4.25
	1.95	19.80±1.34 <sup>D</sup>	29.85±0.12 <sup>C</sup>	30.50±1.97 <sup>C</sup>	38.50±2.15 <sup>AB</sup>	29.66±7.66
	2.60	15.36±0.43 <sup>D</sup>	20.36±3.05 <sup>D</sup>	20.16±4.76 <sup>D</sup>	21.72±0.85 <sup>D</sup>	19.40±2.78
	Average	25.76±14.33	31.65±12.30	28.52±7.57	34.18±10.96	
HDL: LDL ratio	1.30	1.24±0.08 <sup>D</sup>	1.25±0.08 <sup>D</sup>	1.72±0.23 <sup>CD</sup>	1.17±0.17 <sup>D</sup>	1.35±0.25
	1.95	3.39±0.37 <sup>B</sup>	2.27±0.11 <sup>BCD</sup>	2.26±0.32 <sup>BCD</sup>	1.85±0.66 <sup>CD</sup>	2.44±0.66
	2.60	4.59±0.03 <sup>A</sup>	3.05±0.37 <sup>B</sup>	3.21±0.95 <sup>B</sup>	2.62±0.04 <sup>BC</sup>	3.37±0.85
	Average	3.07±1.69	2.19±0.90	2.40±0.75	1.88±0.72	

The average ratio of HDL to LDL in serum of experimental chickens that consume feed supplemented with different lauric sources and antioxidant concentrations provides a very significant interaction ( $P < 0.01$ ) in influencing serum HDL: LDL ratio. The highest average serum HDL: LDL ratio was shown by chicken fed a combination of feed containing 2.6% LA with 5% dietary fiber ( $4.59 \pm 0.03$  mg.dL<sup>-1</sup>), while the lowest HDL: LDL ratio was shown by a combination of 1.3% LA with 8 % FF ( $1.17 \pm 0.17$  mg.dL<sup>-1</sup>). Coronary atherosclerosis is associated with a high plasma cholesterol: HDL: LDL ratio (Mayes et al. 1995). Each 5% increase in energy consumption from saturated fat is proportional to the energy consumption from carbohydrate and causes a 17% increase in the risk of heart disease (Hu et al. 1997). However Kolondam et al. (2008), concluded that administration of virgin coconut oil (VCO) with therapeutic doses (0.95 mL) and two doses in wistar rats (*Rattus norvegicus*) had no effect on blood lipid levels after being consumed for 30 days. Increasing the level of VCO did not cause changes in blood lipid levels during the study. Other results also showed that changes in blood lipid levels after administration of virgin coconut oil did not result in changes in the risk of heart disease in test animals, and

claims about the dangers of saturated fat consumption in VCO against increased blood lipids were not proven.

## CONCLUSIONS

The use of lauric acid 1.95% with 8% feed fiber level provides optimal broiler serum lipid profile.

## ACKNOWLEDGEMENTS

This research work was carried out with the support of Ministry of Research Technology and Higher Education Republic of Indonesia, also was financed from Directorate Research and Community Dedication, Directorate General of Research Development Strengthening, by contract letter No. 210/UN12.13/LT/2019

## REFERENCES

- Al-Taleb, S.S. (2003). Effect of an early feed restriction on productive performance and carcass quality. *J. Biol. Sci.*, 3, 634-641.
- Benevent. (1981). *Quelques Aspects de la Croissance Chez les Animaux Supérieur delevage*. ENSA-Montpellier.

- Bortoluzzi, Fernandes, C., Contini, J.I.M., Gurski, 15  
Esser, T.J., Prokoski, A.F.G. (2013). Quantitative feed  
restriction 15 from 35 to 42 days of age for broiler  
23 chicken. *Rev Bras Saude Prod Anim.*, 14(4), 778-784.  
Broiler Performance and Nutrition Supplement. 2015.  
(http://www.cobb-vantress.com/docs)
- Butzen, F.M., Riberio, A.M.L., Vieira, M.M., Kessler,  
A.M., Dadalt, J.C., Della, M.P. (2013). Early feed  
restriction in broiler. I-Performance, body fraction  
weights, and meat quality. *J. Appl. Poult. Res.*, 22,  
6 251-259.
- DebMandal, M., Mandal, S. 2011. Coconut (*Cocos  
nucifera* L.: Arecaceae): In health promotion and  
disease prevention. *Asian Pac. J. Trop. Med.*, 4(3),  
3 241-247.
- Fernandez-Gines, J.M., Fernandez-Lopez, J., Sayas-  
Barbera, E., Perez-Alvares, J.A. 2005. Meat products  
as functional foods: a review. *J. Food Sci.*, 70(2), 37-  
43.
- Gheorghe, A., Habeanu, M., Tabuc, C., Marin, M.  
(2019). Effects of dietary pea seeds (*Pisum sativum*  
L. cv. Tudor) on performance, carcass traits, plasma  
biochemistry and intestinal microflora in broiler  
1 chicks. *AgroLife Scientific Journal*, 8(1), 99-106.
- Jahanpour, H., Seidavi, A., Qotbi, A.A.A., Van Den  
Hoven, R., Rocha e Silva, S., Laudadio, V., Tuvarelli,  
V. (2015). Effects of the level and duration of feeding  
restriction on carcass component of broiler. *Arch.*  
10 *Anim. Breed.*, 58, 99-105.
- Leeson, S., Summers, J.D. (2008). *Commercial Poultry  
Nutrition*. 3<sup>rd</sup> ed. Nottingham, UK: Nottingham  
University Pr.
- Lohmann Meat. (2007). *Broiler Stock Performance  
Objectives*. Alabama, US: Aviagen.
- Londok, J.J.M.R., Manalu, W., Wiryawan, KG., Sumiati.  
(2017). Growth performance, carcass characteristics  
and fatty acids profile of broilers supplemented with  
lauric acid and natural antioxidant from *Areca  
vestiaria* Giseke. *Pakistan Journal of Nutrition*,  
16(9), 719-730.
- Londok, J.J.M.R., Tulung, B., Kowel, Y.H.S., Rompis.  
J.E.G. (2012). Effect of feed restriction on feed  
efficiency, carcass quality and digestive organs  
characteristics of broiler. Proceeding the 2<sup>nd</sup>  
International Seminar "Feed Safety for Healthy  
Food". July 6-7 2011. AINI and Faculty of Animal  
24 Husbandry, Univesitas Padjajaran. Jatinangor.
- Murtidjo, B.A. (2003). *Pemotongan dan Penanganan*  
4 *Daging Ayam*. Yogyakarta, ID: Kanisius.
- Papamandjaris, A.A., McDogall, D.E., Jones, P.J.H.  
(1998). Medium chain fatty acid metabolism and  
energy expenditure: obesity treatment implications.  
Minireview. *Life Sciences*, 62(4), 1203-1215.
- Prudhon, M 17 Benevent, M., Vezinhet, J. P., Dulong  
A.A.T. (1976). Croissance relative du squalete chez  
l'agneau. Influence du sexe et de la race. *Ann. Biol.  
Anim. Bioch. Biophys.*, 18(1), 5-19
- Robelin, J. (1979). Evaluation de la composition  
corporelle des jeunes bovine males entiers de race  
limosine entre 9 et 19 mois. *Ann. Zootech.*, 26(4),  
11 333-346.
- Santoso, U. (2001). Effect early feed restriction on  
growth, fat accumulation and meat composition in  
unsexed 20 broiler chicken. *J. Anim. Sci.*, 4, 1585-1591.
- [SNI] Standar Nasional Indonesia. 3924-2009. *Mutu  
Karkas dan Daging Ayam*. Jakarta, ID: Badan  
Standarisasi Nasional 13 al-BSN.
- Soeparno (1992). *Ilmu dan Teknologi Daging*. Edisi I.  
Yogyakarta, ID: Gadjah Mada University Press.
- Soeparno (2011). *Ilmu Nutrisi dan Gizi Daging*.  
Yogyakarta, ID: Gadjah Mada University Press.
- Tulung, B. (1999). *Isu, Kontroversi dan upaya  
Penurunan Kolesterol Produk Hewan* 33 Pidato  
Penguahan Guru Besar bidang Ilmu Nutrisi &  
Makanan Ternak Fakultas Peternakan Universitas  
Sam Ratulangi Manado. Manado, ID: Unsrat Press.

# SERUM LIPID PROFILE OF BROILER CONSUMING RATION CONTAINING LAURIC ACID AND FEED FIBER

## ORIGINALITY REPORT

20%

SIMILARITY INDEX

17%

INTERNET SOURCES

15%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.arch-anim-breed.net](http://www.arch-anim-breed.net)

Internet Source

1%

2

[animalsciencejournal.usamv.ro](http://animalsciencejournal.usamv.ro)

Internet Source

1%

3

[www.nature.com](http://www.nature.com)

Internet Source

1%

4

[www.journal.eu-jr.eu](http://www.journal.eu-jr.eu)

Internet Source

1%

5

[es.scribd.com](http://es.scribd.com)

Internet Source

1%

6

[www.australiancurriculum.edu.au](http://www.australiancurriculum.edu.au)

Internet Source

1%

7

[docobook.com](http://docobook.com)

Internet Source

1%

8

[www.koreascience.or.kr](http://www.koreascience.or.kr)

Internet Source

1%

9

[www.pjbs.org](http://www.pjbs.org)

Internet Source

1%

10	"Abstracts 2012", British Poultry Abstracts, 2012 Publication	1 %
11	J.D. Liu, J. Fowler. "Effect of inert fillers with changing energy-protein ratio on growth performance and energy digestibility in broilers", Journal of Applied Poultry Research, 2017 Publication	1 %
12	kipdf.com Internet Source	1 %
13	medpub.litbang.pertanian.go.id Internet Source	1 %
14	penerbit.brin.go.id Internet Source	1 %
15	www.bvs-vet.org.br Internet Source	1 %
16	epdf.tips Internet Source	1 %
17	www.gse-journal.org Internet Source	1 %
18	ojs.unud.ac.id Internet Source	<1 %
19	J. ROBELIN, R. DAENICKE. "Variations of net requirements for cattle growth with liveweight, liveweight gain, breed and sex", Annales de Zootechnie, 1980 Publication	<1 %



---

20	<a href="https://id.scribd.com">id.scribd.com</a> Internet Source	<1 %
21	<a href="http://www.rachelarthur.com.au">www.rachelarthur.com.au</a> Internet Source	<1 %
22	Jola J.M.R. Londok, John E.G. Rompis, Claudya Mangelep. "KUALITAS KARKAS AYAM PEDAGING YANG DIBERI RANSUM MENGANDUNG LIMBAH SAWI", ZOOTEK, 2016 Publication	<1 %
23	<a href="https://api.crossref.org">api.crossref.org</a> Internet Source	<1 %
24	<a href="https://repository.unair.ac.id">repository.unair.ac.id</a> Internet Source	<1 %
25	<a href="https://academic.oup.com">academic.oup.com</a> Internet Source	<1 %
26	<a href="http://garuda.kemdikbud.go.id">garuda.kemdikbud.go.id</a> Internet Source	<1 %
27	<a href="https://2020.agricultureforlife.usamv.ro">2020.agricultureforlife.usamv.ro</a> Internet Source	<1 %
28	A J Bennett, M A Billett, A M Salter, E H Mangiapane, J S Bruce, K L Anderton, C B Marenah, N Lawson, D A White. "Modulation of hepatic apolipoprotein B, 3-hydroxy-3-methylglutaryl-CoA reductase and low-density lipoprotein receptor mRNA and plasma lipoprotein concentrations by	<1 %

---

defined dietary fats. Comparison of trimyristin, tripalmitin, tristearin and triolein", Biochemical Journal, 1995

Publication

---

29

Joseph Oluwatope Adebayo, Olumuyiwa Sunday Adewumi, Simbiat Titilayo Baruwa, Elizabeth Abidemi Balogun et al. "Effects of polyphenols of Cocos nucifera husk fibre on selected indices of cardiovascular diseases in mice", TANG [HUMANITAS MEDICINE], 2016

Publication

---

<1 %

30

MEHMET TUGRUL CABYOGLU, NEYHAN ERGENE, UNER TAN. "THE TREATMENT OF OBESITY BY ACUPUNCTURE", International Journal of Neuroscience, 2009

Publication

---

<1 %

31

Xiaoming Sun, Jinglin Shen, Chang Liu, Sheng Li et al. "L-arginine and N-carbamoylglutamic acid supplementation enhance young rabbit growth and immunity by regulating intestinal microbial community", Asian-Australasian Journal of Animal Sciences, 2020

Publication

---

<1 %

32

[bmcpregnancychildbirth.biomedcentral.com](https://bmcpregnancychildbirth.biomedcentral.com)

Internet Source

---

<1 %

33

[docplayer.info](https://docplayer.info)

Internet Source

---

<1 %

34	<a href="http://horizon.documentation.ird.fr">horizon.documentation.ird.fr</a> Internet Source	<1 %
35	<a href="http://jmums.mazums.ac.ir">jmums.mazums.ac.ir</a> Internet Source	<1 %
36	<a href="http://repository.upnjatim.ac.id">repository.upnjatim.ac.id</a> Internet Source	<1 %
37	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet Source	<1 %
38	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	<1 %
39	Bo Hou, Wencheng Wang, Hui Gao, Shanglang Cai, Chunbo Wang. " Effects of aqueous extract of L. roots on serum lipid metabolism ", Journal of International Medical Research, 2017 Publication	<1 %
40	C. S. Williamson. "Red meat in the diet", Nutrition Bulletin, 12/2005 Publication	<1 %
41	<a href="http://archive.org">archive.org</a> Internet Source	<1 %
42	<a href="http://repository.ub.ac.id">repository.ub.ac.id</a> Internet Source	<1 %
43	<a href="http://repository.unhas.ac.id">repository.unhas.ac.id</a> Internet Source	<1 %
44	<a href="http://www.european-poultry-science.com">www.european-poultry-science.com</a> Internet Source	<1 %

45

[www.science.gov](http://www.science.gov)

Internet Source

&lt;1 %

46

M. Rouhi, S. Sohrabvandi, A. M. Mortazavian. "Probiotic Fermented Sausage: Viability of Probiotic Microorganisms and Sensory Characteristics", *Critical Reviews in Food Science and Nutrition*, 2011

Publication

&lt;1 %

47

Mi-Ja Choi, Kyung Ja Chang. "Chapter 40 The Effects of Dietary Taurine Supplementation on Plasma and Liver Lipid in Ovariectomized Rats", Springer Science and Business Media LLC, 2009

Publication

&lt;1 %

48

Adélia P. Miranda, Jorge de Lucas Junior, Maria C. Thomaz, Gener T. Pereira, Ellen H. Fukayama. "Anaerobic biodigestion of pigs feces in the initial, growing and finishing stages fed with diets formulated with corn or sorghum", *Engenharia Agrícola*, 2012

Publication

&lt;1 %

49

M. A Soljak. "Risk factor thresholds", *BMJ*, 11/9/2002

Publication

&lt;1 %

50

Terry Lichtor, Harry R. Davis, Lydia Johns, Draga Vesselinovitch, Robert W. Wissler, Sean Mullan. "The sympathetic nervous system and atherosclerosis", *Journal of Neurosurgery*, 1987

&lt;1 %

## Publication

---

Exclude quotes Off

Exclude bibliography Off

Exclude matches Off