

Nata_as_a_source_of_dietary.p df *by*

Submission date: 03-Jan-2022 09:13PM (UTC+0700)

Submission ID: 1737113417

File name: Nata_as_a_source_of_dietary.pdf (588.18K)

Word count: 4300

Character count: 23263

Nata as a source of dietary fiber with numerous health benefits

Trina Ekawati Tallei^{1*}, Siti Marfuah¹, Abdul Hawil Abas¹, Andi Amelia Dwi Putri Abram¹, Nelsyani Pasappa¹, Putri Sri Anggini¹, Agus Supandi Soegoto², Fatima Wali³, Talha Bin Emran^{4*}

¹Department of Biology, Faculty of Mathematics and Natural Sciences, Sam Ratulangi University, Manado 95115, North Sulawesi, Indonesia

²Department of Management, Faculty of Economics and Business, Sam Ratulangi University, Manado 95115, North Sulawesi, Indonesia

³Pharmacy Study Program, Faculty of Economics and Business, Sam Ratulangi University, Manado 95115, North Sulawesi, Indonesia

⁴Department of Pharmacy, BGC Trust University Bangladesh, Chittagong 4381, Bangladesh

*Corresponding authors

Trina Ekawati Tallei, PhD
e-mail: trina_tallei@unsrat.ac.id
and Talha Bin Emran, PhD
e-mail: talhabmb@bgctub.ac.bd

Academic editor

Akhi Moni, PhD
ABEx Bio-Research Center,
Bangladesh

Article info

Received: 17 August 2021
Accepted: 04 November 2021
Published: 11 November 2021

Keywords

Dietary fiber; functional food; insoluble fiber; nata; health benefit; organic waste.

ABSTRACT

In recent years, the general public's perception of modern diets and human health has shifted dramatically. Changes in lifestyle make people prefer fast food that is poor in nutrients, especially dietary fiber. Lack of dietary fiber is one of the contributions to the increasing prevalence of non-communicable diseases, such as hypercholesterolemia and cancer, among the public. For this reason, efforts to introduce fiber-rich functional foods need to be encouraged. Nata, which is a high-fiber food made from organic plant sources, can be used as an alternative source of fiber for the community. It is hoped that this article will provide some insight into alternative sources of dietary fiber that can be produced by members of the community on their own.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Sustainable development goals (SDGs) number 2 is to end hunger, achieve food security, improve nutrition, and promote sustainable agriculture [1]. Health care is a necessity so that the state can empower the potential of human resources. Currently, the modern food industry is being focused, among others, on various types of beverages and functional foods, which are aimed at increasing the nutritional value, and especially to obtain the health benefits of these products [2]. Diets high in fiber, such as cereals, nuts, fruits, and vegetables, are beneficial to health because their consumption has been linked to a lower incidence of several diseases [3].

One source of dietary fiber is nata. Nata in Spanish is literally defined as cream, such as nata de coco, which means the cream of coconut water. Nata was originally known as nata de coco, but it now comes in a variety depending on the source of the raw material. Nata de coco is composed of millions of fine cellulose threads, which eventually appear solid white to transparent and are referred to as nata. The dietary fiber content in nata de coco is beneficial to the body because it is required and important for digestion.

Nata is beneficial not only as a source of dietary fiber, but also as a wound healer. Nata de coco, for example, can be further processed into a new material that is very strong, heat resistant, as well as flexible and can even transmit light. In a study, one of the products that may be produced is the monitor screen [4]. Nata de coco is a conductive polymer with a conductivity of 553 S/cm that exhibits excellent mechanical stability [5]. In this review, we will focus exclusively on the health benefits of nata as a source of

dietary fiber. Additionally, various natas derived from a variety of raw materials will be discussed.

HISTORY OF NATA

Nata was first introduced in the Philippines in 1973 as nata de coco in an attempt to utilize coconut water waste and preserve it as a jelly-like substance. Nata de coco (alternatively marketed as "coconut gel") is a chewy, translucent, jelly-like food made from coconut water that gels due to *Acetobacter xylinum* producing bacterial cellulose (BC). Nata de coco is primarily made from coconut water, and as a result, it has a low nutritional profile. However, because it is made of cellulose, it is high in dietary fiber. Cellulose (Figure 1), being the most important component of plant cell walls, is a polysaccharide with the formula $(C_6H_{10}O_5)_m$ which is composed of thousands of (1→4) linked β -D-glucose units [6]. During the 1990s, nata de coco became very popular in Japan. Nowadays, it can be found in a variety of different flavors and shapes, and while it can be eaten raw, it is most often used to make refreshing fruit salads, yogurts, ice cream, and beverages as an ingredient.




Figure 1. The structure of cellulose ($m = 2,000 - 26,000$). Each monomer unit is β -D-glucose, and each beta acetal link connects the carbon atom 1 of one glucose to the carbon atom 4 of the next glucose.

VARIOUS TYPES OF NATA

Nata's main final product is cellulose. As is the case with other microbial products, carbon and nitrogen are the primary ingredient as a source of nutrients for microbes. Additionally, vitamins and minerals, collectively referred to as trace elements, are added in trace amounts to promote the growth of microbial cells and the formation of the desired product. *A. xylinum* is the microorganism responsible for converting these nutrients into BC.

Various fruit-based carbon sources can be used to make nata. Various natas have been developed successfully, including nata de piña. This nata can be made from pineapple juice or pineapple peel waste juice. Typically, pineapple peel waste is only used as animal feed. To add economic value to pineapple peel waste, it can be used as a raw material in the production of nata. Nata de coco piña is another nata product that we developed, which is made from coconut water waste and pineapple peel waste (Figure 2). Figure 3 illustrates the nata de coco piña processing. Essentially, the process of making nata is similar, as it utilizes *A. xylinum*, a cellulose-producing bacteria. The only difference is in the organic raw material sources and the amount of sugar and ammonium sulfate used. Table 1 shows the proximate analysis of nata de coco piña. When it comes to the proximate analysis, the sources of organic matter and the amount of sugar used as a starter are important factors to consider. These materials will also determine the thickness and organoleptic features of the resulting nata.

Nata de soya is made from liquid waste generated during the tofu manufacturing process. This waste is the primary by-product of the tofu manufacturing process and has the potential to pollute the environment. Nata de banana is made from banana peel waste. Cocoa bean pulp can be processed into nata de cacao. Nata de cassava is made from tapioca liquid waste. Nata de pinata is made from palm sap. Nata de cane is made from waste sugarcane stalks left over from bud chips which are often wasted, which actually contains a fairly high juice content. Nata de seaweed is made from any kind of seaweed, for example *Eucheuma cottonii*. With these examples, different types of nata can still be created from other organic wastes.

Table 1. The proximate analysis of nata de coco piña.

Parameter	Concentration (%)
Water content	96.71
Protein	0.33
Fat	1.53
Crude fiber	0.93
Carbohydrate	0.59
Calories	17.36
Phosphor (P)	0,01
Sucrose	0.44
Iron (Fe)	1.26 (ppm)
Calcium	153.95 (ppm)



Figure 2. Nata de coco piña.




Figure 3. The flowchart of nata de coco piña processing.

DIETARY FIBERS

Dietary fiber is a part of plants that can be eaten but cannot be digested by human digestion, but can only be digested or processed into simpler products by bacteria found in the large intestine. Dietary fiber is composed of cellulose, non-cellulosic polysaccharides such as hemicellulose, pectic substances, gums, and mucilages, and lignin, a non-carbohydrate component [3]. Fiber is classified into the following two types based on its solubility in water: insoluble fiber and water soluble fiber. Soluble and insoluble fiber content varies among plant foods, with different properties [7].

Soluble fiber is a type of fiber that dissolves easily in water and generates viscous gels. They are not digested in the small intestine but are readily fermented in the large intestine by the microbiota [8]. Soluble fibers can be found in plant cell walls; among others are arabinoxylans (AX), β -glucans, some hemicelluloses, pectins, gums, and inulin [9]. Soluble fiber is usually found in fruits.

Insoluble fibers do not dissolve in water in the human gastrointestinal tract, so they do not form gels, and are also rarely fermented by the intestinal microbiota [8]. Insoluble fiber is commonly found in tough plant cell walls. Among these are cellulose, hemicelluloses, lignin, and resistant starch [9]. Insoluble fiber is found in whole grains,

fruit skins, cucumbers, tomatoes, rice husks (most commonly brown rice), legumes, and beans [10].

HEALTH BENEFITS OF DIETARY FIBERS

Dietary fiber is essential for human health. However, current intake levels of fibre and fiber-rich foods are still far below recommended levels in the majority of countries around the world [11]. Increasing the intake of dietary fiber has been recommended by a number of health organizations, with specific recommendations of 25-30 grams per day. In recent years, there has been an increase in interest in functional foods that can adjust body function and prevent civilizational lifestyle diseases [12]. Dietary fiber is essential for maintaining a healthy digestive system. The deficiency of fiber is especially relevant in light of the astonishingly high incidence rates of colon cancer in urban societies [13,14].

Dietary administration of fiber changes the niche environment in the gut by providing nutrients and substrates for microbiota growth, which allows microbiota that are able to utilize these nutrients to multiply and spread throughout the body [15]. The low administration of dietary fiber not only suppresses the diversity of microbiota in the digestive tract, but also shifts the metabolic pattern of the microbiota towards the use of substrates that are not in accordance with the benefits of these microbiota (16). This event will in turn disrupt the host's metabolism, which will lead to metabolic disorders that have an impact on host health disorders.

Increased insoluble cereal fiber intake resulted in a considerable improvement in whole-body glucose elimination, leading to an 8% increase in insulin sensitivity [17]. Additionally, insoluble fiber may result in a decrease in appetite and food intake [18]. Insoluble fiber speeds up the flow of food through the digestive tract, resulting in smoother bowel movements [19]. In large prospective cohort studies, ingestion of insoluble cereal dietary fibre ³ has been linked to a lower risk of type 2 diabetes mellitus development (20). Increased whole-grain intake is associated with a decreased risk of various diseases, including coronary heart disease, cardiovascular disease, stroke, respiratory disease, and infectious disease, according to a meta-analysis of prospective cohorts [21-23].

HEALTH BENEFITS OF NATA DE COCO

Nata is a source of insoluble dietary fiber due to its cellulose content. Nata de coco contains about 98% water, 0.2% fat, 0.012% calcium, 0.002% phosphorus, 0.0017% vitamin B3, 51 mg/g sodium, potassium 280 mg/100 g, and 2.46 mg/100g vitamin C. This product has a high fiber content, including cellulose (2.5%), hemicellulose, lignin, and soluble fiber [24,25]. Components of chemical compounds contained in nata de coco include: hexadecanoic acid (7.58%), benzeneacetic acid (7.73%), 22-hydroxyhopane (3.96%), tetradecanoic acid (3.84%), 9-octadecenoic acid (3.65%), *o*-cresol (3.50%), 9-octadecenamide, (Z) (3.00%), phenol, 4-(2-aminoethyl) (2.73%), dodecanoic acid (2.21%), pentadecanoic acid (1.79%), 1-heptadecanecarboxylic acid (1.64%), indole (1.79%), hydrocinnamic acid (1.60%), heptadecanoic acid (1.54%), dan cyclohexanecarboxylic acid (1.47%) [26].

A functional food refers to a food product that contains nutrients but also has the potential to provide additional health benefits, either an improvement in one's health and well-being or a decrease in the risk of contracting disease [27]. With regards to this

definition, nata can be classified as a functional food. The following describes the health benefits of nata as a functional food.

Control of biological functions

Hexadecanoic acid (palmitic acid) is a 16-carbon long-chain saturated fatty acid. It is the most abundant saturated fatty acid in the human body, accounting for between 20% and 30% of total fatty acids. At the cellular and tissue level, palmitic acid performs a variety of fundamental biological functions [28]. In a study, BC inhibits α -amylase significantly in an in vitro chyme model. Additionally, it is capable of adsorbing significant amounts of glucose via its binding to glucose molecules. Thus, BC has the ability to regulate blood sugar levels and can be used as a dietary supplement for patients with hyperglycaemia [29]. BC also plays an important role in increasing cellular adhesion, hence contributes to tissue re-epithelialisation [30].

Antifungal and antimicrobial properties

Nata is rich in a variety of fatty acids (FAs). FAs have been shown to be extremely promising for development as next-generation antibacterial agents for the treatment of a broad spectrum of bacterial infections [31]. Benzeneacetic acid (phenylacetic acid) is well-known for its antifungal properties [32-34]. This compound possesses a broad antimicrobial spectrum and inhibited the growth of several soil-borne phytopathogenic fungi completely [33]. Tetradecanoic acid and hexadecanoic acid have antimicrobial activities against multidrug-resistant bacteria [35].

Wound dressing

Bacterial cellulose, a naturally occurring gelly-like substance produced by *A. xylinum*, is widely used in wound dressings due to its high water-holding capacity and mechanical strength [36]. Wound dressings can help to speed up the healing process of the wound by increasing the permeability and protection of the new tissue [37]. Bacterial cellulose has also been used as an alternative carrier of C_{60} (an effective photosensitizer for photodynamic therapy) in the form of a multifunctional wound dressing in the treatment of skin cancer [38].

Contribution to the control of plasma cholesterol levels

Tetradecanoic acid (myristic acid) is a long-chain saturated fatty acid composed of 14 C atoms. This acid is first extracted from the nutmeg plant. It is related to low plasma HDL cholesterol levels in the Mediterranean population [39]. In hypercholesterolemic women, nata de coco consumption can lower total cholesterol levels in the blood [40].

Contribution to cancer cure and prevention

According to the findings of a large number of epidemiological and experimental studies, dietary fiber may play an important role in colon cancer prevention [41]. In several studies it is said that BC is a suitable material in treating various cancers due to its ability to absorb and deliver drugs [42]. Furthermore, according to the findings of a study, BC-*Garcinia mangostana* extract has the potential to be used as an anti-breast cancer biofilm candidate in the future [43]. As a result, there is still room for

advancement of nata in terms of its potential contribution to cancer cure and prevention.

CONCLUSIONS AND FUTURE PROSPECTS

Nata, a jelly-like substance, is a product of fermentation by the bacterium *A. xylinum*. Nata is primarily composed of cellulose, an insoluble fiber. Insoluble fiber has many health benefits, including facilitating bowel movements, reducing the risk of heart disease, cancer, and diabetes, and lowering cholesterol levels in the blood. This nutritious food can be utilized as a functional food, and it is simple to make in the community by repurposing organic waste resources. Nata can be fortified with the addition of vitamins and a variety of other key nutrients, enhancing its potential as a functional food by providing additional health advantages (Figure 4).




Figure 4. Schematic diagram of summary of the study. Juice made of organic wastes is subjected for fermentation with *A. xylinum*. These provide various kinds of natas. These natas are the potential sources of insoluble fiber and other essential micronutrients with many health benefits.

ACKNOWLEDGEMENT

This research was funded by Sam Ratulangi University, Manado, Indonesia, under the scheme of Development Research for University Excellence 2021 with Letter of assignment Number 1089/UN12/LL/2021 and Contract Number 609/UN12/LL/2021.

AUTHOR CONTRIBUTIONS

TET, F and ASS were involved in conception and design of the experiments. AHA, AADPA, NP, PSA and SM contributed to perform the experiments. TET, F, ASS and

¹³ TBE contributed to drafting the article. TET and TBE contributed to revising it critically for important intellectual content. TET and TBE made the final approval of the version to be published.

CONFLICTS OF INTEREST

There is no conflict of interest among the authors.

¹⁵ REFERENCES

- [1] Gil JDB, Reidsma P, Giller K, Todman L, Whitmore A, van Ittersum M. Sustainable development goal 2: improved targets and indicators for agriculture and food security. *Ambio*. 2019; 48(7):685–98.
- [2] Ali A, Rahut DB. Healthy Foods as Proxy for Functional Foods: Consumers' Awareness, Perception, and Demand for Natural Functional Foods in Pakistan. *Int J Food Sci*. 2019; 2019:6390650. <https://doi.org/10.1155/2019/6390650>
- [3] Dhingra D, Michael M, Rajput H, Patil RT. Dietary fibre in foods: a review. *J Food Sci Technol*. 2012; 49(3):255–261. <https://pubmed.ncbi.nlm.nih.gov/23729846>
- [4] Uetani K, Koga H, Nogi M. Estimation of the Intrinsic Birefringence of Cellulose Using Bacterial Cellulose Nanofiber Films. *ACS Macro Lett*. 2019; 8(3):250–254. <https://doi.org/10.1021/acsmacrolett.9b00024>
- [5] Mulyasuryani A, Mustaghfirah AM. Development of Potentiometric Phenol Sensors by Nata de Coco Membrane on Screen-Printed Carbon Electrode. *J Anal Methods Chem*. 2019; 2019:4608135. <https://doi.org/10.1155/2019/4608135>
- [6] Holtzaple MT. Cellulose. In: Caballero BBT-E of FS and N (Second E, editor. Oxford: Academic Press; 2003. p. 998–1007. Available from: <https://www.sciencedirect.com/science/article/pii/B012227055X001851>
- [7] Mudgil D. Chapter 3 - The Interaction Between Insoluble and Soluble Fiber. In: Samaan RABT-DF for the P of CD, editor. Academic Press; 2017. p. 35–59.
- [8] Latimer JM, Haub MD. Effects of dietary fiber and its components on metabolic health. *Nutrients*. 2010; 2(2):1266–89.
- [9] Ciudad-Mulero M, Fernández-Ruiz V, Matallana-González MC, Morales P. Dietary fiber sources and human benefits: The case study of cereal and pseudocereals. *Adv Food Nutr Res*. 2019; 90:83–134.
- [10] Suharoschi R, Pop OL, Vlaic RA, Muresan CI, Muresan CC, Cozma A, et al. Chapter 3 - Dietary Fiber and Metabolism. In: Galanakis Recovery, and Applications CMBT-DFP, editor. Academic Press; 2019. p. 59–77. Available from: <https://www.sciencedirect.com/science/article/pii/B9780128164952000034>
- [11] Komarek AR. Dietary fibre basics: Health, nutrition, analysis, and applications. *Food Qual Saf*. 2017; 1(1):47–59. <https://doi.org/10.1093/fqsafe/fyx007>
- [12] Yang Y, Ma S, Wang X, Zheng X. Modification and Application of Dietary Fiber in Foods. *J Chem*. 2017; 7:9340427. <https://doi.org/10.1155/2017/9340427>
- [13] O'Keefe SJD. The Need to Reassess Dietary Fiber Requirements in Healthy and Critically Ill Patients. *Gastroenterol Clin North Am*. 2018; 47(1):219–229.
- [14] Masrul M, Nindrea RD. Dietary Fibre Protective against Colorectal Cancer Patients in Asia: A Meta-Analysis. *Open access Maced J Med Sci*. 2019;7(10):1723–1727. <https://pubmed.ncbi.nlm.nih.gov/31210830>
- [15] Deehan EC, Duar RM, Armet AM, Perez-Muñoz ME, Jin M, Walter J. Modulation of the Gastrointestinal Microbiome with Nondigestible Fermentable Carbohydrates To Improve Human Health. *Microbiol Spectr*. 2017; 5(5).
- [16] Makki K, Deehan EC, Walter J, Bäckhed F. The Impact of Dietary Fiber on Gut Microbiota in Host Health and Disease. *Cell Host Microbe*. 2018; 23(6):705–715.
- [17] Weickert MO, Mohlig M, Koenig C, Holst JJ, Namsolleck P, Ristow M, et al. Impact of cereal fibre on glucose-regulating factors. *Diabetologia*. 2005; 48(11):2343–2353.
- [18] Samra RA, Anderson GH. Insoluble cereal fiber reduces appetite and short-term food intake and glycemic response to food consumed 75 min later by healthy men. *Am J Clin Nutr*. 2007; 86(4):972–979.
- [19] Axelrod CH, Saps M. The Role of Fiber in the Treatment of Functional Gastrointestinal Disorders in Children. *Nutrients*. 2018; 10(11).
- [20] Hřjova E, Bertkova I, Štofilyova J. Dietary fibre as prebiotics in nutrition. *Cent Eur J Public Health*. 2019; 4(3):251–255.
- [21] Benisi-Kohansal S, Saneei P, Salehi-Marzjarian M, Larijani B, Esmailzadeh A. Whole-Grain Intake and Mortality from All Causes, Cardiovascular Disease, and Cancer: A Systematic Review and Dose-Response Meta-Analysis of Prospective Cohort Studies. *Adv Nutr*. 2016; 7(6):1052–1065.
- [22] Aune D, Keum N, Giovannucci E, Fadnes LT, Boffetta P, Greenwood DC, et al. Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies. *BMJ*. 2016; 353:i2716.
- [23] Zong G, Gao A, Hu FB, Sun Q. Whole Grain Intake and Mortality From All Causes, Cardiovascular Disease, and Cancer: A Meta-Analysis of Prospective Cohort Studies. *Circulation*. 2016; 133(24):2370–2380.
- [24] Ma T, Ji K, Wang W, Wang J, Li Z, Ran H, et al. Cellulose synthesized by Enterobacter sp. FY-07 under aerobic and anaerobic conditions. *Bioresour Technol*. 2012; 126:18–23.
- [25] Santosa B, Wignyanto W, Hidayat N, Sucipto S. The quality of nata de coco from sawarna and mapanget

- coconut varieties to the time of storing coconut water. *J Food Sci.* 2019; 4:957-963.
- [26] Anam C. Mengungkap Senyawa pada Nata De Coco sebagai Pangan Fungsional. *J Ilmu Pangan dan Has Pertan.* 2019; 3(1):42.
- [27] González-Díaz C, Vilaplana-Aparicio MJ, Iglesias-García M. How Is Functional Food Advertising Understood? Approximation in University Students. *Nutrients.* 2020; 12(11).
- [28] Carta G, Murru E, Lisai S, Sirigu A, Piras A, Collu M, et al. Dietary triacylglycerols with palmitic acid in the sn-2 position modulate levels of N-acyl ethanolamides in rat tissues. *PLoS One.* 2015; 10(3):e0120424–e0120424.
- [29] Zhang L-L, Zhang W, Peng F-B, Chen H, Shu G-W. Effects of bacterial cellulose on glucose metabolism in an in vitro dyme model and its rheological evaluation. *Int J Food Sci & Technol.* <https://doi.org/10.1111/ijfs.15244>
- [30] Swingler S, Gupta A, Gibson H, Kowalczyk M, Heaselgrave W, Radecka I. Recent Advances and Applications of Bacterial Cellulose in Biomedicine. *Polymers.* 2021; 13(3):412. <https://doi.org/10.3390/polym13030412>
- [31] Sillas-Vargas G, Ocasio-Malavé C, Medina S, Morales-Guzmán C, Del Valle RG, Carballeira NM, et al. Antibacterial fatty acids: An update of possible mechanisms of action and implications in the development of the next-generation of antibacterial agents. *Prog Lipid Res.* 2021; 82:101093. <https://doi.org/10.1016/j.plipres.2021.101093>
- [32] Hwang BK, Lim SW, Kim BS, Lee JY, Moon SS. Isolation and in vivo and in vitro antifungal activity of phenylacetic acid and sodium phenylacetate from *Streptomyces humidus*. *Appl Environ Microbiol.* 2001; 67(10):3739-3745.
- [33] Kim Y, Cho J-Y, Kuk J-H, Moon J-H, Cho J-I, Kim Y-C, et al. Identification and Antimicrobial Activity of Phenylacetic Acid Produced by *Bacillus licheniformis* Isolated from Fermented Soybean, Chungkook-Jang. *Microbiol.* 2004; 48(4):312-317. <https://doi.org/10.1007/s00284-003-4193-3>
- [34] Sajid I, Shaaban KA, Hasnain S. Identification, isolation and optimization of antifungal metabolites from the *Streptomyces malachitofuscus* ct9. *Braz J Microbiol.* 2011;42(2):592–604. <https://doi.org/10.1590/S1517-83822011000200024>
- [35] Kima J-E, Seo J-H, Bae M-S, Bae C-S, Yoo J-C, Bang M-A, et al. Antimicrobial Constituents from *Allium hookeri* Root. *Nat Prod Commun.* 2016; 11(2):237-238.
- [36] Yang M, Ward J, Choy K-L. Nature-Inspired Bacterial Cellulose/Methylglyoxal (BC/MGO) Nanocomposite for Broad-Spectrum Antimicrobial Wound Dressing. *Macromol Biosci.* 2020; 20(8):2000070. <https://doi.org/10.1002/mabi.202000070>
- [37] Lin CH, Chen JC, Huang CM, Juang TY. High-performance thermosetting films based on an amino-functionalized poly(ether sulfone). *J Appl Polym Sci.* 2014; 131(21). <https://doi.org/10.1002/app.40980>
- [38] Chu M, Gao H, Liu S, Wang L, Jia Y, Gao M, et al. Functionalization of composite bacterial cellulose with C60 nanoparticles for wound dressing and cancer therapy. *RSC Adv.* 2018; 8(33): 1819-1820. <https://dx.doi.org/10.1039/C8RA03965H>
- [39] Noto D, Fayer F, Cefalù AB, Altieri I, Palesano O, Spina R, et al. Myristic acid is associated with low plasma HDL cholesterol levels in a Mediterranean population and increases HDL catabolism by enhancing HDL particles trapping to cell surface proteoglycans in a liver hepatoma cell model. *Atherosclerosis.* 2016; 246:50-56.
- [40] Purwani NPR, Mulyati T. Pengaruh Pemberian Nata De Coco Terhadap Kadar Kolesterol Total Pada Wanita dengan hiperkolesterolemia. *J Nutr Coll.* 2012; 1(1):249-257. <https://doi.org/10.14710/jnc.v1i1.735>
- [41] Zeng H, Lazarova DL, Bordonaro M. Mechanisms linking dietary fiber, gut microbiota and colon cancer prevention. *World J Gastrointest Oncol.* 2014; 6(2):41–51. <https://doi.org/10.4251/wjgo.v6.i2.41>
- [42] Islam SU, Ul-Islam M, Ahsan H, Ahmed MB, Shehzad A, Fatima A, et al. Potential applications of bacterial cellulose and its composites for cancer treatment. *Int J Biol Macromol.* 2021; 168:301-309.
- [43] Agrippina WRG, Widiyanti P, Yusuf H. Synthesis and Characterization of Bacterial Cellulose - *Garcinia mangostana* Extract as Anti Breast Cancer Biofilm Candidate. *J Biomimetics, Biomater Biomed Eng.* 2017; 30:76–85. <https://doi.org/10.4028/www.scientific.net/BBBBE.30.76>

Nata_as_a_source_of_dietary.pdf

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

%
INTERNET SOURCES

%
PUBLICATIONS

21 %
STUDENT PAPERS

PRIMARY SOURCES

1 Submitted to Joliet Junior College **2** %
Student Paper

2 Submitted to University of Arizona **1** %
Student Paper

3 Submitted to University of Queensland **1** %
Student Paper

4 Submitted to University of West Florida **1** %
Student Paper

5 Submitted to University of Sheffield **1** %
Student Paper

6 Submitted to University of South Florida **1** %
Student Paper

7 Submitted to Universitas Sam Ratulangi **1** %
Student Paper

8 Submitted to Loma Linda University **1** %
Student Paper

9 Submitted to Swinburne University of Technology **1** %
Student Paper

10

Submitted to CVC Nigeria Consortium

Student Paper

1 %

11

Submitted to Universidade de Sao Paulo

Student Paper

1 %

12

Submitted to University of Exeter

Student Paper

1 %

13

Submitted to Ewha Womans University

Student Paper

1 %

14

Submitted to Izmir Katip Æelebi Æniversitesi

Student Paper

1 %

15

Submitted to Johns Hopkins University

Student Paper

1 %

16

Submitted to Higher Education Commission
Pakistan

Student Paper

1 %

17

Submitted to University of Nebraska at
Omaha

Student Paper

1 %

18

Submitted to University of Newcastle upon
Tyne

Student Paper

1 %

19

Submitted to Chester College of Higher
Education

Student Paper

1 %

20 Submitted to Texas A&M University, College Station 1 %
Student Paper

21 Submitted to University of California, Merced 1 %
Student Paper

22 Submitted to University of Edinburgh 1 %
Student Paper

23 Submitted to New York Chiropractic College 1 %
Student Paper

24 Submitted to Queen's University of Belfast 1 %
Student Paper

Exclude quotes Off

Exclude matches < 1%

Exclude bibliography Off