

# Identification of pigment profiles and antioxidant activity of *Rhizophora mucronata* mangrove leaves origin Lembeh, North Sulawesi, Indonesia

*by Darus Sa'adah Johanis Paransa 7*

---

**Submission date:** 30-May-2023 08:22AM (UTC+0700)

**Submission ID:** 2104891212

**File name:** BID\_B\_-\_JURNAL\_INTERNASIONAL\_-\_2021\_-\_antonius\_dkk.pdf (764.98K)

**Word count:** 10372

**Character count:** 57375

## Identification of pigment profiles and antioxidant activity of *Rhizophora mucronata* mangrove leaves origin Lembeh, North Sulawesi, Indonesia

ANTONIUS PETRUS RUMENGAN<sup>1,\*</sup>, ELVI SISKA MANDIANGAN<sup>2</sup>, WENDY ALEXANDER TANOD<sup>3</sup>, DARUS SAJAH JOHANIS PARANSA<sup>1</sup>, CAROLUS PAULUS PARUNTU<sup>1</sup>, DESY MARIA HELENA MANTIRI<sup>1</sup>

<sup>1</sup>Department of Marine Science, Faculty of Fisheries and Marine Science, Universitas Sam Ratulangi, Jl. Kampus UNSRAT Bahu, Manado 95115, North Sulawesi, Indonesia. Tel./fax.: +62-852-4027-3030, \*email: antonius\_rumengan@unsrat.ac.id

<sup>2</sup>Research and Development Agency of North Sulawesi Province, Jl. 17 Agustus, Bumi Beringin, Wenang, Manado 95113, North Sulawesi, Indonesia

<sup>3</sup>Department of Fisheries and Marine, Politeknik Negeri Nusa Utara, Jl. Kesehatan, Mahena, Tahuna, Kepulauan Sangihe 95812, North Sulawesi, Indonesia

Manuscript received: 23 May 2021. Revision accepted: 21 June 2021.

**Abstract.** Rumengan AP, Mandiangan ES, Tanod WA, Paransa DSJ, Paruntu CP, Mantiri DMH. 2021. Identification of pigment profiles and antioxidant activity of *Rhizophora mucronata* mangrove leaves origin Lembeh, North Sulawesi, Indonesia. *Biodiversitas* 22: 2805-2816. Mangroves plants contain unique pigments which can serve both nutraceutical and pharmaceutical purpose. Therefore, this preliminary study aims to identify the pigment profiles of *Rhizophora mucronata* mangrove leaves (HPLC method) and to computationally evaluate the antioxidant mechanism of the pigments (PASS and STITCH analysis). Furthermore, it evaluated the antioxidant capacity of *R. mucronata* leaves extracts (DPPH method), and the pigment profiles detected using HPLC were chlorophyll a (68.61%), chlorophyll b (27.69%), lutein (29.94%), beta-carotene (14.05%), pheophytin a (8.72), violaxanthin (5.19%), and neoxanthin (3.65%). Beta-carotene, lutein, neoxanthin, and violaxanthin were predicted to have potential as antioxidants properties using PASS analysis. While neoxanthin and violaxanthin were predicted as free radical scavengers, beta-carotene was Nrf-2 stimulant. The STITCH analysis showed that the pigments contained in the leaves interacted synergistically by activating as antioxidant enzymes and inhibiting the expression of oxidative stress proteins. The ethanol extract of *R. mucronata* leaves may be a potent antioxidant with an IC50  $20.99 \pm 0.33 \mu\text{g/mL}$ . Therefore, the pigment contained in *R. mucronata* leaves is a potential source of antioxidants.

**Keywords:** Beta-carotene, chlorophyll, lutein, neoxanthin, violaxanthin

### INTRODUCTION

Mangrove plants live in low tide areas, and they play a critical role in the coastal ecosystem by supporting fisheries, tourism, and a genetic reservoir (Liu 2012). It functions as a protector of the coastline, preventing seawater abrasion, the habitat of various aquatic biota, and acts best carbon storage ecosystem (Kepel et al. 2017; Rumengan et al. 2018; Saptiani et al. 2018; Tidore et al. 2018). Furthermore, coastal communities have also used mangroves for house construction, charcoal making materials (Dahdouf, Jebas et al. 2000), functional foodstuffs (Jariyah et al. 2014; Kardiman et al. 2017; Analuddin et al. 2019), textile dyes (Pringgienies et al. 2017), and traditional medicine (Revathi et al. 2014; Rout and Basak 2014; Alhaddad et al. 2019; Andriani et al. 2020). These plants have a self-defense mechanism because they live in areas with fluctuations in salinity and temperature (Dewan et al. 2018). The mechanism designed consists of enzymatic and non-enzymatic antioxidants defense (Sarker et al. 2018a; Sarker and Oba 2018a; 2020a). Pigments in mangroves significantly affect photosynthetic reactions, stress avoidance, and defense mechanisms (Croft and Chen 2017).

Previous studies reported that leaves are the natural sources of pigments such as chlorophyll a, chlorophyll b, betacyanins, betaxanthins, betalains (Sarker et al. 2014;

2015a; 2015b; 2017). These are potential sources of carotenoids pigments such as beta-carotene, alpha-carotene, xanthophylls including lutein, neoxanthin, zeaxanthin, and violaxanthin (Sarker and Oba 2021). In addition, leaves consist of other pigments such as phenolics and different groups of flavonoids, including flavonols, flavones, flavanones (Sarker and Oba 2019a; 2020b; 2020c), and they all exhibit high antioxidant capacity (Sarker et al. 2018b).

Previous studies also showed that mangroves contain unique pigments which may serve nutraceutical and pharmaceutical functions (Banerjee et al. 2017). One of the mangroves has the potential to use its pigment content such as *Rhizophora mucronata* (Pringgienies et al. 2018). Prabhu and Bhute (2015) reported that the brown pigment from the stem of *R. mucronata* was used as a textile dye. Furthermore, it was previously reported that leaves contain chlorophyll and carotenoid pigments (Flores-de-Santiago et al. 2016; Ridlo et al. 2017). However, no study has reported the complete profile of the pigments in the mangrove leaves of *R. mucronata*. The pigments can be used as a source of functional and natural food coloring (Mapari et al. 2005).

The mangrove *Rhizophora mucronata* showed medicinal potential for coastal communities. Meanwhile, the fruit, leaves, bark, and flowers of *R. mucronata* were used to treat various diseases, such as cognitive function

(Suganthy and Pandima Devi 2016), diabetes (Bandaranayake 1998; Aljaghthmi et al. 2017), diarrhea (Puspitasari et al. 2012), hepatitis (Ravikumar and Gnanadesigan 2012), and inflammation (Rohini and Das 2009). The antioxidant capacity of the leaves obtained from India and Asia was widely reported (Imdadul et al. 2011). This is a preliminary study to identify the pigment profiles of *Rhizophora mucronata* mangrove leaves using the high-performance liquid chromatography method and to computationally evaluate the antioxidant mechanism of the pigments. Furthermore, it evaluated the antioxidant capacity of *R. mucronata* leaves extracts using the DPPH method.

## MATERIALS AND METHODS


### Sampling and extraction

Mangrove leaves sample (1000 g) was obtained from Lembah Island, North Sulawesi, Indonesia. *Rhizophora mucronata* leaves were documented and put in a cool temperature container. They were collected in summer with hot sunny conditions (in July 2019) and were taken to the laboratory for extraction. The leaves were finely mashed to a powder, and the extraction was performed three times with ethanol pure grade (Merck) solvent. In addition, the *R. mucronata* leaf powder (100 g) was macerated using ethanol (1:3 w/v) for 48 hours, with occasional shaking. The resultant maceration product was filtered through filter

paper to separate the filtrate from the residue. Furthermore, the filtrate was evaporated (rotary vacuum evaporator Buchi R-300) at 40°C during 4-6 hours to obtain a crude extract (Dewanto et al. 2018). According to Lichtenthaler (1987), the chlorophyll and carotenoid groups have a phytol chain attached to the porphyrin ring system. The possession of the phytol side chain, which is esterified to the ring carboxyl group, gives chlorophyll and carotenoids their lipid character. Therefore, ethanol is one of the polar solvents used to extract fat-soluble pigments from living plant tissues containing water. Figure 1 showed the location of mangrove leaves sampling.

### Identification of pigment profiles

The identification of mangrove leaves pigments was conducted on a High-Performance Liquid Chromatography (HPLC-20AD-Shimadzu) with SP-M20A photodiode array detector (PDA). The pigment analysis was based on the method by Hegazi et al. (1998), and the Shimadzu UV-1800 spectrophotometer was used in determining the wavelength. Meanwhile, the detection of mangrove leaves pigments was conducted by HPLC at 450 nm (carotenoids detection) and 660 nm (chlorophyll detection) respectively. The analytical column was a LiChroCART 250×4 mm I.D. packed with Chromospher 100, RP-18e (5µm spherical particles), and the precolumn was an ODS-hypersil (C18) with a diameter of 5µm, 20×4 mm.



**Figure 1.** Location of *Rhizophora mucronata* mangrove leaves sampling in Lembah Island, North Sulawesi, Indonesia (1° 27' 8.78" N and 125° 14' 41.90" E)

Dry pigment crude extract was dissolved in 5 mL acetone (Paransa et al. 2014) and filtered using a filter membrane (0.2 µm), then 20 µL was injected into HPLC. The pigment elution was conducted at a 1 mL/min flow rate at 20 °C using a gradient elution system from a mixture of methanol, acetone, and ammonium acetate (1 M) solutions. Acetone was used in the pigment extraction because of its amphipathic nature of having a polar and a nonpolar end. Also, it has a significant partial negative and positive charge on the oxygen and carbonyl atom with two nonpolar alpha carbons. However, it is less polar than water and ethanol and can dissolve nonpolar substances. The nature of its small polarity allows it to dissolve polar substances and has fewer properties than water and ethanol. Therefore, acetone is an appropriate solvent, which allows more excellent resolution for detecting pigments using chromatography. It breaks chlorophyll lipid bonds to plant thylakoid structures and suspends pigments in solution (Henriques et al. 2007).

#### Computational analysis

The pigment detected with HPLC predicted biological activity-related antioxidants using the PASS (prediction of activity spectra for substances) analysis (<http://www.pharmaexpert.ru/passonline/index.ph>). PASS is a tool used for predicting the biological activity of compounds (Riyadi et al. 2020). The predicted activity requires a structural formula in the form of canonical SMILE obtained from the National Center for Biotechnology Information (<https://pubchem.ncbi.nlm.nih.gov/>). The pigments were analyzed for their interactions with the STITCH database (search tool for interactions of chemicals) (<http://stitch.embl.de/>).

#### Antioxidant assay

Measurement of the antioxidant capacity of mangrove leaves was carried out using the DPPH method (Oke and Hamburger 2002), and the assays were conducted using 96 well plate microplates. The ethanol extract of *Rhizophora mucronata* leaves was prepared in series concentrations of 10, 20, 40, 80 µg/mL in methanol solution (Merck). Then, 160 µL of extract from each concentration series was fed into the microplate well. Furthermore, 40 µL of DPPH (Merck) 0.76 mM solution was added to each well that contained a sample. A comparison control used vitamin C with a concentration series of 4, 6, 8, 10 µg/mL. As a control sample, each series of dilutions (160 µL) was added to the microplate well before adding 40 µL of methanol. A negative control (without extract) was made by adding 160 µL methanol with 40 µL DPPH, and 200 µL methanol was blank. The microplate was incubated in a dark room at 25–28 °C for 30 minutes, and after that, the absorbance of each well was measured with the Multiskan GO Microplate Spectrophotometer (Thermo Scientific) at a wavelength of 517 nm. The IC<sub>50</sub> determination was measured from the inhibition percentage data in units of µg/mL, and a probit analysis was used to determine the IC<sub>50</sub> value. The

following equation was used to determine the percentage of DPPH inhibition:

$$\% \text{ Inhibition} = \frac{(C - D) - (A - B)}{(C - D)} \times 100\%$$

Where :

A: Sample absorbance

B: Absorbance control sample

C: Absorbance of negative control

D: Absorbance blank

## RESULTS AND DISCUSSION

The dry extract of *Rhizophora mucronata* mangrove leaves was detected for pigment profiles by HPLC at a wavelength of 450 and 660 nm, with a retention time of 20 to 40 minutes. Table 1 and Figure 2 showed the pigment profiles, and the chromatogram for the leaves extract.

Table 1 showed that the *R. mucronata* leaves extract detected seven pigments dominated by chlorophyll a (68.61%) and chlorophyll b (27.69%). Chlorophyll is a unique green pigment in almost every green part of plants, such as leaves (İnanç 2011). Chlorophyll is a pigment with a chlorine ring, where magnesium is surrounded by four nitrogen atoms (Taiz et al. 2006). Meanwhile, in chlorophyll b, the –CHO group replaces –CH<sub>3</sub> on the C7 atom (Pareek et al. 2017). There is also chlorophyll a and b epimer at 11.69% and 5.19%, and Limantara and Heriyanto (2012) stated that they always accompany the presence of chlorophyll. Chlorophyll a and b are common and dominant pigments in the leaves of green plants, such as mangroves (Liu 2012; Dou et al. 2018). Ridlo et al. (2017) reported *R. mucronata* leaves to contain chlorophyll and carotenoid pigments. Furthermore, Flores-de-Santiago et al. (2016) also reported that *Rhizophora mangle* leaves contain chlorophyll a and b. Their levels depend on the season and physiological conditions of the mangroves. Bohn and Walczyk (2004) detected chlorophyll a and the epimer at the wavelength spectrum of 429, 664 nm, while chlorophyll b and the epimer were discovered at 456, 648 nm. Kusmita et al. (2015) detected chlorophyll-b at the wavelength spectrum of 456, 596, 645 nm.

Lutein is a member of the xanthophyll and carotenoid family (Al-ali et al. 2020), and it has a yellowish-orange color, called macular pigment, which is not in abundance (Landrum and Bone 2001; Aruldass et al. 2018). Ngginak et al. (2017) reported detecting lutein on the wavelength spectrum of 417, 443, and 472 nm, while Kurniawan et al. (2020) reported detecting it at 447, 451, 472 nm. Also, Sarker and Oba (2021) reported that leaves contain good macular pigment lutein. Previously, no study has detected lutein pigment in mangrove leaves. The detection was conducted using random sampling of old mangrove leaves that were yellowish in color.

**Table 1.** Identification of pigment profiles from *Rhizophora mucronata* leaves with HPLC

Wavelength detection (nm)	Retention time (min)	Area (%)	Wavelength spectrum (nm)	Pigment profiles	References (Jeffrey et al. 1997; Zapata et al. 2000)	
					Retention time (min)	Wavelength spectrum (nm)
450	20.06	3.65	415, 437, 465	Neoxanthin	–	415.1, 438.5, 467.1
450	21.72	5.19	417, 442, 470	Violaxanthin	21.32	416, 440, 470
450	29.29	24.94	421, 447, 475	Lutein	27.65	422, 446, 475
450	33.91	20.65	462, 598, 647	Chlorophyll b	31.62	462, 599, 648
660		7.04				
450	34.20	5.19	461, 599, 648	Chlorophyll b epimer	31.87	462, 599, 650
450	35.53	11.30	430, 617, 663	Chlorophyll a	33.15	431, 617, 662
660	35.53	57.31				
450	35.89	2.08	430, 617, 663	Chlorophyll a epimer	33.48	430, 615, 664
660	35.89	9.61	430, 616, 663			
660	37.63	8.72	409, 505, 535, 608, 664	Pheophytin a	–	409.5, 505.3, 534.7, 608.9, 665.5
450	38.39	14.05	453, 479	Beta-carotene	35.95	426, 452, 477

Beta-carotene is one of the red-yellow, orange or red-orange carotenoids in natural plants that carry out photosynthesis (Kusbandari and Susanti 2017). It may be fat-soluble, insoluble in water, easily damaged, unstable at high temperatures, and precursor of vitamin A (Strobel et al. 2007). Furthermore, carotenoids including beta-carotene act as an antioxidant. Previous studies showed that they have strong DPPH and ABTS antioxidant activity in different amaranth species such as drought-tolerant amaranth (Sarker and Oba 2020d), *A. gangeticus* (Sarker et al. 2020a), *A. hypochondriacus* (Sarker and Oba 2020e), stem amaranth (Sarker et al. 2020b), *A. blitum* (Sarker and Oba 2020f), green amaranth (Sarker et al. 2020c), weedy amaranth (Sarker and Oba 2019b), and red amaranth (Sarker and Oba 2019c). Furthermore, carotenoids including beta-carotene protect the photosynthetic tissue through direct quenching of triplet chlorophyll. This prevents the generation of singlet oxygen from oxidative damage in abiotic stress like salinity and drought (Sarker and Oba 2020a). It also detoxifies various forms of reactive oxygen species (ROS) (Sarker and Oba 2018b; 2018c) through increasing beta-carotene concentration (Sarker and Oba 2018d; 2018e; 2019d). Beta-carotene is a pigment synthesized by plants (Bogacz-Radomska and Harasym 2018). Radu et al. (2012) reported detecting this pigment on a wavelength spectrum of 445, 472, 498 nm. Meanwhile, it was detected by Mangunsong et al. (2019) at a wavelength spectrum of 460 nm. Beta-carotene was also detected on the wavelength spectrum of 450.20 and 477.60 nm (Kusbandari and Susanti 2017). The literature showed that this pigment was detected in the leaves and roots of mangroves *Avicennia officinalis*, *Excoecaria agallocha*, *Kandelia candel*, and *Rhizophora mucronata* (Ravindran et al. 2012). In addition, it was also detected in the mangroves of *Bruguiera gymnorrhiza*, *Sonneratia alba*, and *Xylocarpus granatum* (Analuddin et al. 2019).

Pheophytin a pigment is a chlorophyll a without  $Mg^{2+}$  ion, and it is dominant in fresh green leaves, which is degraded due to heating and storage processes (Hsu et al. 2013). It is produced naturally by plant leaves and acts as an intermediary for the first electron carrier in the transfer

pathway for photosystem II plants (Eijkelhoff and Dekker 1997). Bohn and Walczyk (2004) detected pheophytin-a on the wavelength spectrum of 405, 661 nm while Kusmita et al. (2015) were on wavelength spectrum of 408, 505, 535, 609, 765 nm.

Violaxanthin is a natural xanthophyll pigment with orange color found in various plants (Giossi et al. 2020). It is biosynthesized from zeaxanthin by an epoxidation reaction and has a 5,6-epoxy double group found in orange fruits, green vegetables, and microalgae (Melendez-Martinez et al. 2008). Furthermore, it plays a role in photoprotection mechanisms such as the ability of plants to adapt to contrasting light environments (Bowen-O'Connor et al. 2013). Wang et al. (2018) detected violaxanthin pigments at the wavelength spectrum of 417.6, 440.9, and 470.1 nm while Ruban et al. (2001) detection was at 470 nm. These pigments were also reported in mangroves with high salinity (Falqueto et al. 2008).

Neoxanthin pigments are carotenoids of xanthophylls groups. It acts as an intermediary for the biosynthesis of the hormone abscisic acid in plants (Perreau et al. 2020). Furthermore, it serves as a protection against photooxidative stress (Dall'Osto et al. 2007). Neoxanthin is the primary xanthophyll pigment found in green plants (Giossi et al. 2020). Chandrika et al. (2005) detected neoxanthin in the wavelength spectrum of 413, 436, 465 nm, and was detected by Gupta et al. (2015) at 415, 437, 465 nm. In mangroves, the neoxanthin pigment was found in mangroves *Avicennia alba* (Sasamoto et al. 2020).

The pigment detected in *R. mucronata* leaves extract predicted its probability to be active (Pa) value as an antioxidant. Bioactivity prediction was conducted using PASS analysis (Filimonov et al. 2014). The Pa value describes the potential activity of a compound. When the  $Pa > 0.7$ , it is estimated to have a high bioactivity potential, both for computational and laboratory assays. Meanwhile, when the value is  $0.3 \leq Pa \leq 0.7$ , then the compound has the computational ability as an antioxidant, but it has not been proven in the laboratory. Also, when the  $Pa < 0.3$ , it is predicted that the compound has a low bioactivity potential (Aisiah et al. 2020; Riyadi et al. 2021). However, the

bioactivity of chlorophyll a and b may not be predicted by PASS analysis because they have a metal element in their structure. The pigments beta-carotene, lutein, neoxanthin, pheophytin-a, and violaxanthin have their potential bioactivity as antioxidants, free radical scavengers, and NF-E2-related factor (Nrf-2) stimulant. Figure 3 showed the probability to be active as the antioxidant of the pigment in the *R. mucronata* leaves extract.

Figure 3 showed that beta-carotene, lutein, neoxanthin, and violaxanthin are predicted to have potential as general antioxidants. Neoxanthin and violaxanthin were also predicted to be free radical scavengers. Meanwhile, beta-carotene was predicted as an Nrf-2 stimulant, which regulates antioxidant protein as protection from oxidative damage (Ma 2013; Cui et al. 2016; Riyadi et al. 2019). Chlorophyll cannot be predicted by PASS analysis as an antioxidant, even though it has antioxidant properties as predicted by previous studies (İnanç 2011; Keleş et al. 2016; Sarker et al. 2018c; 2018d). Durga et al. (2015)

reported chlorophyll a and b from medicinal plants with high potential antioxidants. Pérez-gálvez et al. (2020) stated that chlorophyll b showed higher antioxidant activity than chlorophyll a.

Kurniawan et al. (2020) stated that lutein can act as an antioxidant and maintaining organs such as the eyes, brain, and skin. The content in marigold plant (*Tagetes* spp.) was reported to have a DPPH radical inhibition ability of 89.90% (Ingkasupart et al. 2015). Furthermore, it was reported to work as an antioxidant in the photo-stressed retina (Kamoshita et al. 2016).

Beta-carotene is also reported to be an antioxidant and an anti-carcinogen (Paolini et al. 2003). Berti et al. (2014) stated that it is effective as a radioprotective agent and acts as an antioxidant. Mueller and Boehm (2011) reported that beta-carotene and its derivatives showed antioxidant properties measured by the  $\alpha$ TEAC, chemiluminescence (CL), and ferric reducing activity (FRAP) methods.




Figure 2. HPLC Chromatogram of *Rhizophora mucronata* leaves ethanol extract




Figure 3. Probability to be active of pigments contain in *Rhizophora mucronata* leaves as an antioxidant with PASS analysis

Furthermore, Figure 3 showed the prediction of beta-carotene bioactivity as an Nrf-2 stimulant. The literature showed beta-carotene can activate the Nrf2-ARE (antioxidant response element) pathway to provide a neuroprotective effect from traumatic brain injury (Chen et al. 2019). Ben-dor et al. (2005) reported that beta-carotene stimulates Nrf-2 in the leukemia promyelocytic core body and regulates phase II enzyme expression (associated with cancer-preventing gene activation).

Dall'Osto et al. (2007) stated that neoxanthin acts as an antioxidant in the photosystem II supercomplex in plant thylakoid to protect membrane lipids photooxidation. In addition, Giossi et al. (2020) reported that neoxanthin was directly involved in photoprotection as an antioxidant to increase the activity of ROS scavenger under extreme light conditions. Sarker and Oba (2020g) reported that the neoxanthin contained in *Amaranthus tricolor* showed DPPH radical scavenging activity. Neoxanthin reduces oxidative-induced DNA base damage by less than 50%. In lower concentrations than lutein, it is a better inhibitor of oxidative-induced DNA damage (Şahin et al. 2020).

Dall'Osto et al. (2007) also reported the antioxidant properties of violaxanthin as photoprotection, even though it is lower than neoxanthin. The literature also reports the antioxidant properties of violaxanthin and its derivatives isolated from mangoes, with strong lipid peroxidation inhibition capabilities (Araki et al. 2016). Furthermore, violaxanthin isolated from microalgae *Eustigmatos cf. polyphem* was also reported to have radical scavenger capabilities with 2,2-diphenyl-1-picrylhydrazyl (DPPH), and 2,2-azobis-3-ethylbenzthiazoline-6-sulphonic acid (ABTS) radical assays (Wang et al. 2018). However, Figure 2 showed that pheophytin has a low Pa value as an antioxidant, and it prevents oxidative DNA damage and lipid peroxidation. It works by reducing reactive oxygen species, such as  $OH^{\cdot}$ , and by chelation of metal ions, such as Fe (II) (Hsu et al. 2013). Kusmita et al. (2015) also reported that pheophytin contained in green tea has an antioxidant capacity (DPPH method) with  $IC_{50} = 573 \pm 0.23$  mg/L.

The mechanism of antioxidant action of the pigments in *R. mucronata* leaves extract should be properly understood. Furthermore, the pigments in Table 1 were evaluated by STITCH. This is a tool for integrating information about the interactions of metabolic pathways, crystal structure, binding experiments, and relationships between chemicals (Kuhn et al. 2008). Figure 4 showed the interactions between the pigments.

Figure 4 showed the interaction between the working mechanism of the pigments in the *R. mucronata* leaves extract. Chlorophyll a, chlorophyll b, neoxanthin, and violaxanthin support the performance of lutein and beta-carotene. Furthermore, Beta-carotene pigments activate and catalyze the transcriptional regulation of the BCMO1 protein (beta, beta-Carotene 15,15'-monooxygenase-1), which is a key enzyme in vitamin A metabolism (Lietz et al. 2010). BCMO1 was reported to be expressed in intestinal tissue and plays a role in lipid metabolism (Lietz et al. 2012). It affects the RBP2 protein (Retinol-binding protein 2), which plays a role in the absorption and

metabolism of intracellular vitamin A (Sener et al. 2020). Figure 4 showed that lutein affects the expression of BCMO2 (Beta, beta-carotene 9', 10'-oxygenase) to catalyze beta-carotene. In addition, Lietz et al. (2012) also reported that BCMO2 contributes to the formation of vitamin A to improve antioxidant performance and immune function (Zhang et al. 2019).

Figure 4 also showed that lutein affects the PPARA gene (Peroxisome proliferator-activated receptor alpha), which strengthens antioxidant and anti-inflammatory defense systems (Moran et al. 2014) by increasing regulation of the enzyme expression and inhibiting the activity of NF- $\kappa$ B (nuclear factor kappa b) (Gao and Li 2012). PPARA activates BCMO1 to metabolize vitamin A. It also regulates the expression of SCARB1 (scavenger receptor class B type 1), which is a LDL (high-density lipoprotein) receptor protein that mediates carotenoid uptake (Hoekstra and Sorci-Thomas 2017). Soran et al. (2015) reported that HDL showed an antioxidant effect in protecting oxidative stress due to increased LDL (low-density lipoprotein). Figure 4 showed that lutein affects UCP1 (uncoupling protein 1) expression directly through activation by PPARA. UCP1 gene reduces the production of ROS (reactive oxygen species) in mitochondria (Oelkrug et al. 2014) to prevent oxidative stress in adipose tissue (Shabalina et al. 2006).

Figure 4 showed beta-carotene affects BDNF (Brain-derived neurotrophic factor) expression and it is a gene in the brain that promotes the survival of nerve cells (neurons) by playing a role in the growth, maturation (differentiation), and maintenance of neuron cells. Wu et al. (2016) reported that BDNF increases superoxide dismutases and glutathione reductase expression. It also reduces the oxidative protein damage index (Lee et al. 2009), reduces oxidant protein expression (Wu et al. 2012), restores reduced mitochondrial electron-coupling capacity, and increases mitochondrial uncoupling protein 2 (UCP2). It acts as an antioxidant by reducing the production of superoxide anions (Chan et al. 2010). Figure 4 also showed that beta-carotene and lutein inhibit the activation of the protein MMP9 (Matrix Metalloproteinase-9) through increased BDNF expression. Furthermore, the MMP9 gene regulates the tissue remodeling process by activating cytokines and chemokines, causing inflammation and fibrosis in the tissue (Yabluchanskiy et al. 2013). Figure 4 showed that beta-carotene activates FN1 (Fibronectin 1) action to inhibit the enzyme lipoygenases (LOX) and MMP9 expression. LOX enzyme oxidizes fatty acids and causes inflammation in tissues (Ratnasari et al. 2017). Meanwhile, FN1 is activated through the Nrf-2 pathway (Prestigiacomo and Suter-Dick 2018) to bind LOX. Oxidation and inflammation in the tissue can be inhibited (Fogelgren et al. 2005). In summary, the pigments in the leaves of *R. mucronata* work synergistically by activating antioxidant enzymes and inhibiting the expression of oxidative stress proteins.

The extract of *R. mucronata* leaves was evaluated for its antioxidant capacity as a free radical scavenger using the DPPH method. Meanwhile, DPPH is a stable free radical which accepts electrons or hydrogen atoms to form


stable diamagnetic molecules (Tanod et al. 2019a). The antioxidant capacity was evaluated by calculating DPPH purple light intensity level proportional to the decrease in DPPH concentration. This reduction was caused by the reaction of the 2,2-diphenyl-1-picrylhydrazyl molecule with the hydrogen atoms released by the components of the sample molecule. It formed hydrazone diphenyl picric compound and caused DPPH to change color from purple to yellow (Tanod et al. 2019b). The reactivity of the *R. mucronata* leaves extract with stable free radicals was also evaluated, and the antioxidant capacity was compared with vitamin C (Figure 5). In addition, Vitamin C is commonly used to compare assaying antioxidant activity because it is cheaper and easier to obtain (Lung and Destiani 2014).

Figure 5 showed that the *R. mucronata* leaves extract has antioxidant activity because hydrogen atoms or electrons were donated to react with DPPH radicals.

Increased concentration of the extract also increased the percentage of DPPH free radical inhibition. Table 2 showed that the inhibition percentage was evaluated for IC<sub>50</sub> determination by probit analysis. According to the Blois (1958) category, antioxidant activity can be categorized as very strong (IC<sub>50</sub> < 50 µg/mL), strong (50 ≤ IC<sub>50</sub> ≤ 100 µg/mL), moderate (100 ≤ IC<sub>50</sub> ≤ 150 µg/mL), weak (150 ≤ IC<sub>50</sub> ≤ 200 µg/mL), and very weak (IC<sub>50</sub> > 200 µg/mL).


**Table 2.** The IC<sub>50</sub> value of the *Rhizophora mucronata* leaves extract using the DPPH method was compared with vitamin C

Sample	IC <sub>50</sub> (µg/mL)
<i>R. mucronata</i> leaves	20.99 ± 0.33
Vitamin C	9.62 ± 0.09



**Figure 4.** Mechanism action as an antioxidant of pigments in *Rhizophora mucronata* leaves with STITCH analysis





**Figure 5.** DPPH radical inhibition from the *Rhizophora mucronata* leaves extract compared to vitamin C

Table 2 showed that the *R. mucronata* leaves extract and vitamin C had very strong antioxidant activity, with a DPPH concentration of 0.76 mM. Literature studies showed that the IC<sub>50</sub> of the same sample can vary depending on the DPPH concentration, sample origin, conditions, and the solvent used (Dewanto et al. 2019). Furthermore, fractionation and crude extract of *R. mucronata* mature leaves from Penunggul, East Java, Indonesia showed IC<sub>50</sub> from 82.97 ± 51.15 to 491.78 ± 427.59 µg/mL, while the IC<sub>50</sub> ascorbic acid (vitamin C) was 12.36 µg/mL, with a DPPH concentration of 0.4 mM (Sasmito et al. 2016). The ethanol extract obtained from Sunderban, I (23), was reported to show DPPH radical scavenging (IC<sub>50</sub> 23.65 ± 0.10 µg/mL, with a DPPH concentration of 0.135 mM (Adhikari et al. 2016). Two compounds isolated from the methanol extract of the leaves from Vallarpadam, India, showed IC<sub>50</sub> of 0.76-0.84 mg/mL, with DPPH 0.1 mM (Chakraborty and Raola 2017). Furthermore, the leaves extract from Tugurejo, Central Java, Indonesia, was reported to have IC<sub>50</sub> of 113.41 ppm (methanol), 151.13 ppm (n-hexane), 184.78 ppm (ethyl acetate), with a DPPH concentration of 0.1 mM (Ridlo et al. 2017). The ethanol extract obtained from the coast of Palu Bay showed an IC<sub>50</sub> of 103.95 ± 0.38 µg/mL, with a DPPH concentration of 50 µM (Dewanto et al. 2018). Furthermore, stick balm preparations from methanol extract of *R. mucronata* showed IC<sub>50</sub> of 57.40, 57.14, and 77.32 ppm, with a DPPH concentration of 0.004% (Faiqoh et al. 2020).

Literature studies reported varying IC<sub>50</sub> values of vitamin C with different DPPH concentrations. The IC<sub>50</sub> of vitamin C with a DPPH concentration of 0.1 mM, was 3.90 µg/mL (Pratiwi et al. 2013) and 4.76 µg/mL (Widyowati et al. 2014). Isnindar et al. (2016) reported an IC<sub>50</sub> with a DPPH concentration of 0.4 mM to be 1.83 µg/mL. Aliyu et al. (2017) reported an IC<sub>50</sub> with a DPPH concentration of 0.2 mM to be 13.89 µg/mL. Meanwhile, Johnson et al. (2017) reported an IC<sub>50</sub> with a DPPH concentration of 0.004% (w/v) to be 37.50 µg/mL. Najihudin et al. (2017)

reported an IC<sub>50</sub> with a DPPH concentration of 0.01% (w/v) to be 4.71 µg/mL. Dewanto et al. (2019) reported an IC<sub>50</sub> of vitamin C with a DPPH concentration of 50 µM to be 9.35 ± 0.81 µg/mL.

Mangroves are plants that live in tidal areas, and they experience variations in environmental pressure. In addition, variations in ecological pressure trigger reactive oxygen species (ROS) production and interfere with mangrove growth (Dasgupta et al. 2012). The pigments contained in mangroves produce antioxidant substances useful for the self-defense mechanism of ROS (Thatoi et al. 2014; Sarker and Oba 2018c; Dewanto et al. 2021). Therefore, it can be concluded that chlorophyll a, chlorophyll b, beta-carotene, lutein, neoxanthin, pheophytin a, and violaxanthin are the pigment profiles in *Rhizophora mucronata* mangrove leaves. All pigments detected have strong potential as antioxidants, especially as free radical scavengers and Nrf-2 stimulants. They interact with each other to inactivate antioxidant enzymes and inhibit the expression of oxidative stress proteins. This study showed that the pigment in *R. mucronata* leaves is a potential source of antioxidants for human health. Furthermore, it was developed by isolating the pigments from *Rhizophora mucronata* leaves and comparing the antioxidant power with standards for nutraceuticals and pharmaceuticals preparation.

#### ACKNOWLEDGEMENTS

The authors are grateful to the Rector of Universitas Sam Ratulangi, Manado, Indonesia for research grant No. 042.01.2.400959/2019 (Universitas Sam Ratulangi DIPA research funding), and are also grateful to Andreas Marbun and Maikel Tiolung for their assistance.

#### REFERENCES

- Adhikari A, Ray M, Das AK, Sur TK. 2016. Antidiabetic and antioxidant activity of *Rhizophora mucronata* leaves (Indian sundarban mangrove): An in vitro and in vivo study. *J Res Ayurveda* 37 (1): 76-81. DOI: 10.4103/ayu.AYU.
- Aisiah S, Olga, Tanod WA, Salosso Y, Bambang, Riyadi PH. 2020. Computational analysis of ethyl acetate extract of *Nuclea subdita* (Korth.) Steud. leaves as peptidoglycan glycosyltransferase inhibitor in *Aeromonas hydrophila*. *IOP Conf Ser: Earth Environ Sci* 584 (1): 012022. DOI: 10.1088/1755-1315/584/1/012022.
- Al-ali RM, Al-hilifi SAH, Al-Mossawi AEHJ. 2020. Extraction of lutein from some plant source in different conditions and application in food system. *Plant Arch* 20 (1): 1679-1685.
- Alhaddad ZA, Wahyudi D, Tanod WA. 2019. Antibacterial bioactivity of mangrove leaf extracts *Avicennia* sp. *Jurnal Kelautan: Indones J Mar Sci Technol* 12 (1): 12-22. DOI: 10.21107/jk.v12i1.4752. [Indonesian]
- Aliyu M, Abdullahi A, Ugya A. 2017. Antioxidant properties of selected poaceae species in Kano, Northern Nigeria. *Eur J Biomed Pharm Sci* 4 (5): 577-585.
- Aljaghthmi OH, Heba HM, Abu Zeid IM. 2017. Bioactive compounds extracted from mangrove plants (*Avicennia marina* and *Rhizophora mucronata*): an overview. *Adv Biol Res* 11: 161-170. DOI: 10.1016/j.pathophys.2018.09.002.
- Analuddin K, Septiana A, Nasaruddin, Sabilu Y, Sharma S. 2019. Mangrove fruit bioprospecting: nutritional and antioxidant potential

- as a food source for coastal communities in the Rawa Aopa Watumohai National Park, Southeast Sulawesi, Indonesia. *Intl J Fruit Sci* 19 (4): 423-436. DOI: 10.1080/15538362.2018.1555507.
- Andriani D, Revianti S, Prananingrum W. 2020. Identification of compounds isolated from a methanolic extract of *Acanthus ilicifolius* leaves and evaluation of their antifungal and antioxidant activity. *Biodiversitas* 21 (6): 2521-2525. DOI: 10.13057/biodiv/d210625.
- Araki M, Kaku N, Harada M, Ando Y, Yamaguchi R, Shindo K. 2016. Production of auroxanthins from violaxanthin and 9-cis-violaxanthin by acidic treatment and the antioxidant activities of violaxanthin, 9-cis-violaxanthin, and auroxanthins. *J Agric Food Chem* 64 (49): 9352-9355. DOI: 10.1021/acs.jafc.6b04506.
- Aruldass CA, Dufossé L, Ahmad WA. 2018. Current perspective of yellowish-orange pigments from microorganisms-a review. *J Cleaner Prod* 180: 168-182. DOI: 10.1016/j.jclepro.2018.01.093.
- Bandaranayake WM. 1998. Traditional and medicinal uses of mangroves. *Mangroves Salt Marshes* 2 (3): 133-148. DOI: 10.1023/A:1009988607044.
- Banerjee K, Chowdhury GR, Mitra A. 2017. Astaxanthin level of dominant mangrove floral species in Indian Sundarbans. *MOJ Bioorg Org Chem* 1 (2): 1-5. DOI: 10.15406/mojoc.2017.01.00008.
- Ben-dor A, Steiner M, Gheber L, Danilenko M, Dubi N, Limmewiel K, Zick A, Sharoni Y, Levy J. 2005. Carotenoids activate the antioxidant response element transcription system. *Mol Cancer Ther* 4 (1): 177-187.
- Berti AP, Düsman E, Mariucci RG, Lopes NB, Vicentini VEP. 2014. Antimutagenic and radioprotective activities of beta-carotene against the biological effects of iodine-131 radiopharmaceutical in wistar rats. *Genet Mol Res* 13 (1): 2248-2258. DOI: 10.4238/2014.March.31.5.
- Blaner WS, Brun PJ, Calderon RM, Golczak M. 2020. Retinol-binding protein 2 (RBP2): biology and pathobiology. *Crit Rev Biochem Mol Biol* 55 (2): 197-218. DOI: 10.1080/10409238.2020.1768207.
- Blois M. 1958. Antioxidant determinations by the use of a stable free radical. *Nature* 181 (4617): 1199-1200. DOI: 10.1038/1811199a0.
- Bogacz-Radomska L, Harasym J. 2018.  $\beta$ -Carotene-properties and production methods. *Food Qual Saf* 2 (2): 69-74. DOI: 10.1093/fqsafe/fyy004.
- Bohn T, Walczyk T. 2004. Determination of chlorophyll in plant samples by liquid chromatography using zinc-phthalocyanine as an internal standard. *J Chromatogr A* 1024 (1-2): 123-128. DOI: 10.1016/j.chroma.2003.10.067.
- Bowen-O'Connor CA, VanLeeuwen DM, Bettmann G, Sterling TM, Hilaire RST. 2013. Variation in violaxanthin and lutein cycle components in two provenances of *Acer grandidentatum* L. exposed to contrasting light. *Acta Physiol Plant* 35 (2): 541-548. DOI: 10.1007/s11738-012-1095-7.
- Chakraborty K, Raola VK. 2017. Two rare antioxidant and anti-inflammatory oleanenes from loop root Asiatic mangrove *Rhizophora mucronata*. *Phytochemistry* 135: 160-168. DOI: 10.1016/j.phytochem.2016.12.013.
- Chan SHH, Wu CWJ, Chang AYW, Hsu Ksen, Chan JYH. 2010. Transcriptional upregulation of brain-derived neurotrophic factor in rostral ventrolateral medulla by angiotensin II: significance in superoxide homeostasis and neural regulation of arterial pressure. *Circ Res* 107 (9): 1127-1139. DOI: 10.1161/CIRCRESAHA.110.225573.
- Chandrika UG, Jansz ER, Warnasuriya ND. 2005. Identification and HPLC quantification of carotenoids of the fruit pulp of *Chrysophyllum roxburghii*. *J Nat Sci Found Sri Lanka* 33 (2): 93-98. DOI: 10.4038/jnsf.v33i2.2337.
- Chen P, Li L, Gao Y, Xie Z, Zhang Y, Pan Z, Tu Y, Wang H, Han Q, Hu X, Xin X. 2019.  $\beta$ -carotene provides neuro protection after experimental traumatic brain injury via the Nrf2-ARE pathway. *J Integr Neurosci* 18: 153-161. DOI: 10.31083/jjin.2019.02.120.
- Croft H, Chen JM. 2017. Leaf Pigment Content. Elsevier Inc, Canada. DOI: 10.1016/B978-0-12-409548-9.10547-0.
- Cui T, Lai Y, Janicki JS, Wang X. 2016. Nuclear factor erythroid-2 related factor 2 (Nrf2)-mediated protein quality control in cardiomyocytes. *Front Biosci* 21: 192-202. DOI: 10.2741/4384.
- Dahdouh-Guebais F, Mathenge C, Kairo JG, Koedam N. 2000. Utilization of mangrove wood products around Mida Creek "Kenya" amongst subsistence and commercial users. *Econ Bot* 54 (4): 513-527. DOI: 10.1007/BF02866549.
- Dall'Osto L, Cazzaniga S, North H, Marion-Poll A, Bassia R. 2007. The arabidopsis aba4-1 mutant reveals a specific function for neoxanthin in protection against photooxidative stress. *Plant Cell* 19 (3): 1048-1064. DOI: 10.1105/tpc.106.049114.
- Dasgupta N, Nandy P, Sengupta C, Das S. 2012. Protein and enzymes regulations towards salt tolerance of some Indian mangroves in relation to adaptation. *Trees Struct Funct* 26 (2): 377-391. DOI: 10.1007/s00468-011-0599-x.
- Dewanto DK, Finarti F, Hermawan R, Ndobe S, Riyadi PH, Tanod WA. 2019. Antioxidant activity of soft coral extract from Palu Bay, Central Sulawesi, Indonesia. *Jurnal Pascapenan dan Bioteknologi Kelautan dan Perikanan* 14 (2): 163-178. DOI: 10.15578/jpbkp.v14i2.583 [Indonesian]
- Dewanto DK, Hermawan R, Muliadin, Riyadi PH, Aisiah S, Tanod WA. 2021. GC-MS profile of *Rhizophora apiculata* leaf extract from the coast of Tomini bay, Central Sulawesi with antibacterial and antioxidant activity. *Jurnal Kelautan: Indon J Mar Sci Techno* 14 (1): 30-42. [Indonesian]
- Dewanto DK, Tanod WA, Finarti F, Renol R. 2018. Screening of antiradical activity from some Central Sulawesi mangroves. *Pharmaciana* 8 (1): 155-168. DOI: 10.12928/pharmaciana.v8i1.8187.
- Dou Z, Cui L, Li J, Zhu Y, Gao C, Pan X, Lei Y, Zhang M, Zhao X, Li W. 2018. Hyperspectral estimation of the chlorophyll content in short-term and long-term restorations of Mangrove in Quanzhou Bay Estuary, China. *Sustainability* 10 (4): 1-15. DOI: 10.3390/su10041127.
- Durga Devi M, Banu N. 2015. Study of antioxidant activity of chlorophyll from some medicinal plants. *Indian J Res* 4 (2): 6-8.
- Eijkelhoff C, Dekker JP. 1997. A routine method to determine the chlorophyll a, pheophytin a and B-carotene contents of isolated Photosystem II reaction center complexes. *Photosynth Res* 52 (1): 69-73. DOI: 10.1023/A:1005834006985.
- Faiqoh M, Utami TFY, Pertiwi Y. 2020. Antioxidant assay for stick balm preparation of *Rhizophora mucronata* leaf extract using DPPH method. *Jurnal Ilmiah Jophus: J Pharma UMUS* 2 (1): 51-58. [Indonesian]
- Falqueto A, Silva DM, Fontes RV. 2008. Photosynthetic performance of mangroves *Rhizophora mangle* and *Laguncularia racemosa* under field conditions. *Rev Árvore* 32 (3): 577-582. DOI: 10.1590/S0100-6722008000300018.
- Filimonov DA, Lagunin AA, Glorizova TA, Rudik AV, Druzhilovskii DS, Pogodin PV, Poroikov VV. 2014. Prediction of the biological activity spectra of organic compounds using the pass online web resource. *Chem Heterocyclic Compd* 50 (3): 444-457. DOI: 10.1007/s10593-014-1496-1.
- Flores-de-Santiago F, Kovacs JM, Wang J, Flores-Verdugo F, Zhang C, González-Farías F. 2016. Examining the influence of seasonality, condition, and species composition on mangrove leaf pigment contents and laboratory based spectroscopy data. *Remote Sens* 8 (3): 226-246. DOI: 10.3390/rs8030226.
- Fogelgren B, Polgár N, Szauter KM, Újfaludi Z, Laczkó R, Fong KSK, Csizsar K. 2005. Cellular fibronectin binds to lysyl oxidase with high affinity and is critical for its proteolytic activation. *J Biol Chem* 280 (26): 24690-24697. DOI: 10.1074/jbc.M412979200.
- Gao Z, Li YH. 2012. Antioxidant stress and anti-inflammation of PPAR  $\alpha$  on warm hepatic ischemia-reperfusion injury. *PPAR Res* 2012: 1-8. DOI: 10.1155/2012/738785.
- Giossi CE, Cartaxana P, Cruz S. 2020. Photoprotective role of neoxanthin in plants and algae. *Molecules* 25 (20): 4617-4633. DOI: 10.3390/molecules25204617.
- Gupta P, Sreelakshmi Y, Sharma R. 2015. A rapid and sensitive method for determination of carotenoids in plant tissues by high performance liquid chromatography. *Plant Methods* 11 (1): 1-12. DOI: 10.1186/s13007-015-0051-0.
- Hegazi MM, Perez-ruzafa A, Almela L, Candela ME. 1998. Separation and identification of chlorophylls and carotenoids from *Caulerpa prolifera*, *Jania rubens* and *Padina pavonica* by reversed-phase high-performance liquid chromatography. *J Chromatogr A* 829 (1-2): 153-159. DOI: 10.1016/S0021-9673(98)00803-6.
- Henriques M, Silva A, Rocha J. 2007. Extraction and quantification of pigments from a marine microalga: a simple and reproducible method. In: Mendez-Vilas A (ed) *Communicating Current Research and Educational Topics and Trends in Applied Microbiology*. Formatex Research Center, Badajoz.
- Hoekstra M, Sorci-Thomas M. 2017. Rediscovering scavenger receptor type BI: surprising new roles for the HDL receptor. *Curr Opin Lipidol* 28 (3): 255-260. DOI: 10.1097/MOL.0000000000000413.
- Hsu CY, Chao PY, Hu SP, Yang CM. 2013. The antioxidant and free radical scavenging activities of chlorophylls and pheophytins. *Food Nutr Sci* 4: 1-8. DOI: 10.4236/fns.2013.48a001.

- Imdadul H, Wirakarnain S, Koshy P, Arash R, Shariff HABM, Mat TR. 2011. Valuable antioxidant and antimicrobial extracts from *Rhizophora mucronata* of Asiatic mangrove forests. *Res J Biotechnol* 6 (1): 10-14.
- İnanç AL. 2011. Chlorophyll: structural properties, health benefits and its occurrence in virgin olive oils. *Akademik Gıda* 9: 26-32.
- Ingkasupart P, Manochai B, Song WT, Hong JH. 2015. Antioxidant activities and lutein content of 11 marigold cultivars (*Tagetes* spp.) grown in Thailand. *Food Sci Technol* 35 (2): 380-385. DOI: 10.1590/1678-457X.6663.
- Isnindar, Wahyuono S, Widyarini S, Yuswanto. 2016. Determination of antioxidant activities of buas-buas leaves (*Premna serratifolia* L.) using DPPH (2,2-diphenyl-1-picrylhydrazyl) method. *Tradit Med J* 21 (3): 111-115. DOI: 10.22146/tradmedj.17292.
- Jariyah, Widjanarko SB, Yuniarta, Estiasih T, Sopade PA. 2014. Pasting properties mixtures of mangrove fruit flour (*Sonneratia caseolaris*) and starches. *Int Food Res J* 21 (6): 2161-2167.
- Jeffrey SW, Mantoura RFC, Wright SW. 1997. *Phytoplankton Pigments in Oceanography: Guidelines to Modern Methods*. UNESCO Publishing, Paris.
- Johnson E, Etim E, Archibong E. 2017. Isolation and anti-oxidant potentials of parahydroxybenzaldehyde from the methanol leaf extract of *Aspilia africana* (Pers.) C.D. Adams (Asteraceae). *Niger J Pharm Appl Sci Res* 6 (1): 26-32.
- Kamoshita M, Toda E, Osada H, Narimatsu T, Kobayashi S, Tsubota K, Ozawa Y. 2016. Lutein acts via multiple antioxidant pathways in the photo-stressed retina. *Sci Rep* 6 (1): 1-10. DOI: 10.1038/srep30226.
- Kardiman, Ridhwan M, Armi. 2017. Lindur fruit (*Bruguiera gymnorrhiza*) as food for the Acehnese community. *Serambi Sainia* 5 (2): 51-55. [Indonesian]
- Keleş D, Özgen Ş, Saraçoğlu O, Ata A, Özgen M. 2016. Antioxidant potential of Turkish pepper (*Capsicum annuum* L.) genotypes at two different maturity stages. *Turk J Agric For* 40 (4): 542-551. DOI: 10.3906/tar-1601-24.
- Kepel TL, Suryono DD, Ati RNA, Salim HL, Hutahaean AA. 2017. Important value and economic value estimation of carbon storage of mangrove vegetation in Kema, North Sulawesi. *Jurnal Kelautan Nasional* 12 (1): 19-26. DOI: 10.15578/jkn.v12i1.6170. [Indonesian]
- Kuhn M, Von Mering C, Campillos M, Jensen LJ, Bork P. 2008. STITCH: Interaction networks of chemicals and proteins. *Nucleic Acids Res* 36 (1): 684-688. DOI: 10.1093/nar/gkm795.
- Kumiawan JM, Yusuf MM, Heriyanto, Panintingjati TH, Brotosudarmo. 2020. Literature review on the potential of lutein from local marigold flowers as a health supplement. *Media Penelitian dan Pengembangan Kesehatan* 14 (2): 147-162. DOI: 10.22435/mpk.v30i2.2874. [Indonesian]
- Kusbandari A, Susanti H. 2017. Beta carotene content and free radical scavenging activity of cantaloupe (*Cucumis melo* var. *Cantalupensis* L.) extract against DPPH (1,1-diphenyl-2-picrylhydrazyl) using UV-visible spectrophotometry method. *Jurnal Farmasi Sains dan Komunitas* 14 (1): 37-42. DOI: 10.24071/jpsc.141562. [Indonesian]
- Kusmita L, Puspitaningrum I, Limantara L. 2015. Identification, isolation and antioxidant activity of pheophytin from green tea (*Camellia sinensis* (L.) Kuntze). *Procedia Chem* 14: 232-238. DOI: 10.1016/j.proche.2015.03.033.
- Landrum JT, Bone RA. 2001. Lutein, zeaxanthin, and the macular pigment. *Arch Biochem Biophys* 385 (1): 28-40. DOI: 10.1006/abbi.2000.2171.
- Lee B, Cao R, Choi YS, Cho HY, Rhee AD, Hah CK, Hoyt KR, Obrietan K. 2009. The CREB/CRE transcriptional pathway: protection against oxidative stress-mediated neuronal cell death. *J Neurochem* 108 (5): 1251-1265. DOI: 10.1111/j.1471-4159.2008.05864.x.
- Lichtenhaler HK. 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. In: Khan MU, Mitchell K (eds) *Methods in Enzymology*. Academic Press Inc., Cambridge. DOI: 10.1016/0076-6879(87)48036-1
- Lietz G, Lange J, Rimbach G. 2010. Molecular and dietary regulation of  $\beta$ , $\beta$ -carotene 15,15'-monooxygenase 1 (BCMO1). *Arch Biochem Biophys* 502 (1): 8-16. DOI: 10.1016/j.abb.2010.06.032.
- Lietz G, Oxley A, Boesch-Saadatmandi C, Kobayashi D. 2012. Importance of  $\beta$ , $\beta$ -carotene 15,15'-monooxygenase 1 (BCMO1) and  $\beta$ , $\beta$ -carotene 9',10'-dioxygenase 2 (BCDO2) in nutrition and health. *Mol Nutr Food Res* 56 (2): 241-250. DOI: 10.1002/mnfr.201100387.
- Limantara L, Heriyanto. 2012. Pigment composition and content of fucoxanthin brown seaweed from madura waters with high performance liquid chromatography. *Ilmu Kelautan: Indones J Mar Sci* 15 (1): 23-32. DOI: 10.14710/ik.ijms.15.1.23-32. [Indonesian]
- Liu Y. 2012. Spectral response to varying levels of leaf pigments collected from a degraded mangrove forest. *J Appl Remote Sens* 6 (1): 063501. DOI: 10.1117/1.jrs.6.063501
- Lung JKS, Destiani DP. 2014. Antioxidant activity test of vitamins A, C, E with the DPPH method. *Farmaka* 14 (1): 1-10. [Indonesian]
- Ma Q. 2013. Role of Nrf2 in oxidative stress and toxicity. *Annu Rev Pharmacol Toxicol* 53: 401-426. DOI: 10.1146/annurev-pharmtox-011112-140320.Role.
- Mangunsong S, Assiddiqy R, Sari EP, Marpaung PN, Sari RA. 2019. Determination of  $\beta$ -carotene in carrots (*Daucus carota*) by high performance liquid chromatography (U-HPLC). *ActTion: Aceh Nutr J* 4: 36-41. DOI: 10.30867/action.v4i1.151. [Indonesian]
- Mapari SAS, Nielsen KF, Larsen TO, Frisvad JC, Meyer AS, Thrane, U. 2005. Exploring fungal biodiversity for the production of water-soluble pigments as potential natural food colorants. *Curr Opin Biotechnol* 16 (2): 231-238. DOI: 10.1016/j.copbio.2005.03.004.
- Melendez-Martinez AJ, Britton G, Vicario IM, Heredia FJ. 2008. The complex carotenoid pattern of orange juices from concentrate. *Food Chem* 109 (3): 546-553. DOI: 10.1016/j.foodchem.2008.01.003.
- Moran E, Ding L, Wang Z, Cheng R, Chen Q, Moore R, Takahashi Y, Ma JX. 2014. Protective and antioxidant effects of PPAR $\alpha$  the ischemic retina. *Investig Ophthalmol Vis Sci* 55 (7): 4568-4576. DOI: 10.1167/iovs.13-13127.
- Mueller L, Boehm V. 2011. Antioxidant activity of  $\beta$ -carotene compounds in different in vitro assays. *Molecules* 16 (2): 1055-1069. DOI: 10.3390/molecules16021055.
- Najihudin A, Chaerunisaa A, Subarnas A. 2017. Antioxidant activity of trengguli (*Cassia Fistula* L.) bark extract and fraction using the DPPH method. *Indon J Pharm Sci Technol* 4 (2): 70-78. DOI: 10.15416/ijpst.v4i2.12354.
- Ngginak J, Mangibulude JC, Rondonuwu FS. 2017. The identification of carotenoids and testing of carotenoid antioxidants from sand lobster (*Panulirus homarus*) egg extract. *Ilmu Kelautan* 22 (3): 155-160. DOI: 10.14710/ik.ijms.22.3.155-160. [Indonesian]
- Oelkrug R, Goetze N, Meyer CW, Jastroch M. 2014. Antioxidant properties of UCPI are evolutionarily conserved in mammals and buffer mitochondrial reactive oxygen species. *Free Radic Biol Med* 77: 210-216. DOI: 10.1016/j.freeradbiomed.2014.09.004
- Oke JM, Hamburger O. 2002. Screening of some Nigerian medicinal plants for antioxidant activity using 2,2-diphenyl-picryl-hydrazyl radical. *Afr J Biomed Res* 5 (1-2): 77-79.
- Paolini M, Abdel-Rahman SZ, Sapone A, Pedulli GF, Perocco P, Cantelli-Forti G, Legator MS. 2003.  $\beta$ -carotene: a cancer chemopreventive agent or a co-carcinogen? *Mut Res-Rev Mut Res* 543 (3): 195-200. DOI: 10.1016/S1383-5742(03)00002-4.
- Paransa DSJ, Kemer K, Rumengan AP, Mantiri DM. 2014. Pigment type analysis and antibacterial activity assay of xanthophyll pigment extract on brown algae *Sargassum polycystum* (C. Agardh). *Jurnal LPPM Bidang Sains dan Teknologi* 1 (1): 90-96. [Indonesian]
- Pareek S, Sagar NA, Sharma S, Kumar V, Agarwal T, González-Aguilar GA, Yahia EM. 2017. Chlorophylls: chemistry and biological functions. In: Yahia EM (ed) *Fruit and Vegetable Phytochemicals: Chemistry and Human Health: Second Edition* (Vol. 1, 2<sup>nd</sup> ed). John Wiley & Sons Ltd, New Jersey. DOI: 10.1002/9781119158042.ch14.
- Pérez-gálvez A, Viera I, Roca M. 2020. Carotenoids and chlorophylls as antioxidants. *Antioxidants* 9 (6): 1-39. DOI: 10.3390/antiox9060505.
- Perreau F, Frey A, Effroy-Cuzzi D, Savane P, Berger A, Gissot L, Marion-Poll A. 2020. Abscisic acid-deficient4 has an essential function in both cis-violaxanthin and cis-neoxanthin synthesis. *Plant Physiol* 184 (3): 1303-1316. DOI: 10.1104/pp.20.00947.
- Prabhu KH, Bhute AS. 2015. Plant based dyes and mordant: a review. *J Nat Prod Plant Resour* 2 (6): 649-664.
- Pratiwi D, Wahdaningsih S, Isnindar. 2013. The test of antioxidant activity from bawang mekah leaves (*Eleutherine americana* Merr.) using DPPH (2,2-diphenyl-1-picrylhydrazyl) method. *Trad Med J* 18 (1): 9-16. DOI: 10.1007/s13398-014-0173-7-2.
- Prestigiacomo V, Suter-Dick L. 2018. Nrf2 protects stellate cells from Smad-dependent cell activation. *PLoS ONE* 13 (7): 1-18. DOI: 10.1371/journal.pone.0201044.
- Pringgienis D, Pratiwi AHD, Yudiati E, Azizah R, Susilo ES. 2017. Biopigment tracing of mangrove *Rhizophora mucronata* leaf and bark waste and its application in batik fabrics with various fixation methods. In: Dhramawan HA (eds.) *Proceedings Book The 7th Basic*

- Science International Conference. Universitas Brawijaya, Malang, 7-8 March 2017.
- Pringgienis D, Ridlo A, Dewi LF, Djunaedi A. 2018. The commercial value of mangrove-based pigments as natural dye for batik textiles delianis. *IntechOpen* (Vol. 32). IntechOpen, London.
- Puspitasari YE, Hartiati AM, Suprayitno E. 2012. The potency of *Rhizophora mucronata* leaf extract as anti-diarrhea. *J Appl Sci Res* 8 (2): 1180-1185.
- Radu GL, Litescu SC, Albu C, Teodor E, Truica G. 2012. Beta-carotene and lycopene determination in new enriched bakery products by HPLC-DAD method. *Romanian Biotechnol Lett* 17 (1): 7005-7012.
- Ratnasari N, Walters M, Tsopmo A. 2017. Antioxidant and lipoxigenase activities of polyphenol extracts from oat brans treated with polysaccharide degrading enzymes. *Heliyon* 3 (7): e00351. DOI: 10.1016/j.heliyon.2017.e00351.
- Ravikumar S, Gnanadesigan M. 2012. Hepatoprotective and antioxidant properties of *Rhizophora mucronata* mangrove plant in CCl<sub>4</sub> intoxicated rats. *J Exp Clin Med* 4 (1): 66-72. DOI: 10.1016/j.jecm.2011.11.012.
- Ravindran C, Naveenan T, Varatharajan GR, Rajasabapathy R, Meena RM. 2012. Antioxidants in mangrove plants and endophytic fungal associations. *Bot Mar* 55 (3): 269-279. DOI: 10.1515/bot-2011-0095.
- Revathi P, Jeyaseelan Senthinath T, Thirumalaikolundusubramanian P, Prabhu N. 2014. An overview of anti-diabetic profile of mangrove plants. *Int J Pharm Pharm Sci* 6 (3): 1-5.
- Ridlo A, Pramesti R, Koesoemadji K, Supriyantini E, Soenardjo N. 2017. Antioxidant activity of *Rhizophora mucronata* mangrove leaf extract. *Buletin Oseanografi Marina* 6 (2): 110-116. DOI: 10.14710/buloma.v6i2.16555. [Indonesian]
- Riyadi PH, Romadhon, Anggo AD, Suharto S, Tanod WA, Aryani A. 2021. Anti-inflammatory potential from tilapia (*Oreochromis niloticus*) viscera hydrolysate with bioinformatics analysis (prediction of activity spectra for substances – PASS). *IOP Conf Ser: Earth Environ Sci* 750 (1): 012044. DOI: 10.1088/1755-1315/750/1/012044.
- Riyadi PH, Tanod WA, Wahyudi D, Susanto E, Fahmi AS, Aisiah S. 2020. Potential of tilapia (*Oreochromis niloticus*) viscera bioactive peptides as antiviral for SARS-CoV-2 (COVID 19). *IOP Conf Ser: Earth Environ Sci* 584 (1): 012004. DOI: 10.1088/1755-1315/584/1/012004.
- Riyadi PH, Wahyudi D, Tanod WA. 2019. Effects of dichloromethane *Sarcophyton* spp. extract on the lipopolysaccharide-induced expression of nuclear factor-kappa B and inducible nitric oxide synthase in mice. *Vet World* 12 (12): 1897-1902. DOI: 10.14202/vetworld.2019.1897-1902.
- Rohini RM, Das AK. 2009. A comparative evaluation of analgesic and anti-inflammatory activities of *Rhizophora mucronata* bark extracts. *Pharmacologyonline* 1: 780-791.
- Rout P, Basak UC. 2014. Antioxidant properties in leaf and root extracts of some medicinally important mangrove species of Odisha coast. *Am J Pharmtech Res* 4: 605-617.
- Ruban AV, Pascal AA, Robert B, Horton P. 2001. Configuration and dynamics of xanthophylls in light-harvesting antennae of higher plants. *J Biol Chem* 276 (27): 24862-24870. DOI: 10.1074/jbc.M103263200.
- Rumengan AP, Mantiri DMH, Rompas R, Hutahaean A, Kepel TL, Paruntu CP, Kepel RC, Gerung GS. 2018. Carbon stock assessment of mangrove ecosystem in Totok bay, Southeast Minahasa regency, North Sulawesi, Indonesia. *AAEL Bioflux* 11 (4): 1280-1288.
- Şahin S, Aybastır Ö, Dawbaa S, Karkar B, Çakmak T. 2020. Study of the ability of lutein and neoxanthin as standards and in the extract of *Chlamydomonas reinhardtii* to prevent oxidatively induced DNA base damage using ultrasensitive GC-MS/MS analysis. *Chromatographia* 83: 919-926. DOI: 10.1007/s10337-020-03918-8.
- Saptiani G, Asikin AR, Ardhani F, Hardi EH. 2018. Mangrove plants species from Delta Mahakam, Indonesia with antimicrobial potency. *Biodiversitas* 19 (2): 516-521. DOI: 10.13057/biodiv/d190220.
- Sarker U, Islam MT, Rabbani MG, Oba S. 2014. Genotypic variability for nutrient, antioxidant, yield and yield contributing traits in vegetable amaranth. *J Food Agric Environ* 12: 168-174.
- Sarker U, Islam MT, Rabbani MG, Oba S. 2015a. Variability, heritability and genetic association in vegetable amaranth (*Amaranthus tricolor* L.). *Spanish J Agric Res* 13 (2): e0702. DOI: 10.5424/sjar/2015132-6843.
- Sarker U, Islam T, Rabbani G, Oba S. 2015b. Genotype variability in composition of antioxidant vitamins and minerals in vegetable amaranth. *Genetika* 47 (1): 85-96. DOI: 10.2298/GENSRI1501085S.
- Sarker U, Islam MT, Rabbani MG, Oba S. 2017. Genotypic diversity in vegetable amaranth for antioxidant, nutrient and agronomic traits. *Indian J Genet Plant Breed* 77: 173-176. DOI: 10.5958/0975-6906.2017.00025.6.
- Sarker U, Oba S. 2018a. Drought stress enhances nutritional and bioactive compounds, phenolic acids and antioxidant capacity of *Amaranthus* leafy vegetable. *BMC Plant Biol* 18 (1): 1-15. DOI: 10.1186/s12870-018-1484-1.
- Sarker U, Oba S. 2018b. Catalase, superoxide dismutase and ascorbate-glutathione cycle enzymes confer drought tolerance of *Amaranthus tricolor*. *Sci Rep* 8 (1): 1-12. DOI: 10.1038/s41598-018-34944-0.
- Sarker U, Oba S. 2018c. Drought stress effects on growth, ROS markers, compatible solutes, phenolics, flavonoids, and antioxidant activity in *Amaranthus tricolor*. *Appl Biochem Biotechnol* 186 (4): 999-1016. DOI: 10.1007/s12010-018-2784-5.
- Sarker U, Oba S. 2018d. Augmentation of leaf color parameters, pigments, vitamins, phenolic acids, flavonoids and antioxidant activity in selected *Amaranthus tricolor* under salinity stress. *Sci Rep* 8 (1): 1-9. DOI: 10.1038/s41598-018-30897-6.
- Sarker U, and Oba S. 2018e. Response of nutrients, minerals, antioxidant leaf pigments, vitamins, polyphenol, flavonoid and antioxidant activity in selected vegetable amaranth under four soil water content. *Food Chem* 252: 72-83. DOI: 10.1016/j.foodchem.2018.01.097.
- Sarker U, Islam MT, Oba S. 2018a. Salinity stress accelerates nutrients, dietary fiber, minerals, phytochemicals and antioxidant activity in *Amaranthus tricolor* leaves. *PLoS ONE* 13 (11): e0206388. DOI: 10.1371/journal.pone.0206388.
- Sarker U, Islam MT, Rabbani MG, Oba S. 2018b. Phenotypic divergence in vegetable amaranth for total antioxidant capacity, antioxidant profile, dietary fiber, nutritional and agronomic traits. *Acta Agric Scand Sect B: Soil Plant Sci* 68 (1): 67-76. DOI: 10.1080/09064710.2017.1367029.
- Sarker U, Islam MT, Rabbani MG, Oba S. 2018c. Antioxidant leaf pigments and variability in vegetable amaranth. *Genetika* 50 (1): 209-220. DOI: 10.2298/GENSRI1801209S.
- Sarker U, Islam MT, Rabbani MG, Oba S. 2018d. Variability in total antioxidant capacity, antioxidant leaf pigments and foliage yield of vegetable amaranth. *J Integrative Agric* 17 (5): 1145-1153. DOI: 10.1016/S2095-3119(17)61778-7.
- Sarker U, Oba S. 2019a. Antioxidant constituents of three selected red and green color *Amaranthus* leafy vegetable. *Sci Rep* 9 (1): 1-11. DOI: 10.1038/s41598-019-52033-8.
- Sarker U, Oba S. 2019b. Nutraceuticals, antioxidant pigments, and phytochemicals in the leaves of *Amaranthus spinosus* and *Amaranthus viridis* weedy species. *Sci Rep* 9 (1): 1-10. DOI: 10.1038/s41598-019-50977-5.
- Sarker U, Oba S. 2019c. Protein, dietary fiber, minerals, antioxidant pigments and phytochemicals, and antioxidant activity in selected red morph *Amaranthus* leafy vegetable. *PLoS ONE* 14 (12): e0222517. DOI: 10.1371/journal.pone.0222517.
- Sarker U, Oba S. 2019d. Salinity stress enhances color parameters, bioactive leaf pigments, vitamins, polyphenols, flavonoids and antioxidant activity in selected *Amaranthus* leafy vegetables. *J Sci Food Agric* 99 (5): 2275-2284. DOI: 10.1002/jsfa.9423.
- Sarker U, Hossain MN, Iqbal MA, Oba S. 2020a. Bioactive components and radical scavenging activity in selected advance lines of salt-tolerant vegetable amaranth. *Front Nutr* 7: 1-15. DOI: 10.3389/fnut.2020.587257.
- Sarker U, Oba S, Daramy MA. 2020b. Nutrients, minerals, antioxidant pigments and phytochemicals, and antioxidant capacity of the leaves of stem amaranth. *Sci Rep* 10 (1): 1-9. DOI: 10.1038/s41598-020-60252-7.
- Sarker U, Hossain MM, Oba S. 2020c. Nutritional and antioxidant components and antioxidant capacity in green morph *Amaranthus* leafy vegetable. *Sci Rep* 10 (1): 1-10. DOI: 10.1038/s41598-020-57687-3.
- Sarker U, Oba S. 2020a. The response of salinity stress-induced *A. tricolor* to growth, anatomy, physiology, non-enzymatic and enzymatic antioxidants. *Front Plant Sci* 11: 1-14. DOI: 10.3389/fpls.2020.559876.
- Sarker U, Oba S. 2020b. Phenolic profiles and antioxidant activities in selected drought-tolerant leafy vegetable amaranth. *Sci Rep* 10 (1): 1-11. DOI: 10.1038/s41598-020-71727-y.

- Sarker U, Oba S. 2020c. Polyphenol and flavonoid profiles and radical scavenging activity in leafy vegetable *Amaranthus gangeticus*. BMC Plant Biol 20 (1): 1-12. DOI: 10.1186/s12870-020-02700-0.
- Sarker U, Oba S. 2020d. Nutraceuticals, phytochemicals, and radical quenching ability of selected drought-tolerant advance lines of vegetable amaranth. BMC Plant Biol 20 (1): 1-16. DOI: 10.1186/s12870-020-02780-y.
- Sarker U, Oba S. 2020e. Nutritional and bioactive constituents and scavenging capacity of radicals in *Amaranthus hypochondriacus*. Sci Rep 10 (1): 1-10. DOI: 10.1038/s41598-020-71714-3.
- Sarker U, Oba S. 2020f. Nutrients, minerals, pigments, phytochemicals, and radical scavenging activity in *Amaranthus blitum* leafy vegetables. Sci Rep 10 (1): 1-9. DOI: 10.1038/s41598-020-59848-w.
- Sarker U, Oba S. 2020g. Leaf pigmentation, its profiles and radical scavenging activity in selected *Amaranthus tricolor* leafy vegetables. Sci Rep 10 (1): 1-10. DOI: 10.1038/s41598-020-66376-0.
- Sarker U, Oba S. 2021. Color attributes, betacyanin, and carotenoid profiles, bioactive components, and radical quenching capacity in selected *Amaranthus gangeticus* leafy vegetables. Sci Rep 11: 1-14. DOI: 10.1038/s41598-021-91157-8.
- Sasamoto H, Hayatsu M, Suzuki S. 2020. High allelopathic activity of carotenoid-accumulating callus of a halophilic mangrove plant, *Avicennia alba*: protoplast co-culture method with digital image analysis. J Plant Stud 9 (1): 1-12. DOI: 10.5539/jps.v9n1p1.
- Sasmito BB, Puspitasari YE, Hardoko H. 2016. Antidiabetic and antioxidant activities of tannin extract of *Rhizophora mucronata* leaves. J Chem Pharm Res 8 (3): 143-148.
- Shabalina IG, Petrovic N, Kramarova TV, Hoeks J, Cannon B, Nedergaard J. 2006. UCP1 and defense against oxidative stress: 4-hydroxy-2-nonenal effects on brown fat mitochondria are uncoupling protein 1-independent. J Biol Chem 281 (20): 13882-13893. DOI: 10.1074/jbc.M601387200.
- Soran H, Schofield JD, Durrington PN. 2015. Antioxidant properties of HDL. Front Pharmacol 6 (22): 1-6. DOI: 10.3389/fphar.2015.00222.
- Strobel M, Tinz J, Biesalski HK. 2007. The importance of  $\beta$ -carotene as a source of vitamin A with special regard to pregnant and breastfeeding women. Eur J Nutr 46 (1): 1-20. DOI: 10.1007/s00394-007-1001-z.
- Suganthi N, Pandima Devi K. 2016. In vitro antioxidant and anticholinesterase activities of *Rhizophora mucronata*. Pharma Biol 54 (1): 118-129. DOI: 10.3109/13880209.2015.1017886.
- Taiz L, Zeiger E, Møller IM, Murphy A. 2006. Plant Physiology (Fifth). Sinauer Associates Inc., Massachusetts.
- Tanod WA, Yanuhar U, Maftuch, Wahyudi D, Risjani Y. 2019a. DPPH scavenging property of bioactives from soft corals origin Palu bay, Central Sulawesi, Indonesia. IOP Conf Ser: Earth Environ Sci 236 (1): 012121. DOI: 10.1088/1755-1315/236/1/012121.
- Tanod WA, Dewanto DK, Ndobe S, Riyadi PH, Putra MY. 2019b. Screening of antibacterial and antioxidant activity of soft corals *Sinularia* sp. and *Sarcophyton* sp. from Palu bay, Central Sulawesi, Indonesia. Squalen: Bull Mar Fish Postharvest Biotechnol 14: 73-83. DOI: 10.15578/squalen.v14i2.394.
- Thatoi HN, Patra JK, Das SK. 2014. Free radical scavenging and antioxidant potential of mangrove plants: a review. Acta Physiol Plant 36 (3): 561-579. DOI: 10.1007/s11738-013-1438-z.
- Tidore F, Rumengan AP, Sondak CFA, Mangindaan REP, Runtuwene HCC, Pratasik SB. 2018. Estimation of carbon (C) content in mangrove leaves from Lansia village, Wori sub-district, North Minahasa district. Jurnal Pesisir dan Laut Tropis 6 (2): 53-58. DOI: 10.35800/jplt.6.2.2018.21529. [Indonesian]
- Wang F, Huang L, Gao B, Zhang C. 2018. Optimum production conditions, purification, identification, and antioxidant activity of violaxanthin from Microalga *Eustigmatois* cf. *polyphem* (Eustigmatophyceae). Mar Drugs 16 (6): 190-203. DOI: 10.3390/md16060190.
- Widyowati H, Ulfah M, Sumantri. 2014. Antioxidant activity test of the herba alfalfa ethanolic extract (*Medicago sativa* L.) with the DPPH method (1,1-Diphenyl-2-Picrylhydrazyl). Jumal Ilmu Farmasi dan Farmasi Klinik 11: 25-33. [Indonesian]
- Wu CL, Chen SD, Yin JH, Hwang CS, Yang DI. 2016. Nuclear factor-kappaB-dependent sestrin2 induction mediates the antioxidant effects of BDNF against mitochondrial inhibition in rat cortical neurons. Mol Neurobiol 53 (6): 4126-4142. DOI: 10.1007/s12035-015-9357-1.
- Wu CL, Yin JH, Hwang CS, Chen SD, Yang DY, Yang DI. 2012. C-Jun-dependent sulfiredoxin induction mediates BDNF protection against mitochondrial inhibition in rat cortical neurons. Neurobiol Dis 46 (2): 450-462. DOI: 10.1016/j.nbd.2012.02.010.
- Yabluchanskiy A, Ma Y, Iyer RP, Hall ME, Lindsey ML. 2013. Matrix metalloproteinase-9: many shades of function in cardiovascular disease. Physiology 28 (6): 391-403. DOI: 10.1152/physiol.00029.2013.
- Zapata M, Rodríguez F, Garrido JL. 2000. Separation of chlorophylls and carotenoids from marine phytoplankton: a new HPLC method using a reversed phase C8 column and pyridine-containing mobile phases. Mar Ecol Prog Ser 195: 29-45. DOI: 10.3354/meps195029.
- Zhang T, Wang Z, Wang X, Sun W, Cui X, Li R, Li G. 2019. Effects of vitamin A on antioxidant functions, immune functions and production performance in male sika deer (*Cervus nippon*) during the first antler growth period. Ital J Anim Sci 18: 98-104. DOI: 10.1080/1828051X.2018.1456978.

# Identification of pigment profiles and antioxidant activity of *Rhizophora mucronata* mangrove leaves origin Lembeh, North Sulawesi, Indonesia

## ORIGINALITY REPORT

**20%**  
SIMILARITY INDEX

**12%**  
INTERNET SOURCES

**17%**  
PUBLICATIONS

**4%**  
STUDENT PAPERS

## PRIMARY SOURCES

- |          |   |           |
|----------|---|-----------|
| <b>1</b> | <a href="http://www.mdpi.com">www.mdpi.com</a><br>Internet Source   | <b>1%</b> |
| <b>2</b> | <a href="http://journal.uad.ac.id">journal.uad.ac.id</a><br>Internet Source   | <b>1%</b> |
| <b>3</b> | Dario Rueda, Henry O. Awika, Renesh Bedre, Devi R. Kandel, Kranthi K. Mandadi, Kevin Crosby, Carlos A. Avila. "Phenotypic Diversity and Association Mapping of Ascorbic Acid Content in Spinach", <i>Frontiers in Genetics</i><br>Internet Source | <b>1%</b> |
| <b>4</b> | "Anticancer plants: Properties and Application", Springer Science and Business Media LLC, 2018<br>Publication   | <b>1%</b> |
| <b>5</b> | S Aisiah, Olga, W A Tanod, Y Salosso, Bambang, P H Riyadi. " Computational analysis of ethyl acetate extract of ( <i>Korth.</i> ) <i>Steud.</i> leaves as peptidoglycan glycosyltransferase inhibitor in ", <i>IOP</i>                            | <b>1%</b> |

# Conference Series: Earth and Environmental Science, 2020

Publication

- 
- |    |   |      |
|----|---|------|
| 6  | Wendy Alexander Tanod, Didit Kustantio Dewanto, Samliok Ndobe, Putut Har Riyadi, Masteria Yunovilsa Putra. "Screening of Antibacterial and Antioxidant Activity from the Soft Corals Sinularia sp. and Sarcophyton sp. Origin Palu Bay, Central Sulawesi, Indonesia", Squalen Bulletin of Marine and Fisheries Postharvest and Biotechnology, 2019<br>Publication | 1 %  |
| 7  | Edible Medicinal And Non-Medicinal Plants, 2014.<br>Publication   | <1 % |
| 8  | M. Ezziyani. "Biological Control of Phytophthora Root Rot of Pepper Using Trichoderma harzianum and Streptomyces rochei in Combination", Journal of Phytopathology, 6/2007<br>Publication   | <1 % |
| 9  | <a href="http://www.bioflux.com.ro">www.bioflux.com.ro</a><br>Internet Source   | <1 % |
| 10 | <a href="http://www.researchsquare.com">www.researchsquare.com</a><br>Internet Source   | <1 % |
| 11 | <a href="http://www.scirp.org">www.scirp.org</a><br>Internet Source   | <1 % |
-

12	Submitted to Universiti Teknologi Malaysia Student Paper	<1 %
13	Submitted to Western Governors University Student Paper	<1 %
14	hdl.handle.net Internet Source	<1 %
15	1library.net Internet Source	<1 %
16	v2.smujo.id Internet Source	<1 %
17	"Wild Fruits: Composition, Nutritional Value and Products", Springer Science and Business Media LLC, 2019 Publication	<1 %
18	Chia-Lin Wu, Shang-Der Chen, Jiu-Haw Yin, Chi-Shin Hwang, Ding-I Yang. "Nuclear Factor-kappaB-Dependent Sestrin2 Induction Mediates the Antioxidant Effects of BDNF Against Mitochondrial Inhibition in Rat Cortical Neurons", Molecular Neurobiology, 2015 Publication	<1 %
19	Submitted to Florida State University Student Paper	<1 %
20	Muhammad M Hegazi, Angel Pérez-Ruzafa, Luis Almela, María-Emilia Candela.	<1 %



"Separation and identification of chlorophylls and carotenoids from *Caulerpa prolifera*, *Jania rubens* and *Padina pavonica* by reversed-phase high-performance liquid chromatography", *Journal of Chromatography A*, 1998

Publication

21

[api.intechopen.com](http://api.intechopen.com)

Internet Source

<1 %

22

[ebin.pub](http://ebin.pub)

Internet Source

<1 %

23

[journal.trunojoyo.ac.id](http://journal.trunojoyo.ac.id)

Internet Source

<1 %

24

Martha Reyes-Becerril, Perla Giner, Jorge Silva-Jara, Adriana Macias et al. "Assessment of chemical, biological and immunological properties of "Damiana de California" *Turnera diffusa* Willd extracts in Longfin yellowtail (*Seriola rivoliana*) leukocytes", *Fish & Shellfish Immunology*, 2020

Publication

<1 %

25

[jmhg.springeropen.com](http://jmhg.springeropen.com)

Internet Source

<1 %

26

[www.ukm.my](http://www.ukm.my)

Internet Source

<1 %

27 Zhenyu Ding, Cuiyu Zhang, Baokun Zhang, Qin Li. "Unraveling the Proteomic Landscape of Intestinal Epithelial Cell-Derived Exosomes in Mice", *Frontiers in Physiology*, 2022

Publication

<1 %

28 [dokumen.pub](#)

Internet Source

<1 %

29 [gitisa.it](#)

Internet Source

<1 %

30 [real-d.mtak.hu](#)

Internet Source

<1 %

31 [www.pubfacts.com](#)

Internet Source

<1 %

32 K. Sahin, C. Orhan, M.O. Smith, N. Sahin. "Molecular targets of dietary phytochemicals for the alleviation of heat stress in poultry", *World's Poultry Science Journal*, 2019

Publication

<1 %

33 Lia Kusmita, Ika Puspitaningrum, Leenawaty Limantara. "Identification, Isolation and Antioxidant Activity of Pheophytin from Green Tea (*Camellia Sinensis* (L.) Kuntze)", *Procedia Chemistry*, 2015

Publication

<1 %

34 [www.plantcell.org](#)

Internet Source

<1 %

35

Barthlomew Yonas Chataika, Levi Shadeya-Mudogo Akundabweni, Aristide Carlos Houdegbe, Enoch G. Achigan-Dako et al. "Dietary Phytochemical Screening of Spider Plant (*Gynandropsis gynandra* (L.) Briq.) Accessions From Africa and Asia to Identify Genotypes for Use in Nutraceutical Breeding", *Frontiers in Sustainable Food Systems*, 2021

Publication

---

<1 %

36

Brian Sheng Xian Teo, Rui Yi Gan, Sarah Abdul Aziz, Thanchanok Sirirak, Mohd Fadli Mohd Asmani, Eddy Yusuf. " evaluation of antioxidant and antibacterial activities of extract and its evaluation of the wound healing activity in mice ", *Journal of Cosmetic Dermatology*, 2020

Publication

---

<1 %

37

Chinnarajan Ravindran, Thangaiah Naveenan, Govindaswamy R. Varatharajan, Raju Rajasabapathy, Ram Murthi Meena. "Antioxidants in mangrove plants and endophytic fungal associations", *Botanica Marina*, 2012

Publication

---

<1 %

38

Junjie Wang, Yi Xu, Guofeng Wu. "The integration of species information and soil properties for hyperspectral estimation of leaf

<1 %

biochemical parameters in mangrove forest",  
Ecological Indicators, 2020

Publication

---

39

Meenakshi Sharma, Prashant Kaushik.  
"Vegetable phytochemicals: An update on  
extraction and analysis techniques",  
Biocatalysis and Agricultural Biotechnology,  
2021

Publication

---

<1 %

40

Shyam Ramachandran, Luka A. Clarke, Todd  
E. Scheetz, Margarida D. Amaral, Paul B.  
McCray. "Chapter 12 Microarray mRNA  
Expression Profiling to Study Cystic Fibrosis",  
Springer Science and Business Media LLC,  
2011

Publication

---

<1 %

41

Sunil Pareek, Narashans Alok Sagar, Sunil  
Sharma, Vinay Kumar, Tripti Agarwal, Gustavo  
A. González-Aguilar, Elhadi M. Yahia.  
"Chlorophylls: Chemistry and Biological  
Functions", Wiley, 2017

Publication

---

<1 %

42

[journal.ugm.ac.id](http://journal.ugm.ac.id)

Internet Source

---

<1 %

43

[mjfas.utm.my](http://mjfas.utm.my)

Internet Source

---

<1 %

44

[www.biorxiv.org](http://www.biorxiv.org)

Internet Source

<1 %

45

[www.scilit.net](http://www.scilit.net)

Internet Source

<1 %

46

[www.tandfonline.com](http://www.tandfonline.com)

Internet Source

<1 %

47

A. Stier, P. Bize, C. Habold, F. Bouillaud, S. Massemin, F. Criscuolo. "Mitochondrial uncoupling prevents cold-induced oxidative stress: a case study using UCP1 knockout mice", *Journal of Experimental Biology*, 2013

Publication

<1 %

48

A.P. Berti, E. Düsman, R.G. Mariucci, N.B. Lopes, V.E.P. Vicentini. "Antimutagenic and radioprotective activities of beta-carotene against the biological effects of iodine-131 radiopharmaceutical in Wistar rats", *Genetics and Molecular Research*, 2014

Publication

<1 %

49

Clare Annabel Bowen-O'Connor, Dawn M. VanLeeuwen, Greg Bettmann, Tracy M. Sterling, Rolston St. Hilaire. "Variation in violaxanthin and lutein cycle components in two provenances of *Acer grandidentatum* L. exposed to contrasting light", *Acta Physiologiae Plantarum*, 2012

Publication

<1 %

50 Danuše Tarkowská, Miroslav Strnad. "Isoprenoid-derived plant signaling molecules: biosynthesis and biological importance", *Planta*, 2018  
Publication <1 %

---

51 Fatemeh Khavari, Fateme Asadi, Fatemeh Nouri, Mohammad Taheri, Fateme Mohammadi, Mojdeh Mohammadi, Parisa Habibi, Babak Asghari. "Marine antioxidants from microalgae", Elsevier BV, 2023  
Publication <1 %

---

52 Higor Castro Oliveira, Maria Cristina de Oliveira, Uilcimar Martins Arantes, Early Theodoro Alves de Argyri. "Paprika and/or marigold extracts improve productivity and yolk color in egg-laying quails", *Ciência Animal Brasileira*, 2020  
Publication <1 %

---

53 [academic.oup.com](https://academic.oup.com)  
Internet Source <1 %

---

54 [ejournal.undip.ac.id](https://ejournal.undip.ac.id)  
Internet Source <1 %

---

55 [www.iobc-wprs.org](https://www.iobc-wprs.org)  
Internet Source <1 %

---

56 "Mangroves: Ecology, Biodiversity and Management", Springer Science and Business Media LLC, 2021 <1 %

57

Ahmed M. M. Youssef, Doaa A. M. Maaty, Yousef M. Al-Saraireh. "Phytochemistry and Anticancer Effects of Mangrove (*Rhizophora mucronata* Lam.) Leaves and Stems Extract against Different Cancer Cell Lines", *Pharmaceuticals*, 2022

Publication

---

<1 %

58

Dave Paladin Buenavista, Eefke Maria Mollee, Morag McDonald. "Any alternatives to rice? Ethnobotanical insights into the dietary use of edible plants by the Higaonon tribe in Bukidnon Province, the Philippines", *Regional Sustainability*, 2022

Publication

---

<1 %

59

Fu, H.. "Evaluation of antioxidant activities of principal carotenoids available in water spinach (*Ipomoea aquatica*)", *Journal of Food Composition and Analysis*, 201103

Publication

---

<1 %

60

Kalpesh R. Patil, Purusottam Mohapatra, Harun M. Patel, Sameer N. Goyal, Shreesh Ojha, Chanakya N. Kundu, Chandragouda R. Patil. "Pentacyclic Triterpenoids Inhibit IKK $\beta$  Mediated Activation of NF- $\kappa$ B Pathway: In Silico and In Vitro Evidences", *PLOS ONE*, 2015

Publication

---

<1 %

61

Kusmiati Kusmiati, Fifi Afiati, Chrisna Widhiani, Alpinna Aditia, Destia D. Elviani, Atit Kanti. "The Potential of Lutein Extract of L. Flower as an Antioxidant and Enhancing Phagocytic Activity of Macrophage Cells ", IntechOpen, 2021

Publication

&lt;1 %

62

María Isabel Iñiguez-Luna, Jorge Cadena-Iñiguez, Ramón Marcos Soto-Hernández, Francisco Javier Morales-Flores et al. "Natural Bioactive Compounds of Sechium spp. for Therapeutic and Nutraceutical Supplements", Frontiers in Plant Science, 2021

Publication

&lt;1 %

63

Norain Jamalluddin, Rachael C. Symonds, Sean Mayes, Wai Kuan Ho, Festo Massawe. "Diversifying crops for food and nutrition security: A case of vegetable amaranth, an ancient climate-smart crop", Elsevier BV, 2021

Publication

&lt;1 %

64

Paul D. Fraser, Eugenia M.A. Enfissi, John M. Halket, Mark R. Truesdale, Dongmei Yu, Christopher Gerrish, Peter M. Bramley. "Manipulation of Phytoene Levels in Tomato Fruit: Effects on Isoprenoids, Plastids, and Intermediary Metabolism", The Plant Cell, 2007

Publication

&lt;1 %



65 Saikat Mitra, Abdur Rauf, Abu Montakim Tareq, Shamima Jahan et al. "Potential health benefits of carotenoid lutein: An updated review", Food and Chemical Toxicology, 2021  
Publication <1 %

---

66 Shan Gao, Runhua Li, Nuo Heng, Yu Chen et al. "Effects of dietary supplementation of natural astaxanthin from Haematococcus pluvialis on antioxidant capacity, lipid metabolism, and accumulation in the egg yolk of laying hens", Poultry Science, 2020  
Publication <1 %

---

67 backend.orbit.dtu.dk  
Internet Source <1 %

---

68 eprints.uthm.edu.my  
Internet Source <1 %

---

69 estudogeral.sib.uc.pt  
Internet Source <1 %

---

70 livrepository.liverpool.ac.uk  
Internet Source <1 %

---

71 mspp.org.my  
Internet Source <1 %

---

72 ouci.dntb.gov.ua  
Internet Source <1 %

---

73 www.eurchembull.com  
Internet Source <1 %

---

74

Josie A. Silvaroli, Jacqueline Plau, Charlie H. Adams, Surajit Banerjee et al. "Molecular basis for the interaction of cellular retinol binding protein 2 (CRBP2) with nonretinoid ligands", Journal of Lipid Research, 2021

Publication

&lt;1 %

75

"Ecophysiology, Abiotic Stress Responses and Utilization of Halophytes", Springer Science and Business Media LLC, 2019

Publication

&lt;1 %

76

Carotenoids and Human Health, 2013.

Publication

&lt;1 %

77

Julieta Saba, Juan Turati, Delia Ramírez, Lila Carniglia, Daniela Durand, Mercedes Lasaga, Carla Caruso. "Astrocyte truncated-TrkB mediates BDNF antiapoptotic effect leading to neuroprotection", Journal of Neurochemistry, 2018

Publication

&lt;1 %

78

Morad Bornavard, Hamed Fanaei, Mohammad Ali Mirshekar, Farzaneh Farajian Mashhadi, Alireza Atashpanjeh. "Morphine consumption during pregnancy exacerbates neonatal hypoxia - ischemia injury in rats", International Journal of Developmental Neuroscience, 2020

Publication

&lt;1 %

79

Xiaoming Gong. BMC Molecular Biology, 2006

Publication

<1 %

---

80

van Breemen, R.B.. "Multitargeted therapy of cancer by lycopene", Cancer Letters, 20081008

Publication

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On