

Pigment concentration of red algae, *Kappaphycus alvarezii* (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia

by Darus Sa'adah Johanis Paransa 5


Submission date: 30-May-2023 08:22AM (UTC+0700)

Submission ID: 2104891194

File name: ASIONAL_-_2020_-_DarusSaadah_J_Paransa_Pigment_concentrasion.pdf (534.18K)

Word count: 5289

Character count: 28128



5

Pigment concentration of red algae, *Kappaphycus alvarezii* (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia

Darus S. J. Paransa, Desy M. H. Mantiri, Rene C. Kepel, Antonius Rumengan

14 Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado, North Sulawesi, Indonesia. Corresponding author: D. S. J. Paransa, darusparansa@unsrat.ac.id

Abstract. Cultivation of brownish red algae *Kappaphycus alvarezii* in controlled conditions was conducted in order to obtain information about the growth and concentration of chlorophyll and carotenoid pigments. The cultivation was carried out for 9 weeks namely from March to April 2020 in coastal waters of Nain Island that was quite far from the mainland of Manado City. Pigment concentrations were obtained through extraction based on different polarity in the total pigment extract in petroleum ether added with methanol, KOH in methanol and diethyl ether. Red algae as much as 17 thalli with an initial weight of 100 g were cultured using floating method. The distance between thallus was 25 cm. A total of three thalli was taken every two weeks, namely at week 3, 5, 7, and 9 and grouped into 3 periods. After measuring the weight gain, the algae were then exposed to pigment extraction. *K. alvarezii* experienced weight gain, but the relative growth rate showed a decrease in the 3rd period. The pattern of algae relative growth was identical with the pattern of temperature fluctuation during cultivation. Pigments obtained from the separation using different solvents were chlorophyll a, chlorophyll d, β -carotene and xanthophyll. It was found that the concentration of chlorophyll a and d increased as the weight gain increased. Likewise, the concentration and quantity of β -carotene and xanthophyll pigments were in line with the growth rate pattern of *K. alvarezii*.

Key Words: macroalgae, chlorophyll, carotenoid, carotene, xanthophyll.

7

Introduction. *Kappaphycus alvarezii* is a red macroalgae or seaweed that has an important economic value. This macroalgae is a producer of kappa-carrageenan so that it can be a source of raw material for carrageenan extraction to be used in various industries of food, cosmetics, pharmaceuticals and other industrial applications (Pereira 2016). The carrageenan content of *K. alvarezii* originating from Arakan Village, North Sulawesi has immunostimulating activity (Kreckhoff et al 2019). Nain Island, located close to the capital city of North Sulawesi Province, is one of the macroalgae cultivation areas in North Sulawesi. One type of macroalgae cultivated in this area is *K. alvarezii*.

K. alvarezii has round thalli, cylindrical and flattened, smooth surface, branched irregularly, and has spines that vary in length from 1 to 17 mm (Pereira 2016). Thallus is gelatinous or soft cartilagenous like cartilage (Trono 1997; Prasetyowati et al 2008). In nature, *K. alvarezii* algae generally have two colors, namely brown and green (Calumpong & Meñez 1997). *K. alvarezii* cultivated in the Philippines and Indonesia was also found to be green and brown (Indriatmoko et al 2015; Pereira 2016). The habitat of *K. alvarezii* is in the sublittoral zone in the area of coral reef flats and lives on sandy to rocky substrates where water flow in these areas is low to moderate (Qin 2018).

The chemical composition of *K. alvarezii* is carbohydrate 65.20%, protein 3.40%, fat 1.10% and ash 11.57% (Khalil et al 2018). According to Rajasulochana et al (2012), *K. alvarezii* has antioxidant potential which comes from the content of vitamin C, vitamin E and metals (selenium and magnesium). Furthermore, according to Prabha et al (2013), *K. alvarezii* has antibacterial activity because it has several compounds including alkaloids, flavonoids, terpenoids, tannins and saponins.

1

K. alvarezii cultivation has the potential to be developed because it contains beneficial chemical compounds especially pigments. According to Thirumaran et al (2009), *K. alvarezii* contains carotenoid and chlorophyll pigments. Its main pigment contents are carotene, xanthophyll and xanthophyll derivatives as well as chlorophyll a and chlorophyll derivatives. According to de Fretes et al (2012), the dominant pigments in red algae are chlorophyll a, chlorophyll d, zeaxanthin, lycopene, cryptoxanthine, α -carotene, β -carotene, lutein, and picobillin.

The development of macroalgae in aquaculture activities is not only influenced by the cultivation technique and water quality, but also by photosynthetic pigment, namely chlorophyll-a. Chlorophyll-a is important for the survival of algae or for competing with other organisms in a certain habitat (Ming-Li et al 2010). Meanwhile, carotenoid pigments contained in *K. alvarezii*, have a role as accessory pigments in the photosynthesis process.

According to Lila (2004), each type of pigment has various health benefits. Mantiri & Kepel (1999) stated that carotenoid pigments functioned as natural dyes, could be used as immune materials and as raw materials for pharmaceutical, cosmetic, and foodstuff industries. B-carotene pigments have a physiological function as a precursor of vitamin A. According to Britton et al (2008), β -carotene pigments are useful for the human body and physiological activities especially eye, protecting skin surface tissues from stinging sunlight, as antioxidants and anticancer.

In North Sulawesi, the red algae *K. alvarezii* is abundant because it has been widely cultivated however this alga also was found growing naturally on the Minahasa peninsula which is only a few miles from Nain Island (Kepel et al 2019, 2020). However, there is no research conducted yet concerning pigments based on growth. Therefore, this study aimed to analyze the concentration of carotenoid and chlorophyll pigments in *K. alvarezii* algae cultivated on Nain Island.

Material and Method

Culture of *K. alvarezii*. In general, this type of algae is widely cultivated in the coastal waters of Nain Island, which was used as the research location (Figure 1). Nain Island has a huge coral reef zone that provides a supply of nutrients needed by these algae to grow. According to Trono (1997), coral reef areas were good places for algal growth. This algae cultivation was carried out in March - April 2020 in hot and rainy condition interspersed. The culture was conducted under controlled conditions aiming to measure the the weight gain of the thallus.




Figure 1. Nain Island of North Sulawesi, Indonesia.

K. alvarezii was cultured using the floating method as described by Qin (2018). The culture site had a water depth between 1.5 and 2.0 m during the lowest tide. The culture was adjusted in accordance to the tidal conditions in the coastal waters of Nain Island. A total of 17 thalli were tied to a rope with an initial weight of 100 g and the distance between the thallus was 25 cm. At the end of each rope a ballast was placed (Figure 2).

The culture was carried out for 9 weeks. Measurement of weight gain and collection of algal thalli for pigment analysis were carried out every two weeks.




Figure 2. Culture technique of *Kappaphycus alvarezii*.

Pigment extraction. Algal thalli were taken every two weeks, namely week 3, 5, 7 and 9, each with three replications. Furthermore, the carotenoid and chlorophyll pigments were extracted to determine the pigment concentration during the culture period. The initial extraction was Total Pigment Extract (EPT) from each thallus according to the culture time. Each thallus weighing as much as 50 g was crushed with acetone PA and added with Petroleum ether PA, forming EPT. The total pigment extract in petroleum ether was continued to an extraction step based on different polarity with the addition of a PA methanol solution, so that two layers were formed, the upper layer (PE) and the lower layer (Met²²

A. The top part of the pigment was added with 15 mL of KOH in methanol, so that two layers were formed namely: (1) The yellow top layer was the carotene pigment in Petroleum ether and (2) The green bottom layer was the chlorophyll a in methanol pigment.

B. The lower part of the pigment was added with diethyl ether and KOH in methanol, then it was divided into two layers, namely (3) the yellow upper layer was the xanthophyll pigment in diethyl ether and (4) the green bottom layer was the chlorophyll pigment in methanol.¹²

The results of each extract were then analysed using a UV-VIS GBC CINTRA 10 spectrophotometer with a wave length range of 380-550 nm for carotenoids (Britton et al 1995) and absorption spectrophotometer with a wave length of 350-700 nm for chlorophyll (Harborne 1987). The maximum absorption peak of the spectrophotometer formed was used to obtain the concentration of carotenoid pigments (Britton et al 1995) and chlorophyll pigments based on the mathematical formula of Jeffrey & Humprey (1975) and Harborne (1987).

Analysis of growth data. The formula to determine the growth rate was:

$$RGR = \frac{\ln W_t - \ln W_o}{t}$$

Where:
RGR = relative growth rate
W_o = initial weight of the thallus
W_t = final weight of thallus
t = time

Results and Discussion

Macroalgae growth. The cultivated *K. alvarezii* had a brownish colour as shown in Figure 3. According to Ahda et al (2005), the colour diversity of *K. alvarezii* is caused by environmental factors and is a chromatic adaptation process, namely the adjustment of the proportion of pigments with various lighting qualities.



Figure 3. Macroalgae *Kappaphycus alvarezii* with greenish brown color.

The growth of *K. alvarezii* in the coastal waters of Nain Island within two weeks showed an increase in weight gain. For the 17 thalli, the initial weight of each thallus was 100 g at week 1. At week 3, it was found that the mean weight of thallus was 135 g. Furthermore, at week 5, the average weight was 186 g. At the 7th week, the average weight was 260 g. At the 9th week, the average weight was 355 g. The growth of macroalgae is shown in Figure 4A.

Every two weeks the thallus *K. alvarezii* experienced an increase in weight gain, but the relative weight gain rate showed a decrease in the 3rd period (Figure 4B). The relative weight gain rate showed an increase of 2.26% day⁻¹ during the culture. The relative weight gain rate increased from the 1st period (between the 1st and 3rd week) with 2.14% day⁻¹, from the 2nd period (between the 3rd and 5th week) as much as 2.28% day⁻¹ and continued to increase to 2.39% day⁻¹ in the 3rd period (between the 5th and 7th week). Furthermore, it decreased in the 4th period (between the 7th and 9th week) to be 2.20% day⁻¹. In the fourth period, although the relatively weight gain rate decreased, the absolute weight gain of the thallus continued.

According to Salisbury & Ross (1995), growth is the increase in the size of an individual in terms of volume, weight and number of cells. Growth occurs due to cell division of the organism's body and the results of division will increase in volume or size, so that the stem cell will experience aging. Kasim et al (2019) reported a weight gain of *K. alvarezii* with 50 g of initial weight (Wo) to 262 g during 50 days of cultivation. In that study, the water quality for chemical concentrations including ammonia, nitrate + nitrite and phosphate were in normal conditions.

The rate of relative weight gain of *K. alvarezii* in coastal waters of Nain Island supposed to be influenced by fluctuations in light intensity during cultivation. This was indicated by an increase in temperature in the first three weeks and decreased in week 4. The pattern of relative growth rate of algae was similar to the pattern of temperature fluctuation during cultivation, where a decrease in temperature occurred at week 4 was in line with a decrease in the relative weight gain of the algae (Figure 4C). Jamal et al (2011) stated that the complex interaction between light radiation, temperature, nutrients and current strength at the cultivation site played an important role for the growth of *Kappaphycus*. According to Hurd et al (2014), changes in sea water temperature were directly related to the amount of light reaching the sea. Qin (2018) stated that optimal growth of macroalgae required sunlight in shallow waters, ranging from 30-50 cm. In this research activity, algae culture was carried out at the optimal range of sunlight. Furthermore, Irfan et al (2020) reported that the growth of *K. alvarezii* depended on ecological factors including temperature and brightness suitable for its cultivation activities. Ecological impacts on the sustainability of environmental resources and ecosystems provide benefits for sustainable macroalgae culture activities.

The temperature at the coastal waters of Nain Island during the research fluctuated from 26 to 30°C. Temperature has an important role in the metabolic process of *K. alvarezii*. Organisms are able to perform optimal metabolism at a proper water temperature. According to Lüning (1990), the temperature of water is very specific in influencing the growth of macroalgae. High water temperature, which is more than 36°C, causes mortality to macroalgae. This can affect the photosynthesis process, enzyme damage, and membrane instability. At the low water temperature, which is below 20°C, the protein and fat membranes will be damaged and form crystals. This is very influential for macroalgae life. According to Barsanti & Gualtieri (2006), the most important

parameters that regulate macroalgae growth are the quantity and quality of nutrients, light, pH, turbulence, salinity, and temperature as well as the photosynthesis process. The algae will be able to synthesize all the biochemical compounds essential for growth.




Figure 4. (A) Absolute growth of *Kappaphycus alvarezii*, (B) Relative growth of *K. alvarezii* and (C) Temperature (°C) and salinity (‰) ranges during the research.

Temperature is an important physical factor. Its changes tend to influence various chemical processes that occur simultaneously in plants and animal tissues and can affect the biota as a whole. Ain et al (2014) reported that water temperature can affect several physiological functions of macroalgae such as photosynthesis, respiration, metabolism, growth and reproduction. According to Poncomulyo et al (2006), sea water temperature is affected by sunlight, depth, currents and tides. According to Parenrengi et al (2006) and Setiyanto et al (2008), the water temperature range for *K. alvarezii* growth is 27-30°C.

The salinity in the coastal waters of Nain Island was 33.5-35‰ and the optimum salinity in the present study occurred at the time when the maximum growth was achieved in the 7th week, namely 35‰ (Figure 4C). This indicated that the salinity conditions of the coastal waters of Nain Island could support the growth of *K. alvarezii*

during the culture. According to Choi et al (2010), macroalgae will experience slow growth, if the salinity is too low (less than 15 ppt) or too high (more than 35 ppt) from the salinity range that corresponds to its life requirements for a certain period of time. Qin (2018) stated that the main factor determining the success or failure is selecting the appropriate site. Occasionally, those far from fresh water sources such as small river, etc., have been proven to be good sites. According to Kasim et al (2019), the most prominent physical factor affecting the growth of *K. alvarezii* is salinity because salinity greatly affects the macroalgae metabolism.

Yuliana et al (2015) reported that salinity affected the distribution of macroalgae in the sea. *K. alvarezii* is stenohaline, which is susceptible to wide salinity fluctuations. Most macroalgae have a low tolerance to changes in salinity. Sutresno & Prihastanti (2003) explained that optimum salinity could promote macroalgae to grow optimally, because the functional balance of the cell membrane could be maintained, especially in osmotic regulation in macroalgae and environmental fluids. This balance would facilitate the absorption of nutrients to support photosynthesis, so that macroalgae growth could be optimal.

Pigment concentration

Chlorophyll. The results showed that the pigments obtained were chlorophyll a and chlorophyll d. This is consistent with the statement of Harborne (1987) and Hurd et al (2014), that red algae contains two types of pigments, namely chlorophyll a and chlorophyll d. Figure 5 showed that during the research at weeks 3, 5, 7 and 9, the concentration of chlorophyll a and chlorophyll d recorded an increase. The concentration of chlorophyll d was higher than chlorophyll a. The lowest concentration of chlorophyll a was observed at week 3 (6.14 $\mu\text{g g}^{-1}$). At week 5, the concentration of chlorophyll a increased to 7.38 $\mu\text{g g}^{-1}$ and at week 7 increased to 10.02 $\mu\text{g g}^{-1}$. At the end of research at week 9, the concentration of chlorophyll a continued to rise until 18.32 $\mu\text{g g}^{-1}$. The recorded concentration of chlorophyll d at week 3 was 16.36 $\mu\text{g g}^{-1}$, and slightly increased at week 5 and 7 to be 16.41 $\mu\text{g g}^{-1}$ and 16.87 $\mu\text{g g}^{-1}$, respectively. At week 9, the concentration increased up to 21.42 $\mu\text{g g}^{-1}$ (Figure 5).




Figure 5. Concentration of chlorophyll a and chlorophyll d of *Kappaphycus alvarezii* during the two weeks of culture.

The increasing pattern of chlorophyll a and chlorophyll d concentrations were consistent with the weight gain pattern of the *K. alvarezii* thalli. This was due to the weight of thalli which continued to increase during the cultivation. The increase in weight gain of thalli resulted in the continue growing of cells followed by the development of chloroplast cells through photosynthetic reactions. Thus, the chlorophyll a and d concentrations as pigments contained in *K. alvarezii* would also develop.

During the cultivation of *K. alvarezii*, chlorophyll a and accessories pigments plays a role in absorbing sunlight for the photosynthesis process. Chlorophyll a is the main pigment in red algae (Jupin & Lamant 1999). Hurd et al (2014) stated that the chlorophyll d concentration was found to be higher than the chlorophyll a in cyanobacterium (*Acaryochloris marina*) in waters. According to Lüning (1990), the

dominant pigments found in red algae are chlorophyll a, chlorophyll d, zeaxanthin, cryptoxanthine lycopene, α -carotene, β -carotene, lutein, and picobilin. Chlorophyll d as an accessory pigment can support the role of chlorophyll a as the main pigment in red algae. According to Barsanti & Gualtieri (2006), cells that were accustomed to high radiation levels generally had relatively high concentrations of carotenoid pigments compared to chlorophyll a. Carotenoid pigments such as β -carotene and zeaxanthin pigments do not transfer excitation energy to the reaction center and consequently act as filter cells from excessive light. According to Indriatmoko et al (2015), the changes of chlorophyll and carotenoid concentrations of *K. alvarezii* specifically the one cultivated at the sea in shallow and clear water was very interesting to be studied.

Carotenoids. The content of carotenoid pigments in *K. alvarezii* consists of β -carotene and xanthophyll pigments, namely zeaxanthin, violaxanthin, cryptoxanthine, and lutein (Fretes et al 2011; Andersson et al 2006). Pigments in diethyl ether were extracted from different polarity of the thallus. The pigment content of *K. alvarezii* in the present research was identified as β -carotene.

The results of this study showed that the concentration of β -carotene pigment taken during the research with two weeks interval tended to increase until it reached its maximum point at week 7. Subsequently there was a decrease in concentration until week 9 (Figure 6).




Figure 6. Pigment concentration and carotene amount of *Kappaphycus alvarezii* during the two weeks of culture.

The average concentration of β -carotene pigment extracted based on the separation of different solvents at week 3 was $17.18 \mu\text{g g}^{-1}$ of the dry residual weight. At week 3, the carotene pigment concentration was the highest compared to the pigment concentration of xanthophyll, chlorophyll a and chlorophyll d. At week 5, the carotene pigment concentration increased to $17.63 \mu\text{g g}^{-1}$. It was supposed that at weeks 3 and 5, the carotene pigment played a major role in the photosynthesis process. At week 7, the carotene pigment concentration rose to $27.50 \mu\text{g g}^{-1}$. At the 9th week, the carotene pigment concentration decreased to $7.78 \mu\text{g g}^{-1}$. The concentration pattern of carotene pigments in *K. alvarezii* followed the relative growth rate pattern of the algae.

The average quantity of β -carotene pigment followed the pigment concentration pattern and the relative growth rate pattern of the algae. The average carotene amount at week 3 was $31.53 \mu\text{g}$ from the wet weight of *K. alvarezii*. At week 5, it increased to $39.43 \mu\text{g}$ and at week 7, it increased to $50.81 \mu\text{g}$. At the 9th week there was a decrease to $17.00 \mu\text{g}$. Thus, the average carotene amount would increase along with the increase of growth rate of these algae to carry out the photosynthesis process.

Similar pigment concentration and carotene amount patterns were found by Paransa et al (2002), where the red algae *Kappaphycus striatum* was cultured on the surface of coastal waters of Nain Island for two months. Pigment concentration and carotene amount during the culture increased from week 3 to week 7. At the 7th week there was a decrease until the 9th week. There was a similarity in the pigment concentration and carotene amount even though the culture was carried out in different

months. It was suspected that this similarity occurred because algae came from the same genus, namely *Kappaphycus*, the same rearing location but with different months and years.

The average xanthophyll pigment concentration at week 3 was 13.74 $\mu\text{g g}^{-1}$, and increased to 16.64 $\mu\text{g g}^{-1}$ at week 5. At week 7 increased to 35.55 $\mu\text{g g}^{-1}$, but at week 9 decreased to 14.78 $\mu\text{g g}^{-1}$. The average xanthophyll amount at week 3 was 25.07 μg and increased at week 5 and 7 to 36.43 μg and 65.95 μg , respectively. However, at week 9 it decreased to 35.18 μg (Figure 7).




Figure 7. Concentration and amount of xanthophyll pigment of *Kappaphycus alvarezii* extract during the two weeks of culture.

The average concentration and amount of xanthophyll pigments obtained were the same, where there was an increase both in pigment concentration and xanthophyll amount at week 5 and 7. The highest average quantity was obtained at week 7, while at week 9 the average quantity and quality of xanthophyll pigments decreased due to the age of algae.

Conclusions. The concentration of chlorophyll a and d pigments increased as the weight gain of *K. alvarezii* increased. The increase in pigment concentration and carotene and xanthophyll (carotenoid) amount were in line with the growth rate pattern of *K. alvarezii*.

Acknowledgements. The authors wish to thank Sam Ratulangi University for supporting the financial part of this research through Dana Riset Dasar Unggulan Unsrat Tahun 2020.

References

- Ahda A., Surono A., Imam A., Batubara I., Ismanadji I., Suitha M., Yunaidar R., Setiawan, Kurnia N., Danakusumah E., Sulistijo, Zalnika A., Basmal J., Effendi I., Runtuboy N., 2005 [Profile of Indonesian seaweed]. Jakarta (ID): Direktorat Jenderal Perikanan Budidaya Departemen Kelautan dan Perikanan. [In Indonesian].
- Ain N., Ruswahyuni, Widyarini N., 2014 [The correlation of seaweed density with a different substrate in the waters of Bandengan, Jepara beach]. Diponegoro Journal of Maquares 3(1):99-107. [In Indonesian].
- Andersson M., Schubert H., Pedersén M., Snoeijs P., 2006 Different patterns of carotenoid composition and photosynthesis acclimation in two tropical red algae. *Marine Biology* 149:653-665.
- Barsanti L., Gualtieri P., 2006 *Algae anatomy, biochemistry and biotechnology*. Taylor & Francis Group-CRC Press, 320 p.
- Britton G., Jansen S. L., Pfander H., 1995 *Carotenoids*. Volume 1B: Spectroscopy. Basel, Switzerland, 347 p.
- Britton G., Jansen S. L., Pfander H., 2008 *Carotenoids*. Volume 4: Natural functions. Basel, Switzerland.

- Calumpang P., Meñez H., 1997 Field guide to common mangroves, seagrass and algae of the Philippines. Book Mark, inc. Makati City, Philipines.
- Choi T. S., Kang E. J., Kim J. H., Kim K. Y., 2010 Effect of salinity on growth and nutrient uptake of *Ulva pertusa* (Chlorophyta) from an eelgrass bed. *Algae* 25(1):17-25.
- de Fretes H., Susanto A. B., Limantara L., Prasetyo B., Heriyanto, Brotosudarmo T. H. P., 2011 Composition and content of pigment, photostability and thermostability studies of crudepigment extracts from red, brown, and green varieties of red algae *Kappaphycus alvarezii* (Doty) Doty. Seminar ICONS, Universitas Ma Chung, Malang.
- de Fretes H., Susanto A. B., Budhi P., Heriyanto, Tatas H. P., Brotosudarmo, Limantara L., 2012 [Estimation of degradation product degradation of red algae pigment crude extract, *Kappaphycus alvarezii* (Doty) Doty, variants of red, brown and green: examine the absorption spectrum differences]. *Jurnal Ilmu Kelautan* 17(1):31-38. [In Indonesian].
- Harborne J. B., 1987 Phytochemical methods, guide modern ways to analyse plants. Penerbit ITB, Bandung, Indonesia.
- Hurd C. L., Harrison P. J., Bischof K., Lobban C. S., 2014 Seaweed ecology and physiology. Cambridge Univerity Press, 552 p.
- Indriatmoko, Heriyanto, Limantara L., Brotosudarmo T. H. P., 2015 Composition of photosynthetic pigments in a red alga *Kappaphycus alvarezii* cultivated in different depths. *Jurnal Procedia Chemistry* 14:193-201.
- Irfan M., Abdullah N., Yuliana, Samadan G. M., Subur R., 2020 Sustainability status of seaweed *Kappaphycus alvarezii* cultivation in Posi-Posi waters, South Kayoa Sub-district, Regency of South Halmahera, Province of North Mollucas, Indonesia. *International Journal of Advanced Science and Technology* 29(8):806-817.
- Jamal E., Loupatty, J. W., Pattinasarany M. M., 2011 Seasonal water quality challenges of *Kappaphycus* sp. seaweed culture production in Maluku waters. *Prosiding Seminar Internasional Forum Pimpinan Pascasarjana Se-Indonesia*. Ambon. pp. 81-87.
- Jeffrey S. T., Humphrey G. F., 1975 New spectrophotometric equations for determining chlorophylls a, b, c1 and c2 in higher plants, algae and natural phytoplankton. *Biochemie und Physiologie der Pflanzen* 167(2):191-194.
- Jupin H., Lamant A., 1999 [The photosynthesis]. Dunod, Paris, 268 p. [In French].
- Kasim M., Mustafa A., Ishak E., Ibrahim M. N., Irawati N., Wanurgayah, Munir T., Muzuni, Jalil W. A., 2019 Environmental status of *Kappaphycus alvarezii* cultivation area following temporary eutrophication. *Jurnal AACL Bioflux* 12(4):1102-1113.
- Kepel R. C., Lumingas L. J. L., Tombokan J. L., Mantiri D. M. H., 2019 Biodiversity and community structure of seaweeds in Minahasa Peninsula, North Sulawesi, Indonesia, North Sulawesi, Indonesia. *AACL Bioflux* 12(3):880-892.
- Kepel R. C., Lumingas L. J. L., Tombokan J. L., Mantiri D. M. H., 2020 Community structure of seaweeds in dry season in Minahasa Peninsula, North Sulawesi, Indonesia. *AACL Bioflux* 13(1):393-402.
- Khalil H. A., Yap S. W., Tye Y. Y., Tahir P. M., Rizal S., Fazita M. N., 2018 Effects of corn starch and *Kappaphycus alvarezii* seaweed blend concentration on the optical, mechanical, and watervapor barrier properties of composite films. *BioResources* 13(1):1157-1173.
- Kreckhoff R. L., Ngangi E. L., Undap S. L., Kusen D. J., 2019 Crude extracts of *Kappaphycus alvarezii* algae cultivated in several seaweed production centers in North Sulawesi, Indonesia as immunostimulant. *AACL Bioflux* 12(2):678-686.
- Lila M. A., 2004 Plant pigments and human health. In: *Plant pigments and their manipulation*. Davis K. M. (ed), CRC Press, London, pp. 248-274.
- Lüning K., 1990 *Seaweeds: their environment, biogeography, and ecophysiology*. A Wiley Interscience.
- Mantiri D. H. M., Kepel B., 1999 [Several roles of carotenoid pigment]. *Jurnal Fakultas Perikanan dan Ilmu Kelautan* 1(3). [In Indonesian].
- Ming-Li T., Chu W. L., Phang S. M., 2010 Effect of temperature change on physiology and biochemistry of algae: a review. *Malaysian Journal of Science* 29(2):82-97.

- Paransa D. J., Kepel R. C., Yusa I. K. S. A., 2002 [The growth of the red algae *Kappaphycus striatum* (Schmitz) Doty from the coastal waters of Nain Island]. *Jurnal Perikanan dan Ilmu Kelautan* 1(1):8-14. [In Indonesian].
- Parenrengi A., Sulaeman, Suryati E., Tenriulo A., 2006 [A genetic characterization of the seaweed *Kappaphycus alvarezii* cultivated in South Sulawesi]. *Jurnal Riset Akuakultur* 1(1):1-11. [In Indonesian].
- Pereira L., 2016 *Edible seaweed of the world*. Taylor & Francis Group, 463 p.
- Poncomulyo T., Maryani H., Kristiani L., 2006 [Seaweed culture and processing]. Agromedia. Jakarta, Indonesia, Pp. 14-15. [In Indonesian].
- Prabha V., Prakash D. J., Sudha P. N. 2013 Analysis of bioactive compounds and antimicrobial activity of marine algae *Kappaphycus alvarezii* using three solvent extract. *International Journal of Pharmaceutical Science and Research* 4(1):306-310.
- Prasetyowati, Jasmine A. C., Agustawan D., 2008 [The manufacture of marine flour (*Eucheuma cottonii*) based on the difference in precipitation]. *Jurnal Teknik Kimia* 2(15):27-33. [In Indonesian].
- Qin Y. M., 2018 *Bioactive seaweeds for food applications. Natural ingredients for healthy diets. Part I, Seaweed Bioresources*. Elsevier Academic Press.
- Rajasulochana P., Krishnamoorthy P., Dhamotharan R., 2012 Potential application of *Kappaphycus alvarezii* in agricultural and pharmaceutical industry. *Journal of Chemical and Pharmaceutical Research* 4(1):33-37.
- Salisbury F. B., Ross C. W., 1995 [Plant physiology]. Vol. 3, ITB. Bandung, 343 p. [In Indonesian].
- Setiyanto D., Efendi I., Between K. J., 2008 Growth of *Kappaphycus alvarezii* var. *maumare*, var. *Sacol* and *Eucheuma cottonii* in Musi Buleleng waters. *Journal of Marine Science* 13(3):171-176.
- Sutresno, Prihastanti E., 2003 The effect of salinity on the growth of red algae *Gracilaria verrucosa*. *Buletin Anatomi dan Fisiologi* 11(1):14-20.
- Thirumaran G., Manivannan K. G., Karthikai D., Anantharaman P., Balasbramanian T., 2009 Photosynthetic pigments of different colour strain of the cultured seaweed *Kappaphycus alvarezii* (Doty) Doty ex. P. Silva in Vellar estuary. *Journal of Plant Sciences* 2(3):150-153.
- Trono G. C., 1997 *Field guide: atlas of the seaweed resources of the Philippines*. National Bookstore, Inc., Manila, The Philippines, 306 p.
- Yuliana A., Rejeki S., Widowati L. K., 2015 The effect of different salinity on the growth of latoh seaweed (*Caulerpa lentillifera*) in the Jepara beach, Wikayah (LPWP) Laboratory. *Journal of Aquaculture Management and Technology* 4(4):61-66.

Received: 04 September 2020. Accepted: 12 October 2020. Published online: 19 October 2020.

Authors:

Darus Saadah Johanis Paransa, Sam Ratulangi University, Faculty of Fisheries and Marine Science Unsrat,

Indonesia, North Sulawesi, Manado 95115, Jln. Kampus Unsrat Bahu, e-mail: darusparansa@unsrat.ac.id

Desy Maria Helena Mantiri, Sam Ratulangi University, Faculty of Fisheries and Marine Science Unsrat, Indonesia,

North Sulawesi, Manado 95115, Jln. Kampus Unsrat Bahu, e-mail: dmh_mantiri@unsrat.ac.id

Rene Charles Kepel, Sam Ratulangi University, Faculty of Fisheries and Marine Science Unsrat, Indonesia, North

Sulawesi, Manado 95115, Jln. Kampus Unsrat Bahu, e-mail: renecharleskepel65@gmail.com

Antonius Rumengan, Sam Ratulangi University, Faculty of Fisheries and Marine Science Unsrat, Indonesia,

North Sulawesi, Manado 95115, Jln. Kampus Unsrat Bahu, e-mail: antonius_rumengan@unsrat.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Paransa D. J., Mantiri D. M. H., Kepel R. C., Rumengan A., 2020 Pigment concentration of red algae, *Kappaphycus alvarezii* (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia. *AAFL Bioflux* 13(5):2788-2797.

Pigment concentration of red algae, *Kappaphycus alvarezii* (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia

ORIGINALITY REPORT

12%

SIMILARITY INDEX

8%

INTERNET SOURCES

8%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to Universitas Jenderal Soedirman
Student Paper 2%
- 2** Pang-Hung Yiu, Jian See, Amartalingam Rajan, Choon-Fah J. Bong. "Boric Acid Levels in Fresh Noodles and Fish Ball", American Journal of Agricultural and Biological Sciences, 2008
Publication 1%
- 3** rjoas.com
Internet Source 1%
- 4** Barsanti, . "Algal Culturing", Algae Anatomy Biochemistry and Biotechnology, 2005.
Publication 1%
- 5** thuvien.ntu.edu.vn
Internet Source 1%
- 6** ijsrp.org
Internet Source 1%
- 7** Subir Kumar Mandal. "Invasion potential of *Kappaphycus alvarezii* on corals at Kurusadai 1%

Island, Gulf of Mannar, India", ALGAE,

12/01/2010

Publication

8

www.gjesm.net

Internet Source

<1 %

9

Arini Fadhla Wahyuningtyas, Prayogo, Annur Ahadi Abdillah, M. Nur Ghoyatul Amin, Mochammad Amin Alamsjah. " The Effectivity of Plantation Depth on Seaweed sp. Growth Using Longline Method ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

10

"Sustainable Global Resources Of Seaweeds Volume 1", Springer Science and Business Media LLC, 2022

Publication

<1 %

11

Submitted to Universitas Kristen Satya Wacana

Student Paper

<1 %

12

f1000research.com

Internet Source

<1 %

13

journalppw.com

Internet Source

<1 %

14

repo.unsrat.ac.id

Internet Source

<1 %

15

www.scilit.net

Internet Source

<1 %

16 assets.researchsquare.com <1 %
Internet Source

17 Bastian Niedrée, Anne E. Berns, Harry Vereecken, Peter Burauel. "Do Chernobyl-like contaminations with ¹³⁷Cs and ⁹⁰Sr affect the microbial community, the fungal biomass and the composition of soil organic matter in soil?", *Journal of Environmental Radioactivity*, 2013 <1 %
Publication

18 Eighteenth International Seaweed Symposium, 2007. <1 %
Publication

19 "Sixteenth International Seaweed Symposium", Springer Nature, 1999 <1 %
Publication

20 C. Periyasamy, P. V. Subba Rao, P. Anantharaman. "Spatial and temporal variation of biochemical contents in relation to carrageenan yield of cultivated *Kappaphycus alvarezii* (Doty) Doty at different locations of Palk Bay waters, Tamil Nadu, southeast coast of India", *Journal of Applied Phycology*, 2018 <1 %
Publication

21 Iskandar, Akmal, Rajuddin Syam, and Dody Dh Trijuno. "Content of Carrageenan, Chlorophyll <1 %

a and Carotenoid of *Kappaphycus alvarezii*
Cultivated in Different Seawater Depth
Laikang Village, District of Mangarabombang,
Takalar Regency", *Journal of Applied
Biotechnology*, 2013.

Publication

22

e-journal.uajy.ac.id

Internet Source

<1 %

23

id.scribd.com

Internet Source

<1 %

24

repository.ung.ac.id

Internet Source

<1 %

25

www.pesca.sp.gov.br

Internet Source

<1 %

26

Rene Charles Kepel, Lawrence J. L. Lumingas,
John L. Tombokan, Desy M. H. Mantiri.

"Community structure of macroalgae in
Tanjung Merah, rap-rap and Totok Bay, North
Sulawesi, Indonesia", AIP Publishing, 2023

Publication

27

S.C. Wilson, A.A. Meharg. "Investigation of
organic xenobiotic transfers, partitioning and
processing in air-soil-plant systems using a
microcosm apparatus. Part II: comparing the
fate of chlorobenzenes in grass planted soil",
Chemosphere, 2003

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On