

**Selected as Book Chapter "Pigment concentration of red algae, Kappaphycus alvarezii (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia"**

1 pesan

Editor BPI IN <editor@bpinternational.org>  
Kepada: darusparansa@unsrat.ac.id

13 Mei 2022 pukul 16.4'

To  
**Darus S. J. Paransa**  
Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado, North Sulawesi, Indonesia  
Email: darusparansa@unsrat.ac.id

**Subject: Selected as Book Chapter "Pigment concentration of red algae, Kappaphycus alvarezii (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia"**

Dear Dr. Darus S. J. Paransa,

We are glad that you are reading this mail. We hereby heartily congratulate you for the publication of your following paper. We appreciate your hard work and sincerity behind the publication of this excellent paper.

**"Pigment concentration of red algae, Kappaphycus alvarezii (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia"**

in

AAFL Bioflux, 2020, Volume 13, Issue 5

You are already aware that publication of research work in a peer-reviewed international journal has important value. At the same time publication of research work as a Book chapter in an Edited Book also has a significant importance in career. A research paper which has already been published in a journal can be published as a Book Chapter in an expanded form. It is completely legal and acceptable procedure in the academic world.

We want to share good news with you. We have published more than 82 books in last three months (<https://www.bpinternational.org/>). We are overwhelmed with pleasure for this tremendous response from researchers like you. We are going to publish more than 5 volumes in next two months in the following area. With the same research work you can get double benefit and it will enrich your resume.

*We're happy to inform that your paper has been selected to be included as one of the potential book chapters of the following book. You're also welcome to publish your other research papers, which have been published in other journals. If you have any fresh manuscript, you can submit for review and inclusion as book chapter.*

**Book name: Emerging Challenges in Environment and Earth Science**

Details of charges are available below. Presently 80% discount on the publication charge has been announced for this month.

<https://www.bpinternational.org/wp-content/uploads/2019/04/9.Charges.pdf>

**Features:**

1. Professional review, editing and plagiarism checking.
2. Professional cover-page design and typesetting.
3. Specialized English editing and proofreading
4. Digital Object Identifier (DOI) allotment.
5. ISBN number (online and print version)
6. Online publication of the book and printing in the press.
7. Normally Books will be closed access to increase the possibility of sale and authors will retain the copyright of the book chapter/books. Special request of open access books can also be considered in case of complete book publication.

**Complimentary free services**

1. Social media promotion of the book.
2. Online Press release of the book for better promotion.
3. Advertisement for the book in different News portals

5. Search engine promotion, back-linking and optimization.
6. Promotional video preparation and sharing in YouTube, etc.

**Copyright:** You have published your paper with CC-BY license in the journal. So you hold the copyright of your paper. Therefore, CC-BY license gives you the permission to republish your journal article as book chapter. If you have not published your paper in Open Access journal and publisher posses the copyright, then just ask them permission to republish it as book chapter.

If you are interested please mail us. We shall be eagerly waiting to hear from you.

**Kindly please don't reply to this mail id as this mail id is not monitored.**

**"Please reply us to [editor.2@bookpi.org](mailto:editor.2@bookpi.org) or [submission@bookpi.info](mailto:submission@bookpi.info) and send us your manuscript as an email attachment."**

**Note: Kindly note if you have already responded to this mail earlier, please ignore this mail.**

With Regards

Ms. Sumona Bag

B P International

Email: [editor.2@bookpi.org](mailto:editor.2@bookpi.org) or [submission@bookpi.info](mailto:submission@bookpi.info)

Reg. Offices:

India: B P International, Guest House Road, Street no - 1/6, Hooghly, West Bengal, India, Corp. Firm Registration Number: L77527,

Tele: +91 7439016438

UK: B P International, 27 Old Gloucester Street London WC1N 3AX, UK, Fax: +44 20-3031-1429



**Selected as Book Chapter "Pigment concentration of red algae, Kappaphycus alvarezii (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia"**

1 pesan

Editor BPI <editor@bpipromotion.org>  
Kepada: darusparansa@unsrat.ac.id

27 Mei 2023 pukul 18.00

To

**Darus S. J. Paransa**

**Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado, North Sulawesi, Indonesia**

**Email: darusparansa@unsrat.ac.id**

**Subject: Selected as Book Chapter "Pigment concentration of red algae, Kappaphycus alvarezii (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia"**

Dear Dr. Darus S. J. Paransa,

We are glad that you are reading this mail. We hereby heartily congratulate you for the publication of your following paper. We appreciate your hard work and sincerity behind the publication of this excellent paper.

**"Pigment concentration of red algae, Kappaphycus alvarezii (Doty) Doty ex Silva during the cultivation in the coastal waters of Nain Island, North Sulawesi, Indonesia"**

in

**AAAL Bioflux, 2020, Volume 13, Issue 5**

You are already aware that publication of research work in a peer-reviewed international journal has important value. At the same time publication of research work as a Book chapter in an Edited Book also has a significant importance in career. A research paper which has already been published in a journal can be published as a Book Chapter in an expanded form. It is completely legal and acceptable procedure in the academic world.

We have published more than 1000 books in last two years (<https://www.bpipromotion.org/>). We are overwhelmed with pleasure for this tremendous response from researchers like you. We are going to publish more than 5 volumes in next two months in the following area. Therefore, with the same research paper you can get double benefit and it will enrich your resume.

*We're happy to inform that your paper has been selected to be included as one of the potential book chapters of the following book. You're also welcome to publish your other research papers, which have been published in other journals. If you have any fresh manuscript, you can submit for review and inclusion as book chapter.*

**Book name: Novel Perspectives of Geography, Environment and Earth Sciences**

**Features:**

1. Professional review, editing and plagiarism checking.
2. Professional cover-page design and typesetting.
3. Specialized English editing and proofreading
4. Digital Object Identifier (DOI) allotment.
5. ISBN number (online and print version)
6. Online publication of the book and printing in the press.
7. Normally Books will be closed access to increase the possibility of sale and authors will retain the copyright of the book chapter/ books. Special request of open access books can also be considered in case of complete book publication.

**Complimentary free services**

1. Social media promotion of the book.
2. Online Press release of the book for better promotion.
3. Advertisement for the book in different News portals.
4. Circulation of the book in different digital book stores.
5. Search engine promotion, back-linking and optimization.
6. Promotional video preparation and sharing in YouTube, etc.

**Copyright:** When you publish your paper with CC-BY license in an open access journal, you hold the copyright of your paper. Therefore, CC-BY license gives you the permission to republish your journal article as book chapter. If you have not published your paper in Open Access journal and publisher holds the copyright, then our experts will guide you to get the permission to republish it as book chapter.

If you are interested please mail us. **Kindly please don't reply to this mail id as this mail id is not monitored.**

**"Please reply us to editor.2@bookpi.org or submission@bookpi.info and send us your manuscript as an email attachment."**

**NOTE:** You are very politely requested to note that this invitation is sent to you as a preliminary proposal for your book / book chapter. Without your written consent no book / book chapter will be published.

Thank you for your cooperation and valuable time.

**Note: Kindly note if you have already responded to this mail earlier, please ignore this mail.**

With Regards

Ms. Sumona Bag

B P International

Reg. Offices:

India: B P International, Guest House Road, Street no - 1/6, Hooghly, West Bengal, India, Corp. Firm Registration Number: L77527,  
Tele: +91 7439016438

UK: B P International, 27 Old Gloucester Street London WC1N 3AX, UK, Fax: +44 20-3031-1429

Unsubscribe from this list!

If you'd like to unsubscribe and stop receiving these emails click here .

# Exploration study of the pigment types of stone crab (*Grapsus albolineatus*) as pharmaceutical and cosmetics Ingridients

*by* Darus Sa'adah Johanis Paransa 4

---

**Submission date:** 30-May-2023 08:22AM (UTC+0700)

**Submission ID:** 2104891184

**File name:** JURNAL\_INTERNASIONAL\_-\_2019\_-\_DarusSaadah\_J\_Paransa\_EEC-min.pdf (284.04K)

**Word count:** 3340

**Character count:** 18473



## Exploration study of the pigment types of stone crab (*Grapsus albolineatus*) as pharmaceutical and cosmetics Ingridients

Darus Saadah J. Paransa<sup>1\*</sup>, Desy M.H. Mantiri<sup>1</sup>, James J.H. Paulus<sup>1</sup>, Cysca Lumenta<sup>1</sup>, Medy ompie<sup>1</sup>, Suria Darwisito<sup>1</sup>, Carolus Paruntu<sup>1</sup>, Alfred Kase<sup>2</sup> and Anggreni D.N. Rupidara<sup>2</sup>

<sup>1</sup>Fisheries and Marine Science Study Program of Sam Ratulangi University, Manado, North Sulawesi, Indonesia

<sup>2</sup>Fisheries and Marine Science Study Program of Artha Christian University, Kupang, East Nusa Tenggara, Indonesia

(Received 14 March, 2019; Accepted 27 May, 2019)

### ABSTRACT

*Grapsus albolineatus* is a type of crustacean on the rocky coast of tropical regions like Indonesia. Some of the distribution areas of this crab species are the rocky coastline of, Tanawangko and Manggatasik Village of, Manado City, North Sulawesi. *Grapsus albolineatus* is known to have a high content of carotenoid pigment. Carotenoid pigments are useful for human needs, such as immune substances, and various industrial raw materials such as pharmaceutical, cosmetic and foodstuff industries. In this study, identification of the pigment type was performed on the carapace, gonad, blood, epidermal layer, and hepatopancreas using Thin Layer Chromatography method. The identification of pigments on the carapace was using Column Chromatography and High-Performance Liquid Chromatography (HPLC) methods. The results showed that *Grapsus albolineatus* was identified at the D molting stage. In the Thin Layer Chromatography analysis, there were found some pigment types\* namely  $\beta$ -Carotene, Echinenone, Canthaxanthin, Cryptoxanthin, Adonirubin, Astaxanthin and Astacen. Meanwhile, the separation by Column Chromatography method showed the results of pigments such as  $\beta$ -Carotene, Canthaxanthin, Adonirubin, and Fucoxanthin. The concentrations of pigments contained in the carapace, epidermal layer, hepatopancreas, gonad and blood were respectively 6.88 mg/g, 210.47 mg/g, 454.0 mg/g, 433.33 mg/g, and 153 mg/g. The pigment content of each organ of *G. albolineatus* crabs was 6.95 mg (the carapace), 7.26 mg (epidermal layer), 7.07 mg (hepatopancreas), 34.94 mg (gonad), and 5.86 mg (hemocyanin).

**Key words :** Pigment, Carotenoid, Crustacea, TLC, CC

### Introduction

One of the organisms that become the potential resource of North Sulawesi Province, Indonesia is crabs. The coastal morphology of North Sulawesi is dominated by mangrove coastal waters and rocky coastal waters which are the habitats for crabs to survive. Bliss and Mantel (1985) found that *Scylla*

*serrata* crabs survive by adapting to mangrove areas. Moreover, according to Majchacheep (1989), there are many types of *Grapsus albolineatus* found in rocky coastal waters. Both types of crabs contain carotenoid pigments in the external and internal organs. Carotenoid pigments in these two types of crabs can be used as pharmaceutical and cosmetic raw materials.

Corresponding author's email: [darusparansa@unsrat.ac.id](mailto:darusparansa@unsrat.ac.id)

Based on the results of studies by Paransa *et al.* (2000a) and Paransa *et al.* (2000b), in female and male *S. serrata* crab extracts, there were found same types of pigments, namely  $\beta$ -carotene, zeaxanthin ester, astaxanthin ester, and free phenocoxanthin and astaxanthin. Both female and male *S. serrata* crabs were obtained in the Coastal Waters of Likupang Village, North Minahasa Regency, North Sulawesi Province, Indonesia. Paransa *et al.* (2002) also discovered that in the total extract of *G. albolineatus* crabs captured in the coastal waters of Kalasey Beach, Minahasa Regency, North Sulawesi, Indonesia which is dominantly rocky, there were found some types of pigments, namely  $\beta$ -carotene, echinenone, canthaxanthin in the internal organs (hemocyanin, hepatopancreas, and epidermal layer) and some other types of pigments, namely free phenocoxanthin and astaxanthin, were found in the external organ of carapace. Furthermore, according to Paransa (2005) and Paransa *et al.* (2008), the type of major carotenoid pigment, namely astaxanthin was found separately in the extracted carapace, epidermal layer, hepatopancreas, hemocyanin, and gonad of male *G. albolineatus* crabs captured in the Coastal Waters of Malalayang Beach, Manado, North Sulawesi, Indonesia. The analysis of the carotenoid pigments of *Scylla serrata* and *G. albolineatus* crabs was done using Thin Layer Chromatography (TLC) analysis method with semipolar developer solutions.

Mantiri and Kepel (1999) stated that the carotenoid pigments as an ingredient for natural dyes can be used as immune substances and raw materials for pharmaceutical, cosmetic, and foodstuff industries. Additionally, the results of Naguib's (2000) study suggest that astaxanthin pigments have an antioxidant activity that is ten times stronger than  $\alpha$ -carotene, canthaxanthin, lutein and zeaxanthin pigments. Based on the results of several previous studies above, the carapaces of stone crabs (*G. albolineatus*) contain astaxanthin pigments which can be used as a source of natural colouring agents. Utilization of natural dyes from the extraction of coastal resources of North Sulawesi, Indonesia can support the industrial sector. This research was enhanced by utilizing purer materials. Thus, the analysis of carotenoid pigments was necessarily conducted through the chromatography column stage.

#### Research Method

The crab (*G. albolineatus*) samples were taken at

night during the high tide of the coastal waters of Tanawangko Village, Minahasa Regency, North Sulawesi, Indonesia. The organism species of the samples were then identified and then continued with the molting stage determination. The next stage was taking the internal and external organs, including Carapace (C), Epidermal Layer (EL), Hepatopancreas (H), Gonad (G), and Hemocyanin (D) to be extracted through the grinding process in acetone and petroleum ether solutions. The concentration and quantity of carotenoid pigments in the total pigment extract were analyzed according to Britton *et al.* (1995).

The types of pigments in Carapace (C), Epidermal Layer (EL), Hepatopancreas (H), Gonad (G), and Hemocyanin (D) were determined using Thin Layer Chromatography (TLC) and maximum UV-Vis spectrophotometric absorption analysis methods with a length wave ranging from 380 – 550 nm (Britton *et al.*, 2004). The TLC analysis was done using a semi-polar developer according to Mantiri (1997). The content of the total pigment extract of the carapace organ was analyzed using the separation of Chromatography Column (CC) and the maximum UV-Vis spectrophotometric absorption analysis method according to Britton *et al.* (1995). The CC analysis was undertaken using the dry method principle using a semipolar developer and silica gel powder as a stationary phase (Mantiri, 1997).

The separation of carotenoids using HPLC was achieved in a mixture of methanol: acetone: water (80: 5: 15) and 0.8 mL minute water rate. After 10 minutes, the water content is reduced linearly when the acetone increased into 20% at 50 minutes after injection. The composition of the gradient mixture is methanol: acetone (80:20).

## Results and Discussion

### Sample Identification

The sample identification was aimed to determine the species of the samples obtained from the sampling location. The carapace morphological form of the stone crabs taken from the location had blackish green stripe patterns and the anterior part between the eyes was divided into four blackish green lobes. The first pair of walking legs was the shortest with a line of setae that appeared on the ventral boundary of the walking leg which amounted to eight legs



(four pairs) with no pair of swimming legs. This type of crabs was characterized by a pair of purple claws (as seen in Figure 1). Based on Carpenter (1910), crabs living on the rocky shore are identified as *Grapsus albolineatus* species (Latreille in Milbert 1812) and mostly male. The uropod part of *G. albolineatus* crabs (Latreille in Milbert, 1812) used in this research was observed under a microscope with 10x magnification, and based on Skinner (1962), this belongs to the D moulting stage or also known to be at the premolt stage. In the premolt stage, a new rigid exoskeleton would be formed.

**Analysis of Carotenoid Pigment Type and Content**

The concentrations of pigments contained in the carapace, epidermal layer, hepatopancreas, gonad and blood were respectively 6.88 mg/g, 210.47 mg/

g, 454.0 mg/g, 433.33 mg/g, and 153 mg/g. The pigment content of each organ of *G. Albolineatus* crab was 6.95 mg (carapace), 7.26 mg (epidermal layer), 7.07 mg (hepatopancreas), 34.94 mg (gonad), and 5.86 mg (blood). Types of pigments contained in each organ of *G. albolineatus* crabs (Latreille in Milbert, 1812) can be seen in the table below.

The distribution of pigment types in each organ of *G. albolineatus* crabs (Latreille in Milbert, 1812) presented in the table above shows that the first fraction produced yellow color identified as a  $\beta$ -carotene pigment, contained in the gonad, hemocyanin, and hepatopancreas organs. The second fraction appeared to be orange-colored identified as echinenone pigments, contained in the hepatopancreas, epidermal layer, and carapace organs. Meanwhile, in the third fraction,



**Fig. 1.** Male Stone Crab or *Grapsus albolineatus* (Latreille in Milbert, 1812) Obtained from the Coastline of Tanawangko Village, Minahasa Regency, North Sulawesi when at the D Moulting Stage

**Table 1.** Results of Maximum UV-Vis Spectrophometric Absorption and TLC Separation Analysis Using PE and Acetone Developer Solutions (80:20) from Total Pigment Extract of *G. albolineatus* crabs (Latreille in Milbert, 1812)

F	Rf	Color - TLC	Wave Length (nm) - Maximum Spectrophotometer Absorption	Pigment Type	Description				
					G	D	H	EL	C
1	1.00	Yellow Reddish	427, 452, 478 -	$\beta$ -Carotene -	*	*	*		
2	0,91	Orange	458	Echinenone			*	*	*
3	0,82	Reddish	468	Canthaxanthin			*	*	*
4	0,64	Orange	425,450, 476	$\beta$ - cryptosanthine	*	*	*	*	*
5	0,58	Yellow	462	Adonirubine	*	*	*	*	*
6	0,54	Reddish	470	Astaxanthin	*	*	*	*	*
7	0,48	Yellow	475	Astacen	*	*	*		
8	0,41	Yellow	427	-					
9	0,22	Reddish Yellow	452	-	*	*	*	*	*

Description F = Fraction  
 Rf = Retention factor  
 G = Gonad  
 D = Hemocyanin  
 H = Hepatopancreas  
 EL = Epidermal Layer  
 C = Carapace

canthaxanthin pigments were identified with a reddish color, contained in the hepatopancreas, epidermal layer, and carapace organs. The fourth fraction showed orange color identified as cryptoxanthin pigments, found in the gonad, hemocyanin, hepatopancreas, epidermal layer, and carapace organs. The fifth fraction also produced yellow color identified as adonirubin pigments, contained in the gonad, hepatopancreas, epidermal layer, and carapace organs. In the sixth fraction, the pigments identified were astaxanthin with a reddish color, contained in the gonad, hemocyanin, hepatopancreas, epidermal layer, and carapace organs. The last or seventh fraction produced yellow color identified as astacin pigments, found in the gonad, hemocyanin and hepatopancreas organs.

According to Makalalag *et al.* (2017), female *G. albolineatus* crabs contain  $\beta$ -carotene, echinenone, canthaxanthin, adonirubin, astaxanthin, and astacin in which the quantity of the pigments is concentrated in the gonad and hemocyanin organs. Meanwhile, Paransa (2002) explained that in the total pigment extract of male *S. serrata* crabs, the carapace

and epidermal layer contain some types of pigments covering  $\beta$ -carotene, and free zeaxanthin ester and astaxanthin, as well as have antibacterial activities. Moreover, Thamin *et al.* (2006) found that  $\beta$ -carotene, echinenone, astaxanthin diester, astaxanthin monoester and free astaxanthin were contained in the total pigment extract of carapace and epidermal layer organs of male *G. albolineatus* crabs obtained from the Coastal Waters of Malalayang Dua Beach, Manado, North Sulawesi, Indonesia. His study also shows that astaxanthin and astaxanthin diester pigments of the extracted carapace organ have antibacterial activities from *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella paratyphi B*. On the other side, the results of the study by Paransa *et al.* (2014) suggest that  $\beta$ -carotene and Lutein pigments of the extracted *Sargassum polycystum* brown algae (C. Agardh) have antibacterial activity from *Escherichia coli*, *Klebsiella pneumonia*, *Salmonellaparatyphy b* and *Staphylococcus aureus*.

The pigment metabolic flow of *G. albolineatus* crabs (Latreille in Milbert, 1812) is formed through two metabolic pathways. The first pathway is the




Fig. 2. Forecast of Pigment Metabolic Cycles of *G. albolineatus* Crabs (Lamarck, 1888)

Table 2. Results of Maximum UV-Vis Spectrophometric Absorption and CC Separation Analysis with Hexane and Acetone Developer Solutions from Total Pigment Extracts of *G. albolineatus* Crab (Latreille in Milbert, 1812)

Stage I-CC Separation			Stage II-CC Separation		
BAND	WAVELENGTH (nm)	PIGMENT TYPE	BAND (nm)	WAVE LENGTH	PIGMENT TYPE
JKP1	424, 448 485	$\beta$ - Carotene	-	Not Continued	-
JKP2	465	Canthaxanthin	JKP2P1	460	Echinenone
			JKP2P2	482	Astacen
JKP3	475	Adonirubine	JKP3P1	424,449, 479	Meso-zeaxanthin
			JKP3P2	423,448, 476	Diatoksanthin
			JKP3P3	-	-
JKP4	459, 478	Fucoxanthin	JKP4P1	425,449,476	$\beta$ - cryptoxanthin
			JKP4P2	468	Astaxanthin

addition of carbonyl groups on one side of  $\beta$ -carotene pigments to form echinenone pigments. Moreover, the echinenone pigments receive carbonyl groups in other cyclic chains to form canthaxanthin pigments, to subsequently create astaxanthin and astacin pigments. In the next metabolic pathway,  $\beta$ -carotene pigments receive hydroxyl groups to form  $\beta$ -cryptoxanthin pigments and subsequently adonirubin pigments. Furthermore, major pigments, generally consisting of astaxanthin and astacin pigments, are formed in crustaceans (Mantiri *et al.* 1999).

Based on the results of the study by Mantiri *et al.* (1999), there is a pigment metabolic pathway formed in *Panulirus versicolor* lobster (Latreille, 1804), starting from  $\beta$ -carotene pigments which subsequently form echinenone, canthaxanthin and astaxanthin pigments in a series. The pigment metabolism occurs in the hemocyanin, hepatopancreas, epidermal layer, carapace and gonad organs.

#### Determination of Pigment Types Using Column Chromatography

The total carapace pigment extract of *G. albolineatus* crabs (Latreille in Milbert, 1812) captured in the research location was separated through the stage I-CC separation by using semipolar developers of hexane and acetone (80:20). The extract resulted from the CC separation was analyzed using UV-Vis spectrophotometer absorption. Then, the pigment types of the formed maximum UV-V is spectrophotometric absorption peak was identified according to Britton *et al.* (2004). The results of the stage 1-CC separation formed four bands identified as  $\beta$ -carotene, canthaxanthin, adonirubin, and fucoxanthin pigments. The results of Column Chromatography analysis using semipolar developer solutions are

shown in the table below.

The extract of canthaxanthin pigments contained echinenone and astacin pigments. Both pigment types were found through further separation using the same semipolar developer solutions. Moreover, through the CC separation, adonirubin pigments were found in the third band of the total pigment extract of carapaces of *G. albolineatus* crabs (Latreille in Milbert, 1812). The extract of adonirubin pigments was further separated using CC, resulting in meso-zeaxanthin and diatoxanthin pigments. Meanwhile, from the further CC separation of the fucoxanthin pigment extract, two bands were found, each of which was identified using maximum UV-Vis spectrophotometric absorption analysis method. The results showed that the two bands represented  $\beta$ -Cryptoxanthin and Astaxanthin pigments. Overall, the total carapace pigment extract resulted from the Stage II-CC separation showed the same pigment types with that resulted from TLC separation, namely echinenone,  $\beta$ -cryptoxanthin, astaxanthin and astacin. In conclusion, the pigments obtained from crustaceans which are utilized as natural dyes are generally taken from the carapace organ. The results of the separation of KCKT in the isolation extract of second fraction from TLC (Figure 3) are identified as canthaxanthin and Meso-zeaxanthin.

Carotenoids have a good effect on human health, serving as a precursor to vitamin A, antioxidants, and anticancer (Jaswir *et al.*, 2011). In humans, carotenoids can function as antioxidants (Yeum *et al.*, 2011). Carotenoids are known to work as well as tocopherols (vitamin E) which can stabilize reactive oxygen when photooxidative stress occurs (Stahl *et al.*, 2000). The use of carotene compounds by humans increases from year to year because carotenoids positively affect human health, such as re-




Fig. 3. HPLC chromatogram profile from KLT extract 2nd fraction of *Grapsus albolineatus* crab (Latreille in Milbert, 1812) with methanol, acetone and water solvent. The second and third peaks were identified as canthaxanthin and Meso-zeaxanthin pigments



ducing cancer risk and improving immune system function due to the role of antioxidants (Das *et al.*, 2005).

### Conclusion

*Grapsus albolineatus* is one of crab in the rocky coastline of Tanawangko and Manggatasik Village of Manado city, North Sulawesi. *Grapsus albolineatus* have high content of carotenoid pigment. Each organ of *Grapsus albolineatus* have different concentration of carotenoid. The concentration of pigment contained in carapace was 6.88 mg/g, epidermal layer 210.47 mg/g, hepatopancreas 454.0 mg/g, gonad 433.33 mg/g, and blood 153 mg/g. The pigment content of each organ *G. albolineatus*, ie at 6.95 mg carapace, epidermal layer 7.26 mg, hepatopancreas 7.07 mg, gonad 34.94 mg, and blood 5.86 mg. Carotenoid pigment of *Grapsus albolineatus* are consist of  $\beta$ -Carotene, Echinenone, Cantaxanthin, Cryptoxanthin, Adonirubin, Astaxanthin and Astacene. Carotenoid is a color pigment that have some benefit as a raw material of industry, pharmacy, or food additive.

### Acknowledgment

This research is funded in the 2018 UNSRAT PNPB DIPA budget. Therefore, all the researchers would like to say a greatfull thanks to the funders who gave the opportunity to conduct this research

### References

- Bliss, D.E. and Mantel, L.H. 1985. *Integument, Pigments and Hormonal Process in The Biology of Crustacea*. Volume 9. Academic Press Inc. Orlando-Florida.
- Britton, G., Liagen – Jansen, S. and Pfander, H. 1995. *Carotenoids*. Volume 1B. Spectroscopy. Basel, Switzerland.
- Britton, G., Liagen – Jansen, S. and Pfander, H. 2004. *Carotenoids Handbook*. Basel. Switzerland. Birkhauser Verlag AG
- Majchacheep, S. 1989. *Marine Animals of Thailand*. Published by Prae Pittaya. Bangkok, Thailand. 269 Hal.
- Mantiri, D.H.M., Paransa, D.S.J. and Mulianty Dunggio 1999. Telaah kandungan Dan jenis Pigmen Udang barong (*Panulirus versicolor*, Latreille). *Jurnal Warta – Wiptek Unsrat*. Nomor 22/10. ISSN 0854 – 0667.
- Mantiri, D.H.M. 1997. These. Nature, Localization et Metabolisme des Carotenoides et des Complexes Caratenoproteiques au Cours de L'Evolution Embryonnaire et Larvaire du Hommarid Europeen *Homarus gammarus*. Universite De Droit, D' Economie et des Sciences D' Aix Marseille.
- Mantiri, D.M.H. and Kepel, B. 1999. Beberapa Peran Pigmen Karotenoid. *Jurnal Fakultas Perikanan dan Ilmu Kelautan*. 1(3) Oktober 1999.
- Paransa, D.D.M.H, Mantiri, C. Abadi, 2000a. Penentuan Jenis dan Kandungan Pigmen Karotenoid Pada Kepiting Bakau (*Scylla serrata* Forskal, 1775) Betina dari Pesisir Likupang. *Jur. Fak. Perikanan Universitas Sam Ratulangi*. 1(4) Januari ISSN 0853-2559 : Hal 94-100
- Paransa, D.S.J., Mantiri, D.M.H. and Kasim, S. 2000b. Penentuan Kandungan Dan Jenis Pigmen Karotenoid Pada Kepiting bakau (*Scylla serrata*: Forskal 1775) Jantan, Dari Perairan Likupang. *Jur. Perikanan and Ilmu Kelautan UNSRAT*.
- Paransa, D.S.J. 2002. Uji Aktifitas secara *In vitro* Dari Ekstrak 3 Jenis Pigmen Karotenoid Pada Karapas Dan Lapisan Epidermis Kepiting Bakau *Scylla serrata* (Forskal, 1775) Jantan. *Jur Perikanan Dan Ilmu Kelautan Unsrat*. 1(2) : Hal 39 - 47
- Paransa, D.S.J., Desy, M.H. Mantiri. Florence Korompis. 2002. Penentuan Kandungan Pigmen Karotenoid Pada Kepiting *Grapsus alboneatus* (Lamarck) Betina Berdasarkan Beda Larutan Pengembang Pada Kromatografi Lapis Tipis. *Jur. Perikanan dan Ilmu Kelautan UNSRAT*. 1(3) : 1-10. ISSN 1411-9234
- Paransa D.S.J 2005. Pemurnian Jenis Pigmen Karotenoid Pada Ekstrak Kepiting (*Grapsus alboneatus*) (Lamarck) Jantan. *Jurnal Warta – WIPTEK*. No: 26/ Th.2005/Oktober ISSN : 1411-9234. Hal. 37-44
- Paransa, D. and Kemer, K. 2006. Isolasi Pigmen Xantofil Kepiting (*Grapsus alboneatus*. Lamarck) Jantan Menggunakan Prinsip Reduksi dan Saponifikasi. *Jurnal. Warta-Wiptek*. No.28/2006/Oktober.
- Paransa D. S. J., K. Kemer., A. P. Rumengan dan D. M. H. Mantiri. 2014. Analisis Jenis Pigmen Dan Uji Aktifitas Antibakteri Ekstrak Pigmen Xantofil pada alga coklat *Sargassum polycystum* (C. Agardh). *Jurnal LPPM Bidang Sains and Teknologi*. 1 (1): 90-96.

# Exploration study of the pigment types of stone crab (*Grapsus albolineatus*) as pharmaceutical and cosmetics Ingredients

## ORIGINALITY REPORT

6%

SIMILARITY INDEX

5%

INTERNET SOURCES

4%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://erepo.unud.ac.id">erepo.unud.ac.id</a> Internet Source	1%
2	<a href="http://academic.oup.com">academic.oup.com</a> Internet Source	1%
3	Submitted to Universitas Negeri Surabaya The State University of Surabaya Student Paper	1%
4	<a href="http://rjoas.com">rjoas.com</a> Internet Source	1%
5	Fatika Sari Mokoginta, Darus Saadah Johanis Paransa, Kurnati Kemer, James J. H. Paulus, Nickson J. Kawung, Henky Manoppo. "Analysis of Carotenoid Pigment Types in the Carapace of the Male Crab <i>Grapsus albolineatus</i> Latreille in Milbert 1812", Jurnal Ilmiah PLATAX, 2021 Publication	1%
6	<a href="http://123dok.com">123dok.com</a> Internet Source	<1%

7

[epjournal.edpsciences.org](http://epjournal.edpsciences.org)

Internet Source

<1 %

8

[fpik.unsrat.ac.id](http://fpik.unsrat.ac.id)

Internet Source

<1 %

9

[pubmed.ncbi.nlm.nih.gov](http://pubmed.ncbi.nlm.nih.gov)

Internet Source

<1 %

10

Irna Rustikasari, Darus S. J. Paransa, Erly Y. Kaligis, Medy Ompi, Wilmy E. Pelle, Silvester B. Pratasik. "Morphological identification of crabs in the rocky coast of Manado Bay", *Jurnal Ilmiah PLATAX*, 2021

Publication

<1 %

11

M. Lotocka. "Changes in carotenoid composition in different developmental stages of copepods: *Pseudocalanus acuspes* Giesbrecht and *Acartia* spp.", *Journal of Plankton Research*, 02/01/2004

Publication

<1 %

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On