

Growth and feed efficiency enhancement by probiotic originating from intestine of carp, *Cyprinus carpio*

by Reiny Antonetha Tumbol 6

Submission date: 10-May-2023 06:40AM (UTC+0700)

Submission ID: 2088985571

File name: Growth_and_feed_efficiency_enhancement_Manoppo_2019.pdf (265.86K)

Word count: 4220

Character count: 22825



3

Growth and feed efficiency enhancement by probiotic originating from intestine of carp, *Cyprinus carpio*

¹Henky Manoppo, ²Reiny A. Tumbol, ²Henky J. Sinjal, ³Jeane Sumaraw

⁵¹
⁴⁷ ¹ Aquaculture Study Program, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado, Indonesia; ² Aquatic Science Study Program, Graduate School of Sam Ratulangi University, Manado, Indonesia; ³ Board of Fish Quarantine, Quality Control and Safety of Fisheries Product, Manado, Indonesia. Corresponding author: H. Manoppo, hmanoppo@yahoo.com

¹⁷ **Abstract.** The objectives of this study were to evaluate the effects of probiotic isolated from digestive tracks of carp on growth and feed efficiency. Carp fingerlings with an average weight of 2.09 ± 0.27 g were gathered from Tatele Freshwater Aquaculture Fisheries Center. The probiotic *Lactobacillus* sp. was previously isolated from carp intestine at Fish Health, Environment and Toxicology Laboratory of the Fisheries and Marine Science Faculty Sam Ratulangi University Manado, Indonesia. After purification, probiotic was subsequently mixed into feed. The density of probiotic bacteria added to the feed was 1×10^6 cfu mL⁻¹, 1×10^7 cfu mL⁻¹, 1×10^8 cfu mL⁻¹, 1×10^9 cfu mL⁻¹. Commercial feed without addition of probiotic was used as a control treatment. Probiotic-supplemented feed was fed to carp fingerling having an average weight of 2.09 ± 0.27 g for 30 days at a dose of 5%/body weight, twice a day at 09.00 am and 16.00 pm. Data collected included fish weight, feed efficiency and feed conversion ratio measured at the end of the study. The results found the addition of probiotic bacteria into fish significantly affected weight gain, daily growth, feed efficiency, feed conversion ratio ($p < 0.01$) with the best results obtained in fish fed probiotic pellet supplemented with 1×10^8 cfu mL⁻¹. Thus, incorporation of probiotic bacteria isolated from carp intestines is able to enhance growth, feed efficiency as indicated by the increase in growth, feed efficiency and reduce food conversion ratio.

Key Words: food conversion ratio, friendly aquaculture, growth promoter, immunomodulator.

Introduction. Carp (*Cyprinus carpio* L.) is a freshwater fish species widely cultivated in North Sulawesi. Intensification of aquaculture in limited land with high density has caused problems especially the declining quality of the environment due to high organic matter discharge, stress on fish, poor growth, and increased disease events which result in a significant decrease in production (Manoppo et al 2016; Hoseinifar et al 2015).

Antibiotics and chemicals have long been used to prevent or treat diseases in cultured fish. This has resulted in the emergence of antibiotic resistant pathogens in aquaculture environments and accumulation of residues in aquatic products (Tan et al 2019). Ullah et al (2018) also stated that antibiotics have been proven to have harmful effects on the environment and health of fish and have been banned in many countries. In addition, disease control through the use of antibiotics/chemicals had created problems such as bioaccumulation, pollution, and suppression of the fish's immune system (Biswas et al 2012; Babu et al 2013; Wu et al 2013). Somplang & Sudthidol (2016) reported food of animal origin has been found to contain antibiotic residues in meat due to excessive use of antibiotics and if the meat is consumed by humans then the residue will be accumulated in the human body causing resistance to drugs.

At present, the use of herbs, immunostimulants, bioflocs and probiotics has attracted much attention to be used as an alternative to antibiotics and chemicals (Nguyen et al 2019; Adipu et al 2019; Dawood & Koshio 2016; Kuhn et al 2010; Crab et al 2012; Burrells et al 2011; Payung et al 2017). The main challenge of fish and shrimp farmers is to increase growth and at the same time reduce the incidence of disease. The application of probiotics is a safe alternative to antibiotics and has been used well by

farmers as feed additives and biological control agents (Ullah et al 2018; Wang et al 2018; de La Banda et al 2010; Wang et al 2008). FAO defines probiotics as living microorganisms which if given in appropriate quantities can improve fish health (FAO 2002). Prabhurajeshwar & Chandrakanth (2019) stated that probiotics are living organisms that can improve health if added to feed. Probiotic bacteria not only function as biocontrol for fish diseases, but also as an activator to increase the nutritional value of feed (Kumar et al 2013; Ullah et al 2018). Probiotics produce protease, amylase and lipase enzymes, as well as growth factors such as vitamins, fatty acids and amino acids so that absorption of nutrients becomes more efficient. Probiotics also produce antimicrobial ingredients so that they can kill or inhibit pathogens, increase growth, function as immunostimulants that can enhance the fish's immune system against disease, and regulate digestive microbial balance (Wang et al 2019). So the use of probiotics is a good management strategy to increase growth to control pathogens in fish and crustaceans.

Currently there are several commercial probiotics derived from several bacterial species such as *Bacillus* sp., *Lactobacillus* sp., *Enterococcus* sp., *Carnobacterium* sp. and yeast *Saccharomyces cerevisiae* (Cruz et al 2012). Probiotic microorganisms can originate from the host or from other sources but it is not yet known whether probiotics originating from the host are better than those from other sources (Nguyen et al 2019).

The present research has been carried out from April to September 2019 to evaluate the effects of probiotic isolated from digestive tracks of carp on growth and feed efficiency of such hosts.

Material and Method

Fish. The fish used were carp fingerlings with an average weight of 2.0-0.27 g. The fish was gathered from Freshwater Aquaculture Fisheries Center, Tatelu. The fish taken were placed in an oxygenated plastic bag, then transported to the Aquaculture Technology Laboratory of the Faculty of Fisheries and Marine Sciences for further use.

Probiotic bacteria. The probiotic used was previously isolated from the intestine of carp weighing about 100 g at the Laboratory of Fish Health, Environment and Toxicology, Faculty of Fisheries and Marine Science. Briefly, one gram of intestine was cut into small pieces, ground using a mortar and added to a 9 mL NaCl. The solution was centrifuged twice and the supernatant was removed. It was then diluted to 10^{-2} and 10^{-3} , spread onto a Petri dish containing MRS agar and placed in an incubator at 28°C for 24-48 hours. The growing bacteria were then purified through several times of reculture in the same media. The bacteria obtained were gram-positive, rod-shaped, and biochemically identified as *Lactobacillus* sp.

Feed preparation. The concentration of probiotics used as treatments consisted of 1×10^6 , 1×10^7 , 1×10^8 and 1×10^9 cfu mL⁻¹. Each treatment was supplemented by spraying thoroughly to commercial pellets (Hi-Pro-Vite FF999) as much as 1%. The composition of the feed was 35% crude protein, 2% lipid, 3% crude fiber, 13% ash, and 12% water. After air-dried, the feed was coated with 2% yellow eggs, air-dried again, placed into a plastic bag and stored at the refrigerator. Pellet without the addition of probiotics but coated with yellow eggs was used as a control treatment.

Experimental design. The experimental design used in this research was completely randomized design (CRD) with 5 treatments, each with 3 replications. The fish were acclimatized for one week before running the experiment. During the adaptation period, the fish were fed with pellet without the addition of probiotics at a dose of 5%/body weight/day. Feeding frequency was twice a day at 09.00 am and 16.00 pm.

After acclimatization process, the fish were distributed into 15 aquaria (60x40x40 cm each) with a density of 20 individuals per aquarium. Each aquarium was equipped with an aerator and a submersible water pump for recirculation system. Fish were fed probiotic-supplemented pellets that had been previously prepared for 30 days at 5% per

body weight/day, two times a day (09.00 am and 16.00 pm). Uneaten food and wastes of fish were removed by syphoning. Water replacement was also carried out every 2-3 days as much as 30% of the water volume. The parameters measured were weight gain, daily growth rate, feed efficiency, and food conversion ratio. Fish growth and feed efficiency were calculated based on the formula applied by Zokaeifar et al (2012) and Dawood et al (2016) as follows:

Weight gain, WG (g) = Wt - Wo
 Where: Wt: final weight of fish (g)
 Wo: initial weight of fish (g)

Specific growth rate, $SGR (\%) = \frac{\ln(Wt - Wo)}{t} \times 100$
 Where t: feeding duration (days)

Feed efficiency, FE (%) = $\frac{Wt - Wo}{\text{Feed intake (g)}} \times 100$

Feed conversion ratio, $FCR = \frac{\text{Total feed given (g)}}{WG}$

Data analysis. The effect of different concentrations of probiotics on growth and feed efficiency was analyzed through one-way Analysis of variance. The different effect between treatments was analyzed by Duncan's Test using SPSS 24 for windows. The effect was considered significant at p<0.05.

Result and Discussion. The addition of probiotics isolated from carp intestine into the diets with different concentrations had a very significant effect on weight gain and specific growth of carp fingerlings (p<0.01). After the fish were fed for 30 days, the best weight gain (WG) was achieved in fish fed pellet supplemented with probiotics 1x10⁸ cfu/mL (Treatment D), followed by concentrations of 1x10⁷ cfu/mL (Treatment C). The best specific growth rate (SGR) was also achieved in treatment D, then in treatment C (Table 1). Duncan test showed that WG and SGR of fish treated with treatment D and C were significantly different compared to WG and ADG of control fish (treatment A), as well as fish treated with treatment E (1x10⁹ cfu/mL) and B (1x10⁶ cfu/mL). However, there was no significant difference between treatments D and C. WG and ADG of fish treated with treatment E were significantly different from those of control fish as well as fish in treatment B.

Table 1
 Weight gain, specific growth, FCR and FE of fish fed pellet added with probiotic

Parameters	Treatments				
	A	B	C	D	E
Final weight (g)	3.89±0.39	4.04±0.37	5.64±0.11	6.09±0.10	5.20±0.17
WG (g)	1.80±0.39 ^a	1.94±0.37 ^a	3.55±0.11bc	4.00±0.10 ^c	3.11±0.17 ^b
SGR (%)	2.20±0.36 ^a	2.33±0.33 ^a	3.54±0.07 ^{bc}	3.82±0.06 ^c	3.25±0.11 ^b
FE (%)	48.72±7.57 ^a	49.55±7.27 ^a	82.68±1.43 ^b	89.68±6.44 ^b	77.73±7.71 ^b
FCR	2.08±0.32 ^a	2.05±0.31 ^a	1.21±0.02 ^b	1.12±0.08 ^b	1.29±0.12 ^b

Means with different superscript in the same row are significantly different. A - control without addition of probiotic; B - probiotic 1x10⁶ cfu mL⁻¹; C - probiotic 1x10⁷ cfu mL⁻¹; D - probiotic 1x10⁸ cfu mL⁻¹; E - probiotic 1x10⁹ cfu mL⁻¹; WG - weight gain (g); SGR - specific growth rate (%); FCR - feed conversion ratio; FE - feed efficiency (%).

The present research proves that probiotics originating from the digestive organs of carp have the potential to spur fish growth. Previous research report showed that probiotic isolated from catfish intestines can increase the growth of carp juveniles (Manoppo et al 2019). Probiotics isolated from the digestive tract of gourami (*Osphronemus goramy*) had

also been reported to increase the growth of tilapia (Mulyasari et al 2016). In red sea bream (*Pagrus major*), supplementation of *Lactobacillus rhamnosus* in feed and given for 56 days had a significant effect on the increase of fish growth and immunity (Dawood et al 2016). Furthermore, Lin et al (2019) reported that giving *Lactobacillus* spp. to Lined sea horse (*Hippocampus erectus*) had a significant influence on growth, feed utilization, feed digestibility, protease activity, and fish immune response. Several research reports also proved that probiotics derived from fish digestive organs could induce growth, feed conversion ratio and feed efficiency (Mohammadian et al 2019; Doan et al 2018; Marlida et al 2014).

In the fish given probiotic with higher concentration (Treatment E), fish growth tended to be slower than the growth of fish in lower concentration (treatment D and C). This occurred because of the probiotic concentration added had probably exceeded the required dose. Conversely at lower dose (treatment B), administration of probiotic had no effect on the growth. It was suspected that the amount of probiotics given was not enough to function effectively in promoting fish growth. This condition occurred because the administration of an ingredient such as probiotic was strongly influenced by the dose and duration of administration (Sakai 1999).

Supplementation of probiotic into feed with different concentrations also had a very significant effect on feed conversion ratio (FCR) and feed efficiency (FE) ($p < 0.01$). After feeding for 30 days, the best FCR value was achieved in fish in treatment D, then C and E. The best FE value was also observed in fish in treatment D, followed by C and E. Based on Duncan's Test, the FCR and FE values in these three treatments were not significantly different one to each other, but significantly different if compared with control fish and fish in treatment B. There were no significant differences between treatments A and B (Table 1).

The present research proved that probiotics isolated from the carp intestine could reduce the FCR and increase the FE value. Similar research reports showed that probiotic (*Lactobacillus* sp.) isolated from digestive organ of catfish and given to carp for 30 days significantly increased FCR and FE (Manoppo et al 2019). Probiotic isolated from the digestive tract of Humpback groupers could also reduce the value of feed conversion, increase growth, protein and fat retention (Marlida et al 2014). In addition, several studies found that the use of probiotics in aquaculture could improve growth, digestibility and feed efficiency, appetite, the fish's immune system, tolerance to stress and resistance to pathogens (Midhun et al 2019; Mulyasari et al 2016; Ling et al 2018; Rahmawan et al 2014; Hemaiswarya et al 2013). The report by Midhun et al (2019) showed the addition of probiotics in feed both single and in combination between *Lactobacillus plantarum* N11 and *Bacillus velezensis* H3.1 improved growth performance, specific growth rate, weight gain, final weight and feed conversion ratio in tilapia. Ullah et al (2018) reported that the use of commercial probiotics significantly increased growth, digestive enzymes activities and immune response of Mori fish (*Cirrhinus mrigala*) in the polyculture system. Furthermore, Tan et al (2019) reported that supplementation of probiotics (*Rummeliibacillus stabekisii*) isolated from the stomach of tilapia into the feed and given to tilapia (mean 4.1 ± 0.34 g) for 8 weeks increased fish growth, immunity, disease resistance and gut microflora and health status of tilapia. In shrimp, Kongnum & Hongpattarakere (2012) reported that the addition of probiotics isolated from wild shrimp and fed for 6 weeks significantly increased the relative growth rate, reduced FCR and increased the survival of *Penaeus vannamei*, as compared to control shrimp. Wang et al (2019) found that pellet supplemented with a mixture of several species of probiotics for 56 days improved growth performance, survival and health status of shrimp (*P. vannamei*). Xie et al (2019) also reported the use of a mixture of probiotics in white shrimp increased growth, FE, nonspecific immune response, digestive enzymes activity, and reduce FCR.

Conclusions. Supplementation of probiotic isolation from digestive tract of carp able to enhance growth performance and feed efficiency as indicated by the increase in weight gain, specific growth, feed efficiency and reduce food conversion ratio. The best results were obtained in fish fed pellet with the addition of 1×10^8 cfu mL⁻¹.

Acknowledgements. The authors thank to the Ministry of research, technology and higher education of the Republic of Indonesia for the financial support through the fiscal year 2019. Special thank was also addressed to the head of Aquaculture Technology Laboratory of Faculty of Fisheries and Marine Science for allowing us to use the laboratory facilities during the research.

References

- Adipu Y., Lumenta C., Mangindaan R. E. P., Manoppo H., 2019 Growth performance of *Litopenaeus vannamei* grown in biofloc system produced from different carbohydrate sources. *AAFL Bioflux* 12(2):472-479.
- Babu D. T., Antony S. P., Joseph S. P., Bright A. R., Philip R., 2013 Marine yeast *Candida aquaetextoris* S527 as a potential immunostimulant in black tiger shrimp *Penaeus monodon*. *Journal of Invertebrate Pathology* 122:243-252.
- Biswas G., Korenaga H., Takayama H., Kono T., Shimokawa H., Sakai M., 2012 Cytokine responses in the common carp, *Cyprinus carpio* L. treated with baker's yeast extract. *Aquaculture* 356-357:169-175.
- Burrels C., Williams P. D., Fomo P. F., 2001 Dietary nucleotide: A novel supplement in fish feed effects on resistance to disease in Salmonids. *Aquaculture* 199:159-169.
- Crab R., Defoirdt T., Bossier P., Verstraete W., 2012 Biofloc technology in aquaculture: Beneficial effects and future challenges. *Aquaculture* 356-357:351-356.
- Cruz P. M., Ibanez A. L., Hermosillo O. A. M., Saad H. C. R., 2012 Use of probiotics in aquaculture. *International Scholarly Research Network, ISRN Microbiology*, Volume 2012, Article ID 916845, 13 pages doi:10.5402/2012/916845.
- Dawood M. A. O., Koshio S., 2016 Recent advances in the role of probiotics and prebiotics in carp aquaculture: A review. *Aquaculture* 454:243-251
- Dawood M. A. O., Koshio S., Ishikawa M., Yokoyama S., Basuini M. E. E., Hossain M. S., Nhu T. H., Dossou S., Moss A. S., 2016 Effect of dietary supplementation of *Lactobacillus rhamnosus* or/and *Lactococcus lactis* on the growth, gut microbiota and immune responses of red sea bream, *Pagrus major*. *Fish and Shellfish Immunology* 49:275-285.
- de La Banda I. G., Lobo C., Leon-Robio J. M., Tapia-Paniagua S., Balebona M. C., Morinigo M. A., Moreno-Ventas X., Lucas L. M., Linares F., Arce F., Arijó S., 2010 Influence of two closely related probiotics on juvenile Senegalese sole (*Solea senegalensis*, Kaup 1858) performance and protection against *Photobacterium damsela* subsp. *piscicida*. *Aquaculture* 306:281-288.
- Doan H. V., Hoseinifar S. H., Khanongnuch C., Kanpiengjai A., Srichaiyo S., 2018 Host-associated probiotics boosted mucosal and serum immunity, disease resistance and growth performance of Nile tilapia (*Oreochromis niloticus*). *Aquaculture* 491:94-100.
- Hemaiswarya S., Raja R., Ravikumar R., Carvalho I. S., 2013. Mechanism of action of probiotics. *Brazilian Archives of Biology and Technology* 56(1):113-119.
- Hoseinifar S. H., Esteban M. A., Cuesta A., Yun-Zhang Sun Y. Z., 2015 Prebiotics and fish immune response: A review of current knowledge and future perspectives. *Reviews in Fisheries Science & Aquaculture* 23:315-328.
- Kongnum K., Hongpattarakere T., 2012 Effect of *Lactobacillus plantarum* isolated from digestive track of wild shrimp on growth and survival of white shrimp (*Litopenaeus vannamei*) challenged with *Vibrio harveyi*. *Fish and Shellfish Immunology* 32:170-177.
- Kuhn D. D., Lawrence A. L., Boardman G. D., Patnaik S., Marsh L., Flick Jr. G. J., 2010 Evaluation of two types of bioflocs derived from biological treatment of fish effluent as feed ingredients for pacific white shrimp, *Litopenaeus vannamei*. *Aquaculture* 303:28-33.
- Kumar S., Raman R. P., Pandey P. K., Mohanty S., Kumar A., Kumar K., 2013 Effect of orally administered azadirachtin on non-specific immune parameters of goldfish *Carassius auratus* (Linn. 1758) and resistance against *Aeromonas hydrophila*. *Fish and Shellfish Immunology* 34:564-573.

- Lin T., Liu X., Xiao D., Zhang D., Cai Y., Zhu X., 2019 *Lactobacillus* spp. as probiotics for prevention and treatment on enteritis in the lined seahorse (*Hippocampus erectus*) juveniles. *Aquaculture* 503:16-25.
- Ling Y., Zhang R., Ke C., Hong G., 2018 Effects of dietary supplementation of probiotics on growth, immune responses, and gut microbiome of the abalone *Haliotis diversicolor*. *Aquaculture* 493:289-295.
- Manoppo H., Tumbol R. A., Sinjal H. J., Novitarizky I. A., 2019 The use of probiotic isolated from Sangkuriang catfish (*Clarias gariepinus* var. Sangkuriang) intestine to improve growth and feed efficiency of carp, *Cyprinus carpio*. *AAFL Bioflux* 12(1):239-245.
- Manoppo H., Kolopita M. E. F., Malatunduh R., 2016 Growth promoter effect of garlic (*Allium sativum*) on carp (*Cyprinus carpio* L). *International Journal of PharmTech Research* 9(4):283-288.
- Marlida R., Suprayudi M. A., Widanarni, Harris E., 2014 Isolation, selection and application of probiotic bacteria for improvement the growth performance of humpback groupers (*Cromileptes altivelis*). *International Journal of Science: Basic and Applied Research* 16(1):364-379.
- Midhun S. J., Arun D., Neethu S., Vysakh A., Radhakrishnan E. K., Jyothis M., 2019 Administration of probiotic *Paenibacillus polymyxa* HGA4C induce morphometric, enzymatic and gene expression change in *Oreochromis niloticus*. *Aquaculture* 508: 52-59.
- Mohammadian T., Nasirpour M., Tabandeh M. R., Mesbah M., 2019 Synbiotic effects of B-glucan, mannan oligosaccharide and *Lactobacillus casei* on growth performance, intestine enzymes activities, immune-hematological parameters and immune-related gene expression in common carp, *Cyprinus carpio*: an experimental infection with *Aeromonas hydrophila*. *Aquaculture Volume* 511, 15 September 2019, 634197.
- Mulyasari, Widanarni, Suprayudi M. A., Zairin Jr. M., Sunarno M. T. D., 2016 Screening of probiotics from the digestive tract of gouramy (*Osphronemus goramy*) and their potency to enhance the growth of tilapia (*Oreochromis niloticus*). *AAFL Bioflux* 9(5):1121-1132.
- Nguyen T. L., Park C. I., Do-Hyung Kim D. H., 2019 Improved growth rate and disease resistance in olive flounder, *Paralichthys olivaceus*, by probiotic *Lactococcus lactis* WFLU12 isolated from wild marine fish. *Fish and Shellfish Immunology* 471:113-120.
- Payung C. N., Tumbol R. A., Manoppo H., 2017 Dietary ginger (*Zingiber officinale*) enhance resistance of Nile tilapia (*Oreochromis niloticus*) against *Aeromonas hydrophila*. *AAFL Bioflux* 10(4):962-968.
- Prabhurajeshwar C., Chandrakanth K., 2019. Evaluation of antimicrobial properties and their substances against pathogenic bacteria in-vitro by probiotic Lactobaccilli strains isolated from commercial yogurt. *Clinical Nutrition Experimental* 23:97-115.
- Rahmawan M. E. A, Suminto V., Herawati E., 2014 Application of probiotic into artificial food to improve feed efficiency, growth and survival rate of Dumbo catfish. *Journal of Aquaculture Management and Technology* 3(4):257-264.
- Sakai M., 1999. Current research status of fish immunostimulants. *Aquaculture* 172:63-92.
- Sornplang P., Sudthidol P., 2016 Probiotic isolates from unconventional sources: A review. *Journal of Animal Science and Technology* 58: 26, doi: 10.1186/s40781-016-0108-2.
- Tan H. Y., Sai-Wei Chen S. W., Shao-Yang Hu S. Y., 2019 Improvements in the growth performance, immunity, disease resistance, and gut microbiota by the probiotic *Rummeliibacillus stabekisii* in Nile tilapia (*Oreochromis niloticus*). *Fish and Shellfish Immunology* 92:265-275.
- Ullah A., Zuberi A., Ahmad M., Shah A. B., Younus N., Ullah S., Khattak M. N. K., 2018 Dietary administration of the commercially available probiotics enhanced the survival, growth, and innate immune responses in Mori (*Cirrhinus mrigala*) in a natural earthen polyculture system. *Fish and Shellfish Immunology* 72:266-272.
- Wang H., Wang C., Tang Y., Sun B., Huang J., Song X., 2018 Pseudoalteromonas probiotics as potential biocontrol agents improve the survival of *Penaeus vannamei*

- challenged with acute hepatopancreatic necrosis disease (AHPND)-causing *Vibrio parahaemolyticus*. *Aquaculture* 494:30-36.
- Wang Y. C., Hu S. Y., Chiu C. S., Liu C. H., 2019 Multiple-strain probiotics appear to be more effective in improving the growth performance and health status of white shrimp, *Litopenaeus vannamei*, than single probiotic strains. *Fish and Shellfish Immunology* 84:1050-1058.
- Wang Y. B., Li J. R., Lin J., 2008 Probiotics in aquaculture: Challenges and outlook. *Aquaculture* 281:1-4.
- Wu Y. R., Gong Q. F., Fang H., Liang W. W., Chen M., He R. J., 2013 Effect of *Sophora flavescens* on non-specific immune response of tilapia GIFT (*Oreochromis niloticus*) and disease resistance against *Streptococcus agalactiae*. *Fish and Shellfish Immunology* 34:220-227.
- Xie J. J., Liu Q. Q., Liao S., Fang H. H., Yin P., Xie S. W., Tian L. X., Liu Y. J., Niu J., 2019 Effects of dietary mixed probiotics on growth, non-specific immunity, intestinal morphology and microbiota of juvenile pacific white shrimp, *Litopenaeus vannamei*. *Fish and Shellfish Immunology* 90:456-465.
- Zokaeifar H., Balcázar J. L., Saad C. R., Kamarudin M. S., Sijam K., Arshad A., Nejat N., 2012 Effects of *Bacillus subtilis* on the growth performance, digestive enzymes, immune gene expression and disease resistance of white shrimp, *Litopenaeus vannamei*. *Fish and Shellfish Immunology* 33:683-689.
- *** FAO, 2002 Guidelines for the evaluation of probiotics in food. London, Ontario, Canada. Available at: https://www.who.int/foodsafety/fs_management/en/probiotic_guidelines.pdf. Accessed: February, 2019.

Received: 08 November 2019. Accepted: 16 December 2019. Published online: 23 December 2019.

Authors:

Henky Manoppo, Sam Ratulangi University, Faculty of Fisheries and Marine Science, Aquaculture Study Program, Indonesia, 95115 Manado, e-mail: hmanoppo@yahoo.com

Reiny Antonetha Tumbol, Graduate School of Sam Ratulangi University, Aquatic Science Study Program, Indonesia, 95115 Manado, e-mail: reinytumbol@yahoo.com

Henky Julius Sinjal, Graduate School of Sam Ratulangi University, Aquatic Science Study Program, Indonesia, 95115 Manado, e-mail: juliushengky@yahoo.com

Jeane Sumaraw, Board of Fish Quarantine, Quality Control and Safety of Fisheries Product, Indonesia, 95258 Manado, e-mail: jeanesu1104@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Manoppo H., Tumbol R. A., Sinjal H. J., Sumaraw J., 2019 Growth and feed efficiency enhancement by probiotic originating from intestine of carp, *Cyprinus carpio*. *AAFL Bioflux* 12(6):2188-2194.

Growth and feed efficiency enhancement by probiotic originating from intestine of carp, *Cyprinus carpio*

ORIGINALITY REPORT

20%

SIMILARITY INDEX

14%

INTERNET SOURCES

14%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	jurnal.unigo.ac.id Internet Source	1%
2	mail.scialert.net Internet Source	1%
3	www.proquest.com Internet Source	1%
4	Toledo, Tárík Massucci, Bruno Corrêa Silva, Felipe do Nascimento Vieira, José Luiz Pedreira Mouriño, and Walter Quadros Seiffert. "Effects of different dietary lipid levels and fatty acids profile in the culture of white shrimp <i>Litopenaeus vannamei</i> (Boone) in biofloc technology: water quality, biofloc composition, growth and health", <i>Aquaculture Research</i> , 2014. Publication	1%
5	docsdrive.com Internet Source	1%
6	onlinelibrary.wiley.com Internet Source	

1 %

7

www.deepdyve.com

Internet Source

1 %

8

performfish.eu

Internet Source

1 %

9

www.semanticscholar.org

Internet Source

1 %

10

Mohammad Jalil Zorriehzahra, Somayeh Torabi Delshad, Milad Adel, Ruchi Tiwari, K. Karthik, Kuldeep Dhama, Carlo C. Lazado. "Probiotics as beneficial microbes in aquaculture: an update on their multiple modes of action: a review", *Veterinary Quarterly*, 2016

Publication

<1 %

11

Submitted to Padjadjaran University

Student Paper

<1 %

12

Ahmed H. Sherif, Mofeed Y. Gouda, Eman T. Al - Sokary, Mohamed M. Elseify. " enhances immunity of Nile tilapia challenged with ", *Aquaculture Research*, 2020

Publication

<1 %

13

Intan A. Novitarizky, Henky Manoppo, Sammy N.J. Longdong. "Isolasi bakteri probiotik

<1 %

Lactobacillus sp dari usus ikan mas (Cyprinus carpio)", e-Journal BUDIDAYA PERAIRAN, 2018

Publication

14

Submitted to Lira University

Student Paper

<1 %

15

Maria B. Flores-Pérez, Enrico A. Yépez, Agustín Robles-Morúa, Martín Villa-Ibarra et al. "Eco-efficiency assessment of disease-infected shrimp farming in Mexico using environmental impact assessment tools", Science of The Total Environment, 2023

Publication

<1 %

16

Monika M. Wantah, Sammy N.J. Longdong, Reni L. Kreckhoff, Desy M.H. Mantiri, Reiny A. Tumbol, Henky Manoppo. "Efikasi ekstrak tanaman obat Binahong Anredera cordifolia (Ten) Steenis untuk meningkatkan pertumbuhan ikan Nila Oreochromis niloticus", e-Journal BUDIDAYA PERAIRAN, 2018

Publication

<1 %

17

www.acarindex.com

Internet Source

<1 %

18

www.omicsonline.org

Internet Source

<1 %

19

fas.biomedcentral.com

Internet Source

<1 %

20

peerj.com
Internet Source

<1 %

21

uokerbala.edu.iq
Internet Source

<1 %

22

vdoc.pub
Internet Source

<1 %

23

Henky Manoppo. "Uji efektivitas ekstrak daun Andong (*Cordyline fruticosa*) sebagai bahan antimikroba alami", e-Journal BUDIDAYA PERAIRAN, 2020
Publication

<1 %

24

Hien Van Doan, Seyed Hossein Hoseinifar, Chartchai Khanongnuch, Apinun Kanpiengjai et al. "Host-associated probiotics boosted mucosal and serum immunity, disease resistance and growth performance of Nile tilapia (*Oreochromis niloticus*)", *Aquaculture*, 2018
Publication

<1 %

25

Y S Ismail, Febriani, C Yulvizar, R Ramadhani. " Identification of the bacterium isolate from Mackerel Fish sp.) using 16S rRNA Gene ", IOP Conference Series: Earth and Environmental Science, 2019
Publication

<1 %

26

dspace.lpu.in:8080
Internet Source

<1 %

- | | | |
|----|--|------|
| 27 | eclass.upatras.gr
Internet Source | <1 % |
| 28 | eprints2.undip.ac.id
Internet Source | <1 % |
| 29 | erl.ucc.edu.gh:8080
Internet Source | <1 % |
| 30 | journals.plos.org
Internet Source | <1 % |
| 31 | Elham Awad, Amani Awaad. "Role of medicinal plants on growth performance and immune status in fish", <i>Fish & Shellfish Immunology</i> , 2017
Publication | <1 % |
| 32 | Hien Van Doan, Seyed Hossein Hoseinifar, Chanagun Chitmanat, Sanchai Jaturasitha et al. "The effects of Thai ginseng, <i>Boesenbergia rotunda</i> powder on mucosal and serum immunity, disease resistance, and growth performance of Nile tilapia (<i>Oreochromis niloticus</i>) fingerlings", <i>Aquaculture</i> , 2019
Publication | <1 % |
| 33 | Maedeh Rashmehi, Seyed Pezhman Hosseini Shekarabi, Mehdi Shamsaie Mehrgan, Hamed Paknejad. "Assessment of dietary chaste tree (<i>Vitex agnus-castus</i>) fruit extract on growth performance, hemato-biochemical parameters, and mRNA levels of growth and | <1 % |

appetite-related genes in goldfish (Carassius auratus)", Aquaculture and Fisheries, 2021

Publication

34

Md. Abdul Kader, Nursyafiqah Zahidah Azahar, Shumpei Iehata, Mahbuba Bulbul et al. " Dietary supplementation of host - associated lactic acid bacteria modulates growth, metabolic activities, and immune - related gene expression in giant freshwater prawn, ", Journal of the World Aquaculture Society, 2020

Publication

35

Mulyasari, I Taufik, L Setijaningsih, Subaryono. "Effect of agar and alginate on growth performance and levels of leukocytes and erythrocytes of Tilapia Oreochromis niloticus", IOP Conference Series: Earth and Environmental Science, 2021

Publication

36

Rocío Simón, Félix Docando, Noelia Nuñez-Ortiz, Carolina Tafalla, Patricia Díaz-Rosales. "Mechanisms Used by Probiotics to Confer Pathogen Resistance to Teleost Fish", Frontiers in Immunology, 2021

Publication

37

ciba.res.in
Internet Source

<1 %

<1 %

<1 %

<1 %

38

Internet Source

<1 %

39

japu.gau.ac.ir

Internet Source

<1 %

40

kb.psu.ac.th

Internet Source

<1 %

41

ntnuopen.ntnu.no

Internet Source

<1 %

42

scholarworks.aub.edu.lb

Internet Source

<1 %

43

sciendo.com

Internet Source

<1 %

44

www.researcherslinks.com

Internet Source

<1 %

45

Milad Adel, Abdel-Fattah M. El-Sayed, Sakineh Yeganeh, Maryam Dadar, Sib Sankar Giri.

"Effect of Potential Probiotic Lactococcus lactis Subsp. lactis on Growth Performance, Intestinal Microbiota, Digestive Enzyme Activities, and Disease Resistance of Litopenaeus vannamei", Probiotics and Antimicrobial Proteins, 2016

Publication

<1 %

46

Sib Sankar Giri, Jin Woo Jun, Saekil Yun, Hyoun Joong Kim et al. "Effects of dietary heat-killed Pseudomonas aeruginosa strain VSG2 on

<1 %

immune functions, antioxidant efficacy, and disease resistance in *Cyprinus carpio*", *Aquaculture*, 2019

Publication

47

Suryadi N.N. Tatura, Stefanus Gunawan, Janno Bernadus, Sianne Sandjoto.

"*Plasmodium falciparum* found in the bone marrow of a child in Manado City, East Indonesia: A case report", *Asian Pacific Journal of Tropical Medicine*, 2017

Publication

48

da Silva, Bruno Corrêa, Felipe do Nascimento Vieira, José Luiz Pedreira Mouriño, Norha Bolivar, and Walter Quadros Seiffert.

"Butyrate and propionate improve the growth performance of *Litopenaeus vannamei*", *Aquaculture Research*, 2014.

Publication

49

repository.ub.ac.id

Internet Source

50

A. Mikani, H.R. Etebarian, H. Aminian, A.

Alizadeh. "Interaction Between *Pseudomonas fluorescens* Isolates and Thiabendazole in the Control of Gray Mold of Apple", *Pakistan Journal of Biological Sciences*, 2007

Publication

51

Diana Shintawati Purwanto, Yanti Meilen Mewo, Edmond Leonard Jim, Richardo Jordan

<1 %

<1 %

<1 %

<1 %

<1 %

Laloan et al. "The Analysis of Electrolyte Levels on Hypertensive Patients in Noongan Regional General Hospital, North Sulawesi, Indonesia", Science Journal of Clinical Medicine, 2019

Publication

52

Patricia Martínez Cruz, Ana L. Ibáñez, Oscar A. Monroy Hermosillo, Hugo C. Ramírez Saad. "Use of Probiotics in Aquaculture", ISRN Microbiology, 2012

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On