

Sedimentation Rate and Total Suspended Solid (TSS) In Melombo Area, Salurang Village, Sangihe Archipelagic Regency

by Lefrand Manoppo 2

Submission date: 17-May-2023 10:03AM (UTC+0700)

Submission ID: 2095114331

File name: 3._Sedimentation_Rateand_Total_Suspended_Solid_TSS.pdf (529.82K)

Word count: 4770

Character count: 22315

Sedimentation Rate and Total Suspended Solid (TSS) In Melombo Area, Salurang Village, Sangihe Archipelagic Regency

Lefrand Manoppo¹, Marsoedi², S. Muhammad², S. Berhimpon⁴

¹ Doctorate Program Student of Agricultural Sciences

^{2,3} Teaching Staff of Brawijaya University, Malang

⁴ Teaching Staff of Sam Ratulangi University, Manado

Abstract: This study was aimed to know the sedimentation rate and the total suspended solid of the estuary in relation to the implementation of melombo culture practice area. The sedimentation rate was measured using sediment traps placed in 3 set station and data were taken for 8 months of two weeks interval. The sediment traps were made of PVC pipe, 11.5 cm long and 5 cm diameter and located at 20 cm above the sea bottom. Sediment samples were firstly soaked in freshwater for approximately 4 hours to remove salt content. Sedimentation rate was determined as m^2/h time length, while Total Suspended Solid (TSS) followed the formula of Eaton, et al (1995). Results showed that the sediment trapped in station 1 ranged from 5.6 cm (109.9 cm^3) – 7.7 cm (151.11 cm^3) with an average of 6.61 cm (129.7 cm^3), station 2 from 5.3 cm (104 cm^3) to 8.5 cm (166.8 cm^3) with an average of 7.18 cm (140.9 cm^3) and station 3 from 3.8 cm (74.6 cm^3) to 7.6 cm (149.2 cm^3) with an average of 6.8 cm (133.8 cm^3), respectively. Total suspended solids (TSS) ranged from 18.28 to 50.60 with an average of 29.165 (29.59%) for station 1, 27.84 to 47.48 with an average of 38.99 (39.56%) for station 2, and 18.28 to 50.60 with an average of 30.405 (30.85%) for station 3, respectively. Based upon the decree of Living Environment Minister Numbered 51, 2004, for marine biota, the TSS has been above the standard seawater quality.

Keywords: Melombo, Sediment traps, sedimentation rate, TSS.

I. Introduction

Sangihe Archipelagic Regency is geographically located between $4^{\circ} 4' 13'' - 4^{\circ} 44' 22''$ U and $125^{\circ} 9' 28'' - 125^{\circ} 56' 57''$ E, at the position between Sulawesi Island and Mindanao Island (Philippine Republic). Its capital is Tahuna with a distance of 142 sea miles from Manado, the capital of North Sulawesi. It has a total area of 11,863.58 km^2 consisting of land of 736.98 km^2 or 6.2 % (60 % flat, 40 % slope) and sea of 11,126.61 km^2 (93.8%). Forest area is about 13,820 Ha comprising protected forests of $\pm 12,672$ Ha and mangrove forests of $\pm 1,148$ Ha. Field conditions showed that the forests have mostly become critical land due to plantation, protected forest clear-cut, timber theft and other damages. Recent data show that the critical land in the forest area is $\pm 9,322$ Ha, and that outside the forest is $\pm 53,213$ Ha (BAPPEDA, 2011).

As an archipelagic area in general, this region often has natural phenomena that could sometimes cause very high loss, such as sea storm/tidal wave, tornado, volcano, flood and landslide. The climate is influenced by monsoons, with mean air temperature per month of 27°C over 2010 following the measurements of Naha Meteorological Station. Topographic condition is mountainous and hilly terrestrial area with labile land (latosol and aluvial) with 0 - 750 m above sea level and a transpacific volcano chain from Hawaii, Japan, Philippines ending in Mollucas so that the area is vulnerable to natural disaster.

Melombo area is an estuary located between coral reefs and terrestrial in the river mouth of Salurang. Estuary is a body of water where occurring mixtures of tide affected-seawater and freshwater from the land so that this water condition is highly dependent upon the seawater and freshwater conditions going into the area. The higher the suspended materials contained in the water the higher the mud deposition in an estuary. Nybakken (1992) stated that deposit formation got also marine influences, since seawater contains sufficient number of suspended materials.

Recent increase in human activities along Salurang watershed has had some impact on the estuarine ecosystem. These are illegal gold mining in Salurang watershed and forest clear cut in the upstream. Gold mining activities were done by taking apart of the mound using high pressure pumps. These activities bring about increased soil erosion along the watershed. As a result, number of sediment (suspended solid) in the river increases and makes superficiality in melombo area. Factors affecting the sedimentation process in the river mouth are waves and current patterns (Efrijeldi, 1999).

Sedimentation is the entry of sediment load in a certain aquatic environment through water medium and suspended in the environment. Coastal sedimentation will be a problem if it occurs in locations where there are human activities requiring relatively better water conditions with sufficient depth, such as melombo area, which is scad fishing ground, and sailing paths or those who need clear water condition, such as marine

tourisms, coral reef ecosystem or seagrass beds. In other words, sediment is defined as materials concentrated in a water mass, either in organic or inorganic forms. Factors affecting sedimentation in the rivermouth are watershed topography, climate, soil type and texture, river morphometric, hydrological system and tidal energy in the rivermouth (Pickard, 1967).

Indonesia coastal waters possess various important roles, such as food source, transportation mode and harbour, industrial area, agroindustry, tourism, residential area and fisheries (Dahuriet al., 2001). Basic concept of fishing fisheries development is Indonesia national sustainable development concept, including natural resources and environmental protection (Men. LH, 1997), since the coastal area holds various problems, from pollution to habitat degradations, and other longer implication, such as sea level change.

Melombo waters, in the range of 0.5 – 5.0 m deep at low tide (Appendix 1), is an implementation area of melombo activities in Salurang village located in the district of Central South Tabukan, Sangihe Archipelagic Regency. The use of melombo coastal area is for yellowstripe scad (*Selaroides leptolepis*) fishing activity and local sea transportation in which these activities could result in some impact on the sustainability of fish exploitation and natural resources that will eventually put strong ecological pressures due to water quality decline.

The sustainability of local wisdom “melombo” in the coastal area of Salurang is highly determined by the environmental condition of the estuary, and therefore, the aquatic physical condition study was carried out in order to promote the sustainability of the cultural wisdom melombo directing to know the sedimentation rate and the total suspended solid of the estuary on the sustainability of cultural wisdom, melombo.

II. Method

Sediment measurements were done by placing sediment traps at 3 established stations, then checked at a 2 week interval for 8 months. The sediment traps used were made of PVC pipe, 11.5 cm long and 11 cm diameter, and then placed at 20 cm above the sea bottom. Moreover, the sediment trap placement used a 40 cm x 40 cm x 6 cm concrete block which upper part attached a 20 cm cylindrical concrete block as a base of the PVC pipe placement put on the sea bottom (Fig. 1). Trap construction followed Gardner (1980) in Manengkei (2003), and it was laid at 3 – 4 m depth.

Sediment samples collected in the sediment traps were measured the height trapped, then put into the plastic bag and held in the iced box for laboratory analysis. Before sediments were measured/weighed, the samples were previously soaked in freshwater for approximately 4 jam to remove salt content in clean sediment. Then, the sediment at each station was diluted to be 250 ml solution, taken 10% and placed on a Ø 47 mm and 0.45 µm mesh size “Whatman” filter paper whose weight was known. The sediment was then placed on the filter and water filtering used ‘vac Gast Benton Harbon, MI’-typed vacuum pump, weighed and dried in a “Mommert” oven at 90°C for 24 hours. Afterwards, final weight was taken to obtain the dry weight.

Sedimentation rate calculations used the $\pi r^2 h$ formula divided by length of time, where π = constant, r = ½ pipe diameter, h = sediment height in the pipe, but Total Suspensi Solid (TSS) was analyzed using the formula of Eaton et al (1995) as follows:

$$\text{TSS (mg/L)} = \frac{(A-B) \times 1000}{\text{Sample volume (mg/L)}}$$

where A = filter weight + dry residue weight (mg) and B = filter weight

III. Results and Discussion

A. Sedimentation Rate

Based on the sediment falling into the sediment trap, the sediment height in station 1 ranged from 5.6 cm (109.9 cm³) – 7.7 cm (151.11 cm³) with an average of 6.61 cm (129.7 cm³), station 2 from 5.3 cm (104 cm³) – 8.5 cm (166.8 cm³) with an average of 7.18 cm (140.9 cm³) and station 3 from 3.8 cm (74.6 cm³) – 7.6 cm (149.2 cm³) with an average of 6.8 cm (133.8 cm³), respectively (Table 1 and Fig. 2).

The highest sediment was recorded in station 2, 7.18 cm, due to its position in the rivermouth, then followed by station 3. It could probably result from mining activities in Salurang watershed and global climate change that affects the rainfalls in Sangihe island. Based on meteorological data, rainfalls from March to September were difficult to predict due to natural anomaly from global climate change.




Figure 1. Sediment Traps

Table 1. Sediment height distribution in each sediment trap from March to September 2012

Observation	Station					
	(1) 03°28'33.6"N 125°40'00.6"S		(2) 03°28'27.4"N 125°39'57.5"S		(3) 03°28'30.3"N 125°39'58.4"S	
	Height (cm)	Volume (cm ³)	Height (cm)	Volume (cm ³)	Height (cm)	Volume (cm ³)
1	7,3	143.26	7,1	139.34	7,5	147.19
2	6,3	123.64	6,4	125.60	7,6	149.15
3	6,5	127.56	7,7	151.11	7,2	141.30
4	6,7	131.49	6,9	135.41	6,7	131.49
5	5,9	115.79	7,1	139.34	7,1	139.34
6	6,3	123.64	7,3	143.26	7,0	137.34
7	6,2	121.67	7,2	141.30	6,9	135.41
8	6,9	135.41	7,3	143.26	6,8	133.45
9	6,3	123.64	7,5	147.19	7,6	149.15
10	6,4	125.60	7,7	151.11	5,5	107.94
11	5,6	109.90	5,3	104.01	3,8	74.57
12	6,6	129.53	7,0	137.37	7,2	141.30
13	7,2	141.30	7,2	141.30	7,0	137.37
14	7,7	151.11	8,5	166.81	7,2	141.30
15	6,9	135.41	7,1	139.34	6,8	133.45
16	7,0	137.37	7,6	149.15	7,3	143.26
Mean	6.61	129.77	7.18	140.93	6.82	133.94




Figure 2. Sediment height distribution in each station from March to September 2012.

B. Suspended Solid (TSS)

Table 2 shows that total suspended solid (TSS) in station 1 ranges from 18.28 to 50.60 ppm with an average of 30.40 ppm (29.59%) (Table 2, Figure 3 A), in station 2 from 27.84 to 47.48 ppm with an average of 38.99 ppm (39.56%) (Table 2, Figure 3 B) and in station 3 from 18.28 to 44.68 ppm with an average of 30.405 ppm (30.85%), respectively (Table 2, Figure 3 C). These values have exceeded the standard seawater quality based upon the decree of Living Environment Minister numbered 51, 2004, for marine biota < 20 ppm.


1 *Sedimentation Rate and Total Suspended Solid (Tss) In Melombo Area, Salurang Village, Sangihe*

7 Table 2. Calculation table and figure of TSS mean distribution in station 1, 2, and 3,

Station 1				Station 2					
No.	9 filter Weight (mg)	Wet Weight (mg)	Dry Weight (mg)	TSS/ltr	No.	4 filter Weight (mg)	Wet Weight (mg)	Dry Weight (mg)	TSS/ltr
1	0.4574	6.89	6.96	27.84	1	0.4673	9.89	6.96	27.84
2	0.4574	14.32	12.65	50.6	2	0.4764	14.32	11.62	46.48
3	0.4639	9.2	6.82	27.28	3	0.4628	11.37	8.82	35.28
4	0.4819	10.42	9.24	36.96	4	0.4729	10.42	11.35	45.4
5	0.4615	10.05	9.75	39	5	0.4645	10.05	7.85	31.4
6	0.4562	11.45	8.81	35.24	6	0.4548	11.45	9.81	39.24
7	0.4632	13.51	11.17	44.68	7	0.4612	13.51	11.17	44.68
8	0.4553	10.44	8.42	33.68	8	0.4567	10.44	8.42	33.68
9	0.4768	6.92	5.54	22.16	9	0.4565	13.92	10.55	42.2
10	0.4774	6.89	4.57	18.28	10	0.4784	14.36	11.21	44.84
11	0.5747	6.79	4.62	18.48	11	0.5749	15.24	11.87	47.48
12	0.4672	9.87	6.42	25.68	12	0.4589	12.42	9.78	39.12
13	0.4572	9.59	6.32	25.28	13	0.4789	11.78	9.57	38.28
14	0.4642	8.87	6.57	26.28	14	0.4539	10.34	9.07	36.28
15	0.4658	9.17	5.93	23.72	15	0.4563	11.23	8.94	35.76
16	0.4774	10.23	7.83	31.32	16	0.4592	11.12	8.97	35.88
Mean				30.405	Mean				38.99

Station 3

No.	4 filter Weight (mg)	Wet Weight (mg)	Dry Weight (mg)	TSS/ltr
1	0.4575	7.23	4.96	19.84
2	0.4594	8.26	7.23	28.92
3	0.463	9.20	6.72	26.88
4	0.4829	10.42	10.15	40.6
5	0.4615	10.05	9.75	39
6	0.456	8.44	5.81	23.24
7	0.4602	13.51	11.17	44.68
8	0.4563	9.57	8.32	33.28
9	0.4765	6.92	7.1	28.4
10	0.4774	6.89	4.57	18.28
11	0.5746	6.79	4.68	18.72
12	0.4579	9.87	6.42	25.68
13	0.4759	11.56	9.75	39
14	0.4562	9.98	9.43	37.72
15	0.457	7.76	5.23	20.92
16	0.4678	7.3	5.37	21.48
Mean				29.165






Figure 3. Mean TSS distribution in each station from March to September 2012.

Figure 3 exhibits that TSS distribution in each station is relatively higher than that near the river mouth, so that the area easily turns turbid when flooding and big wave occurrence. According to Manik (2003), suspended solids are derived from erosion in terrestrial area, such as residence, agriculture, mining activity and river mouth. Suspended solids are cause of water turbidity, insoluble and cannot directly sink to the bottom (Fardiaz, 1992). On the bottom, it will gradually cover the benthic organisms and could affect the food web (Canter and Hill, 1979). Very high concentration of suspended solids could inhibit light penetration into the water column and decrease the photosynthetic process in the water (Effendi, 2003). It could also disturb fish activities in the area.

Based on climate classification of Schmidt and Ferguson (1951) and 10 year (2002-2011) rainfall data collected from Geophysical and Meteorological Center station in Naha (Bappeda, 2011), Sangihe Island belongs to type A climate with $Q = 7.1\%$, meaning that *melombo* area is relatively wet with relatively high rainfalls. Meanwhile, during this period maximum rainfalls were 3.770 mm, minimum 1.448 mm and the average 2.521 mm. Beside that, based on humidity and air temperature data, it was apparent that the relative humidity was 83.52%, while the average air temperature was 27.39°C.

Looking at the biogeophysical elements mentioned above, there are tendency of reduction in forest area and relatively high rainfalls along the year in Salurang watershed, which is supported by topographic conditions, mostly way to hilly, labile latosol and aluvial soil vulnerable to erosion, and dendritic patterned river fast flowing the water that could easily cause erosion and sedimentation *melombo area*. Land use is any form of human intervention over the land to fulfill their living needs. Each type of land use will have impacts on environmental damages, such as critical land occurrence, flood and drought. Thus, land management without considering the land carrying capacity and suitability will result in many losses, as a result of increasingly higher erosion so that the equilibrium of natural resources could be disturbed (Sitorus, 2002). The bigger the erosion the lower the soil productivity will be.

There are two events showing the sedimentation in *melombo* area. First, the sink of umbrella rock in the estuary (around station 2) (Figure 4), in which in 1970s to the end of 1980s, it was still clearly seen at the depth of 5 – 10 m. The present study, based on bathymetry, the depth ranged between 1 – 5 m (Appendix 1) proving that




Fig. 4. Umbrella Rock

the umbrella rock has mostly been covered by sediment and the upper part is left (Manansang, 2012). Second, seagrass ecosystem behind the reef flat was lost and most corals on the reef flat disappeared due to sediment cover. In low tide and wavy condition, sediments will be reworked by wave and current so that the sediment is lifted up to the water column and eventually fell in *melombo* area. Since the coral reef condition was found badly damaged, it reflects that the destruction of surrounding coral reefs will still occur due to human activities or natural mortality from high sedimentation. According to DKP-COREMAP (2004), the coral reef getting stressed from environmental pressures tends to release mucus which is then attached with sediment particles.

Coral coverage condition was badly damaged indicating that coral degradation has occurred from sedimentation, flow opening for fisherman's boat and natural mortality. The coral damages are mostly caused by human activities, such as illegal gold mining so that it is feared that it could suffer from more damages if there is no management effort involving all stakeholders. Damages from sedimentation could be attributed with widespread dead corals in several locations. It occurs in *melombo* area and needs more attention from the communities, government and private sectors to avoid bigger coral damages around the area.

Ocean circulation was also a major cause behind the global climate change, and significantly influenced the terrestrial climate. Seasonal change in sunlight and ocean temperature was a reflection of coral environmental conditions, including the equator and a parameter of changes in body tissue system of the coral. Other damage type is derived from long exposure to the air and the sunlight at low tide, usually at the reef peak. Storms could affect the coral life and break the reefs and promote sedimentation in large scale seas well (Barnes & Hughes, 1990).

The more the suspended solid carried in the water, the higher the mud sediment in the estuary. Nybakken (1992) stated that sediment formation was also affected by seawater, since it contained enough suspended materials. In this study, forest degradation caused by illegal mining activities and illegal logging in the upper watershed brought about a lot of forest areas turn to gold mining, plantation and agricultural areas. This condition has caused high erosion rate in the upstream impacting on high sedimentation in the downstream.

Water visibility ranged from 0.2 – 6 m in station 1 with an average of 1.79 m, 0.5 – 3.0 m in station 2 with an average of 1.21 m, and 1.5 – 10 m in station 3 with an average of 3.13 m, respectively (Table 3). Average visibility of three stations ranged from 1.2 – 3.13 m indicating that water visibility was below the standard seawater quality for marine biota established by the decree of Living Environment Minister numbered 51, 2004 > 5 m. Water quality parameters of *melombo* area are in general good, and therefore could support the coral reef growth. Coral growth needs a depth range of 3 – 10 m where water depth influences corals in relation to light intensity (Thamrin, 2003 in Kasmir, 2011). All hermatyphic corals require enough light for photosynthesis. These are corals capable of constructing the reefs. They live symbiotically with algae (zooxanthellae) conducting photosynthetic process. Zooxanthellae are autotrophic macroalgae belonging to dinoflagellates. Sunlight penetration highly determines the depth of coral reef habitats (Thamrin, 2006 in Kasmir, 2011).

Low light penetration is usually brought about by high suspended particles-containing riverwater entering the sea (Lalamentik, 2001). Turbidity and sedimentation of *melombo* area have been beyond the tolerable limit for corals. Present study showed visibility range of 1.21 – 3.13 m. Sedimentation and turbidity are highly potential to occur in small islands near the mainland, so that they could be a threat to the coral reef ecosystem. These could put impacts on coral growth and morphology. Rich nutrient-induced turbidity in the water will threaten the corals and even be able to kill the corals, and promote algal growth, so that it could result in coral-algae competition involving a number of interactions. Coral community replacement dominated by macroalgae often indicates external disturbances, not only because of overgrowth competition, but recruitment competitive inhibition as well, so that the coral reef recovery is inhibited. The recovery could occur through coral recruitment in the coastal area of Salurang.

Factors affecting the sedimentation process in the rivermouth are waves and current patterns. Salurang river brought sediment materials and wastes from the upstream and along the watershed that will be deposited in the rivermouth. According to Triatmodjo and Takeda (1993), sedimentation process is affected by tides, currents

and waves. Wave energy functioning as longshore current producer could also cause abrasion. Sedimentation and erosion are two interchanging processes in relatively short time in order to achieve an equilibrium and part of river flow dynamics. Hence, fisheries resources utilization particularly in the estuary requires wise actions oriented in optimal and sustainable utilization. The exploitation patterns should also consider the environmental carrying capacity.

Table 3. Visibility measurement in *melombo* waters from March to September 2012

Measurement	Station		
	03°28'33.6"N 125°40'00.6"S	03°28'27.4"N 125°39'57.5"S	03°28'30.3"N 125°39'58.4"S
March	1.5	1.0	3.8
	1.0	0.8	2.8
	1.0	0.5	2.5
April	1.5	1.0	2.8
	1.5	1.0	2.8
May	2.0	0.6	1.5
	1.8	0.6	1.5
June	1.5	1.0	2.8
	1.5	1.0	2.5
	1.8	1.0	2.5
July	6.0	3.5	10.0
	2.5	1.5	3.0
August	4.2	1.0	0.2
	2.6	1.5	3.5
September	1.2	1.35	3.0
	2.0	1.25	3.75
Mean	2.03	1.21	3.13

IV. Conclusion

Basic problems in Melombo area were erosion and sedimentation from illegal mining activities in Salurang watershed. People's activities were also major cause of high sedimentation rate in Melombo. In the last 8 months, sedimentation rate ranged between 6.61 cm (129.7 cm³) and 7.18 cm (140.9 cm³). Total suspended solid (TSS) at each observation site has reached the seawater quality standard of <20 ppm. Thus, it is necessary to have more comprehensive policy of the local government concerning land management and land use, particularly the watershed areas in Sangihe Archipelagic Regency.


REFERENCE

- [1.] Bappeda, 2011, Profil Kabupaten Kepulauan Sangihe.
- [2.] Barnes, R. S. K. and Hughes, R. N. 1999. An Introduction to Marine Ecology. 3th ed. Blackwell Science. United Kingdom. 286 hal.
- [3.] Canter, W.L. and L.G. Hill. 1979. Handbooks of Variable or Environmental Impact Assessment. Ann Arbor Sci. Pub. Inc. USA.
- [4.] Dahuri, R., Rais, J., Ginting, S.P., and Sitepu, M.J. 2001. Pengelolaan Sumberdaya Pesisir dan Lautan Secara Terpadu (Integrated Coastal and Marine Resource Management). PT. Paradnya Paramita, Jakarta.
- [5.] DKP-COREMAP. 2004. Pedoman Umum Pengelolaan Terumbu Karang Departemen Kelautan dan Perikanan-Coral Reef Rehabilitation and Management Program. Jakarta.
- [6.] Eaton, A.D., L.S. Clesceri, and A.E. Greenberg, 1995. Standard Methods for the Examination of Water and Wastewater. 19th Edition.
- [7.] Effendi, H. 2003. Telaah Kualitas Air Bagi Pengelolaan Sumber Daya dan Lingkungan Perairan, Kanisius, Yogyakarta.
- [8.] Efriyeldi, 1999. Sebaran Spasial Karakteristik Sedimen dan Kualitas Air Muara Sungai Bantan Tengah Bengkalis dengan Budidaya Keramba Jaring Apung Fakultas Perikanan Universitas Riau. Eidman. Gramedia Jakarta.
- [9.] Fardiaz, S. 1992. Polusi air dan Udara. Kerjasama Pusat Antar Universitas Pangan dan Gizi Institut Pertanian Bogor. Penerbit Kanisius.
- [10.] Kasmir M., 2011. Analisis Aspek Ekologi Penata Kelolaan Minawisata Bahari Kepulauan Spermonde Kabupaten Pangkep, Sulawesi Selatan. *Fakultas Perikanan dan Ilmu Kelautan UMI, Jurnal Ilmu Kelautan. Vol. 16 (2) 61-69 ISSN 0853-7291.*
- [11.] Lalamentik, L.T.X., 2001. Coral Reef Conditions Around the Gold Mining Area of PT. Newmont Minahasa Raya A Monitoring Study in Rataotok and Adjacent Waters, The District of Minahasa North Sulawesi. (Survey Dated September, 2001) Sam Ratulangi University, Manado.
- [12.] Manansang 2012. Mantan ketua Adat (Perosanal Communication).
- [13.] Manengkey H, W 2003. Tingkat Sedimentasi dan Pengaruhnya pada Ekosistem Terumbu Karang di Perairan Teluk Buyat dan sekitarnya Provinsi Sulawesi Utara.
- [14.] Manik, K.E.S. 2003. Pengelolaan Lingkungan Hidup. Djambatan. Jakarta.
- [15.] Menteri Lingkungan Hidup No. 51 tahun 2004. Tentang Baku Mutu Air Laut. Kantor Menteri Negara Lingkungan Hidup, Jakarta : 32 hal.

Sedimentation Rate and Total Suspended Solid (Tss) In Melombo Area, Salurang Village, Sangihe

- [16.] Nybakken, J.W. 1992. Biologi Laut. Suatu Pendekatan Ekologis. Gramedia, Jakarta. Penerjemah: Eidman dkk. 459 Hal.
- [17.] ODUM, E.P. 1971. *Fundamental of Ecology*. W.B. Saunder Com. Philadelphia 125 pp.
- [18.] Pickard G.L., 1975. *Descriptive physical oceanography. An Introduction*, Pergamon Press, Oxford, p.214.
- [19.] Pickard, G.L., and W.J Emery 1982. *Decriptive Physical Oceanography. An Introduction* . Pergamon Press. Oxford.
- [20.] Satriadi, A, 2004, Analisis Oseanografi Terhadap Distribusi Sedimen di Muara Sungai Grindulu Kabupaten Pacitan Jawa Timur, Laporan Penelitian, Surabaya.
- [21.] Schmidt, F.H. and J.H.A. Ferguson, 1951. Rainfall types based on wet and dry period ratios for Indonesia with Western New Guinea. Kementerian perhubungan, Djawatan Meteorologi dan Geofisika, Jakarta.
- [22.] Sitorus, S.R.P., 2002. Evaluasi sumberdaya Lahan. Penerbit Tarsinto Bandung.
- [23.] Triatmojo, B. 1999. Teknik Pantai Edisi Kedua. Beta Offset. Yogyakarta.
- [24.] Umi M dan S. Agus., 2002. Pengantar Kimia dan Sedimen Dasar Laut. Badan Riset Kelautan Dan Perikanan: Jakarta.

Appendix 1. Bathymetric data



Sedimentation Rate and Total Suspended Solid (TSS) In Melombo Area, Salurang Village, Sangihe Archipelagic Regency

ORIGINALITY REPORT

5%

SIMILARITY INDEX

4%

INTERNET SOURCES

3%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	es.slideshare.net Internet Source	2%
2	archive.org Internet Source	<1%
3	www.tandfonline.com Internet Source	<1%
4	Hylka, V.W.. "Ontogeny of embryonic chicken lung: Effects of pituitary gland, corticosterone, and other hormones upon pulmonary growth and synthesis of surfactant phospholipids", <i>General and Comparative Endocrinology</i> , 198310 Publication	<1%
5	Submitted to Heriot-Watt University Student Paper	<1%
6	ejournal.unsrat.ac.id Internet Source	<1%
7	dx.doi.org Internet Source	<1%

8

Dede Aulia Rahman, Georges Gonzalez, Stéphane Aulagnier. " Benefit of camera trapping for surveying the critically endangered Bawean deer (Temminck, 1836) ", *Tropical Zoology*, 2016

Publication

<1 %

9

Pablo Wappner, Luis A. Quesada-Allué. "Water loss during cuticle sclerotization in the medfly *Ceratitis capitata* is independent of catecholamines", *Journal of Insect Physiology*, 1996

Publication

<1 %

10

S Sucahyanto, C Setiawan, R Septiani. "Land Cover Changes and Its Effect to Sediment Growth in Cacaban Reservoir, Tegal District, Central Java Province, Indonesia", *IOP Conference Series: Earth and Environmental Science*, 2018

Publication

<1 %

11

acikerisim.pau.edu.tr:8080

Internet Source

<1 %

Exclude quotes OnExclude matches OffExclude bibliography On