

Dear Muhammad Rahimi Bahar, Fahrul Nurkolis, Jeanette Irene Christiene Manoppo, Nelly Mayulu, William Ben Gunawan, Siti Chairiyah Batubara, Piko Satria Augusta, Nurpudji Astuti Taslim, Max F. J. Mantik,

We have reached a decision regarding your submission to *Annals of the Romanian Society for Cell Biology*, "Cookies High in Antioxidants from the Combination of *Lactobacillus Reuteri* with Mango Juice as a Functional Food Candidate for Children with Functional Digestive Problems", Manuscript ID = ARSCB_0235.

Our decision is to: **Accept your manuscript for publication in ARSCB after minor revision.**

*Please the Authors provide the Author Copyright Transfer Form

Reviewer A:

Manuscript "Cookies High in Antioxidants from the Combination of *Lactobacillus Reuteri* with Mango Juice as a Functional Food Candidate for Children with Functional Digestive Problems" by Bahar et al aimed to process mango juice with *L. reuteri* into functional biscuits. It is very attractive and composed in a suitable method. Innovations that have been researched can be further developed at the mouse/animal trial stage. Please state the bacterial strain used and provide data on the standard deviation of each observed data

Reviewer B:

Please provide additional appropriate references; In substance it is acceptable.

Editor in Chief: Ryon Oelen, Wageningen University & Research, Romania

editor@annalsofscb.ro

Annals of the Romanian Society for Cell Biology

ISSN: 1583-6258

← Balas

→ Teruskan

Annals of the Romanian Society for Cell Biology

ISSN : 1583-6258

mobile : +91 98840 78453 / whatsapp : +91 98840 78453

website : www.annalsofrscb.ro/index.php/journal

Acceptance Letter

Dear Author(s): Muhammad Rahimi Bahar¹, Fahrul Nurkolis², Jeanette Irene Christiene Manoppo¹, Nelly Mayulu³, William Ben Gunawan⁴, Siti Chairiyah Batubara⁵, Piko Satria Augusta⁶, Nurpudji Astuti Taslim⁷, Max F. J. Mantik¹

Paper ID	ARSCB_0235
Paper Title	Cookies High IN Antioxidants from the Combination of Lactobacillus Reuteri with Mango Juice as a functional food candidate for children with functional digestive problems

This is to enlighten you that the above manuscript is reviewed and appraised by the review committee members of BioLEAGUES and it is accepted for the purpose of publication in “**Annals of the Romanian Society for Cell Biology ISSN: 1583-6258**”.

Copyright Form

Journal Title: Annals of Romanian Society for Cell Biology

Title of the Paper: COOKIES HIGH IN ANTIOXIDANTS FROM THE COMBINATION OF LACTOBACILLUS REUTERI WITH MANGO JUICE AS A FUNCTIONAL FOOD CANDIDATE FOR CHILDREN WITH FUNCTIONAL DIGESTIVE PROBLEMS

Corresponding Author: Muhammad Rahimi Bahar

Affiliation: Department of Pediatric, Faculty of Medicine, Sam Ratulangi University, Indonesia

Address: Manado, Indonesia

E-mail: rahimibahar17@gmail.com Tel: + +62 822-5566-2007

Terms & Conditions

1. The copyright transfer covers the exclusive right to reproduce and distribute the contribution, including reprints, translations, photographic reproductions, microform, electronic form, or any other reproductions of similar nature.
2. The Author may not publish his/her contribution anywhere else without the prior written permission of the publisher unless it has been changed substantially.
3. The Author promise that the article is original work, has not previously been published, and is not currently under consideration by another publication.
4. The Author warrants that his/her contribution is original, except for such excerpts from copyrighted works as may be included with the permission of the copyright holder and author

10. No authors will be added or removed after the submission of the article.
11. The corresponding author also declares that on the acceptance of the manuscript s/he is responsible for the payment of the publication fee in those journals that carry a required fee after the acceptance.
12. After signing the copyright form, withdrawn of Article from the respected Journal is not acceptable and Article Processing Fee is not refundable.
13. Journal have the right to reject the article and BIOLEAGUES is not involved in the decision making process.
14. BIOLEAGUES do not involve in any process of enlisting the articles into the indexed databases like SCOPUS, PubMed, Web of Science, UGC, etc.
15. BIOLEAGUES do not involve in any selection of the Journal while publishing the article. BIOLEAGUES will not force any authors to publish their articles in any particular journal.
16. For any dispute or related violence, it will be discussed and considered only in front of 'Judiciary of Chennai' at Chennai, Tamil Nadu, India.

I have read and agreed to the above Terms and conditions

Date: 13 April 2021

Muhammad R. Bahar

Corresponding Author Signature

Cookies High in Antioxidants from the Combination of *Lactobacillus Reuteri* with Mango Juice as a Functional Food Candidate for Children with Functional Digestive Problems

Muhammad Rahimi Bahar¹, Fahrul Nurkolis², Jeanette Irene Christiene Manoppo³, Nelly Mayulu⁴, William Ben Gunawan⁵, Siti Chairiyah Batubara⁶, Piko Satria Augusta⁷, Nurpudji Astuti Taslim⁸, Max F. J. Mantik⁹

^{1,3,9}Pediatrics, Faculty of Medicine, Sam Ratulangi University, Manado, Indonesia.

²Biological Sciences, Faculty of Sciences and Technology, State Islamic University of Sunan Kalijaga, Yogyakarta, Indonesia.

⁴Nutrition, Faculty of Medicine, Sam Ratulangi University, Manado, Indonesia.

⁵Nutrition, Faculty of Medicine, Diponegoro University, Semarang, Indonesia.

⁶Food Technology, Faculty of Food Technology and Health, Sahid University of Jakarta, South Jakarta, Indonesia.

⁷Medical Sciences, Faculty of Medicine, Brawijaya University, Malang, Indonesia.

⁸Professor Clinical Nutrition, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia.

¹Corresponding Author: rahimibahar17@gmail.com

ABSTRACT

Lactobacillus reuteri is one of the good microorganisms that play a role in the digestive health of children, such as diarrhea and constipation. Mango fruit (*Mangifera indica*) has an abundant content of antioxidants, dietary fiber, and beta-carotene. This research aims to process mango juice with *L. reuteri* into functional biscuits. There were 3 variations of mango juice formulation, namely S1, S2, and S3, based on mango and CO₂-free water content (S1 = 1:0.5; S2 = 2:1; and S3 = 3:2). Then, all product samples were inoculated with *Lactobacillus reuteri* LRE02-DSM 23878 5% b/v for 14 days under anaerobic conditions. The part of mangoes used was pure ripe flesh. The fermented products were made into flour with a freeze dryer, then the powder was mixed and stirred with the addition of 5% water using a mixer with a power of 102-189 rpm for 30 minutes and then put in the oven for 15 minutes with a temperature of 70 - 90°C so that it became cookies. Sample variation was done to determine the average significance of the antioxidant content in it. In the next step, Vitamin C analysis from 3 samples of cookies was done using the Titration Iodometric Method to determine the amount of Vitamin C (mg/100g) and also the antioxidant activity with 2,2-diphenyl-1-picrylhydrazyl (DPPH). The results of vitamin C and antioxidant activity in each sample of cookies; S1 was 89.64mg/100g with 25.78% antioxidant activity, S2 was 81.55mg/100g with 23.77% of antioxidant activity, and S3 was 96.92mg/100g with 28.67% antioxidant activity. The formulation for the number of cookie samples containing the highest vitamin C was S3. There was a significant difference ($P < 0.05$) which determined the vitamin C level between the sample formulations..

Keywords:

Cookies; antioxidants; *L. reuteri*; functional food; digestive problems.

1.Introduction

Mango (*Mangifera indica*) is a tropical fruit plant that contains high levels of nutrients, fiber, macronutrients, micronutrients, and minerals as well as abundant bioactive compounds (Maldonado-Celis *et al.*, 2019). The well-known high content in mangoes, including vitamin C, beta-carotene, polyphenol types of quercetin, and kaempferol (Nurkolis *et al.*, 2020; Mantik *et al.*, 2021). Recent research has shown that increasing the intake of foods high in antioxidants and polyphenols such as Vitamin C, beta-carotene, quercetin and kaempferol can increase the body immunity against viral infection (Levy *et al.*, 2020; Pitsillou *et al.*, 2020; Suhail *et al.*, 2020). Therefore, the consumption of food sources of antioxidants can be one of the efforts to minimize inflammation, one of which is inflammation caused by functional digestive diseases. Epidemiological studies have shown that approximately 14-25% of infants, young children, and

adolescents suffer from at least one gastrointestinal function disorder (Robin, et al.,2018). Probiotics are living microorganisms that, if administered in sufficient quantities, can provide health benefits to the host by regulating an unbalanced gastroenteric microbiota (Margiotta, et al.,2021). One of those probiotics is *Lactobacillus reuteri* LRE02-DSM 23878. *Lactobacillus reuteri* (*L. reuteri*) is a well-studied probiotic bacteria that can colonize a large number of mammals (Mu, Tavella and Luo, 2018). In humans, *L. reuteri* is found in different body sites, including the gastrointestinal tract, urinary tract, skin, and breast milk. The abundance of *L. reuteri* varies among different individuals (Mu, Tavella and Luo, 2018). Some beneficial effects of *L. reuteri* have been noted. First, *L. reuteri* can produce antimicrobial molecules, such as organic acids, ethanol, and reuterin (Mu, Tavella and Luo, 2018). Due to its antimicrobial activity, *L. reuteri* can inhibit the colonization of pathogenic microbes and overhaul the composition of the commensal microbiota inside the host. Second, *L. reuteri* can be beneficial for the immune system. For example, some strains of *L. reuteri* may reduce the production of pro-inflammatory cytokines while promoting the development and function of regulatory T cells. Third, with the ability to strengthen the intestinal barrier, colonization of *L. reuteri* can decrease the translocation of microbes from the intestinal lumen to the tissues. Microbial translocation across the intestinal epithelium has been hypothesized as an inflammatory trigger (Mu, Tavella and Luo, 2018). Therefore, inflammatory diseases, including those located in the intestines as well as in distant tissues, can be corrected by increasing the colonization of *L. reuteri*. In particular, the decrease in the abundance of *L. reuteri* in humans in recent decades correlated with an increase in the incidence of inflammatory diseases over the same period. Direct supplementation or modulation of *L. reuteri* prebiotics may be an interesting means of prevention and/or therapy against inflammatory diseases (Mu, Tavella and Luo, 2018). This study aims to process mango juice fermented by *L. reuteri* into cookies that are high in antioxidants as a functional food for children with functional digestive problems.

2. Material and Methods

There were 3 variations of mango juice formulation, namely S1, S2, and S3, based on mango and CO₂-free water content (S1 = 1:0.5; S2 = 2:1; and S3 = 3:2). Then, all product samples were inoculated with *Lactobacillus reuteri* LRE02-DSM 23878 5% b/v for 14 days under anaerobic conditions. The part of mangoes used was pure ripe flesh. The fermented products were made into flour with a freeze dryer, then the powder was mixed and stirred with the addition of 5% water using a mixer with a power of 102-189 rpm for 30 minutes and then put in the oven for 15 minutes with a temperature of 70 - 90°C so that it became cookies. Sample variation was done to determine the average significance of the antioxidant content in it. In the next step, Vitamin C analysis from 3 samples of cookies was done using the Titration Iodometric Method to determine the amount of Vitamin C (mg/100g) and also the antioxidant activity with 2,2-diphenyl-1-picrylhydrazyl (DPPH).

$$\text{Vitamin C} \left(\frac{\text{mg}}{100\text{g}} \right) = \frac{V I 2 \times 0,88 \times Fp \times 100}{Ws \text{ gram}}$$

The determination of water content used was the AOAC drying method (thermogravimetry). The principle of this method is based on the evaporation of water in the material by heating, then weighing it to a constant weight. The weight reduction that occurs is the water content contained in the material. An empty plate is heated in an oven at 105° C for 30 minutes, cooled down in a desiccator for 15 minutes, then weighed (W0). A 2-gram sample was then put in a plate with known weight, weighed (W1), then dried in an oven at 105° C for 3 hours, cooled in a desiccator

for 15-30 minutes, then the plates and contents were weighed and dried again for another one hour, cooled in the exicator, and weighed again (W2). The water content was calculated using the following formula:

$$\text{Water Content (\%)} = \frac{W1 - W2}{W1 - W0} \times 100$$

The procedure for determining the ash content was carried out using the AOAC 2005 method. Meanwhile, the procedure is as follows: the cup was dried in an oven at 105° C for 1 hour. Then, it was cooled in the cup for 15 minutes in a desiccator and weighed. 2 grams of the sample was put into a furnace where the temperature was 550° C for 3 hours. It was then cooled outside the furnace to a temperature of ± 120° C and put in a desiccator. The plates and ashes were weighed so that a constant weight was obtained. Calculation of the ash content was carried out using the following formula:

$$\text{Ash Content (\%)} = \frac{\text{Weight of Bowl After Heated} - \text{Constant Weight of Empty Bowl}}{\text{Sample Weight}} \times 100$$

3.Results and Discussion

The results of vitamin C and antioxidant activity in each sample of cookies; S1 was 89.64mg/100g with 25.78% antioxidant activity, S2 was 81.55mg/100g with 23.77% of antioxidant activity, and S3 was 96.92mg/100g with 28.67% antioxidant activity. The formulation for the number of cookie samples containing the highest vitamin C was S3. There was a significant difference (P <0.05) which determined the vitamin C level between the sample formulations. The higher the antioxidant activity, the higher the antioxidant levels, and the less food needed to reduce free radicals (Lisdawati and Kardono., 2012). This shows that the formulation of mango juice which was made into cookies contains vitamin C and has antioxidant activity. This makes formulated cookies a potential healthy snack that is high in vitamin C and antioxidants during the COVID-19 pandemic. The average vitamin C level in the three cookie samples was 89.37mg/100g. S3 showed the best activity, namely antioxidant activity against 2,2-diphenyl-1-picrylhydrazyl (DPPH) of 28.67%.

The average ash content of the three samples was 1.85% and water content was 1.87%, which corresponds to the Indonesian National Standard (SNI) 01-2973-1992. Higher ash content in cookies indicates the higher mineral contents in cookies such as calcium, potassium, and iron (Andarwulan *et al.*, 2014). It is clinically known that mineral intake can improve the respiratory system, especially in tuberculosis sufferers (Taslim *et al.*, 2020).

Fermented mango with *L. reuteri* has a great potential to be developed into healthy snack cookies. The vitamin C and antioxidants content in cookies from the fermentation of mango may be a great substitute for snacks for children with functional digestive problems since antioxidants and vitamin C can improve immunity and anti-inflammatory response. These cookies are also good prebiotics for the gut microbiome which plays a good role in the digestive system and immune system. It needs clinical trials in humans to find out more about its effects on human health and the authors are very open to joint research collaborations.

Acknowledgments

All authors contributed to the writing and revisions contained in the paper; and all authors have read and approved the final manuscript.

Conflict of Interest

The authors declare that there are no conflicts of interest

References

- [1] Andarwulan, N., Kusnandar, F. and Herawati, D., (2014). Pengelolaan Dara Analisis Pangan. Modul I.
- [2] Levy, E., Delvin, E., Marcil, V. and Spahis, S. (2020). Can phytotherapy with polyphenols serve as a powerful approach for the prevention and therapy tool of novel coronavirus disease 2019 (COVID-19)? In American Journal of Physiology - Endocrinology and Metabolism (Vol. 319, Issue 4, pp. E689–E708). American Physiological Society. <https://doi.org/10.1152/ajpendo.00298.2020>
- [3] Lisdawati, V. and Kardono, L. B. S. (2006). Aktivitas Antioksidan Dari Berbagai Fraksi Ekstrak Daging Buah dan Kulit Biji Mahkota Dewa (*Phaleria macrocarpa*). Media Penelitian dan Pengembangan Kesehatan, 16(4), 161424.
- [4] Maldonado-Celis, M. E., Yahia, E. M., Bedoya, R., Landázuri, P., Loango, N., Aguillón, J., Restrepo, B. and Guerrero Ospina, J. C. (2019). Chemical Composition of Mango (*Mangifera indica* L.) Fruit: Nutritional and Phytochemical Compounds. *Frontiers in Plant Science*, 10. <https://doi.org/10.3389/fpls.2019.01073>
- [5] Mantik, K.E.K., Nurkolis, F., Mayulu, N., Kurniatanty, I., Widowati, E., Tanner, M.J. and Hardinsyah. (2021) 'Functional sugar from mango (*Mangifera indica*) rich in antioxidants and polyphenols potential for antidiabetic', *Annals of the Romanian Society for Cell Biology*, 25(1), pp. 2545–2553. Available at: <http://annalsofrscb.ro> (Accessed: 9 March 2021).
- [6] Margiotta G, Ferretti S, Graglia B, Gatto A, Capossela L, Bersani G, Curatola A, Chiaretti A. Effect of *Lactobacillus reuteri* LRE02-*Lactobacillus rhamnosus* LR04 combination and gastrointestinal functional disorders in an Emergency Department pediatric population. *Eur Rev Med Pharmacol Sci*. 2021 Apr;25(7):3097-3104. doi: 10.26355/eurev_202104_25564. PMID: 33877678.
- [7] Mu, Q., Tavella, V. J., & Luo, X. M. (2018). Role of *Lactobacillus reuteri* in Human Health and Diseases. *Frontiers in microbiology*, 9, 757. <https://doi.org/10.3389/fmicb.2018.00757>
- [8] Nurkolis, F., Surbakti, F. H., Sabrina, N., Azni, I. N. and Hardinsyah, H. (2020). Mango Sugar Rich in Vitamin C: A Potency for Developing Functional Sugar Rich in Antioxidants. *Current Developments in Nutrition*, 4(Supplement_2), 765–765. https://doi.org/10.1093/cdn/nzaa052_034
- [9] Pitsillou, E., Liang, J., Ververis, K., Lim, K. W., Hung, A. and Karagiannis, T. C. (2020). Identification of Small Molecule Inhibitors of the Deubiquitinating Activity of the SARS-CoV-2 Papain-Like Protease: in silico Molecular Docking Studies and in vitro Enzymatic Activity Assay. *Frontiers in Chemistry*, 8, 623971. <https://doi.org/10.3389/fchem.2020.623971>
- [10] Robin SG, Keller C, Zwiener R, Hyman PE, Nurko S, Saps M, Di Lorenzo C, Shulman RJ, Hyams JS, Palsson O, van Tilburg MAL. Prevalence of pediatric functional

gastrointestinal disorders utilizing the Rome IV criteria. *J Pediatr* 2018; 195: 134-139

- [11] Suhail, S., Zajac, J., Fossum, C., Lowater, H., McCracken, C., Severson, N., Laatsch, B., Narkiewicz-Jodko, A., Johnson, B., Liebau, J., Bhattacharyya, S. and Hati, S. (2020). Role of Oxidative Stress on SARS CoV (SARS) and SARS-CoV-2 (COVID-19) Infection: A Review. *The Protein Journal*, 1-13. Springer. <https://doi.org/10.1007/s10930-020-09935-8>
- [12] Taslim, N. A., Rasyid, H., Atmanegara, M. K., Angriavan, S. and Amelia, R. (2020). Effect of Chocolate Soybean Drink on Nutritional Status, Gamma Interferon, Vitamin D, and Calcium in Newly Lung Tuberculosis Patients. *Open Access Macedonian Journal of Medical Sciences*, 8(T2), pp. 210-214. doi: 10.3889/oamjms.2020.5233.

Table 1. Vitamin C Content, Anti-Oxidant Activity, Water and Ash Content in Cookies

Sample	Vitamin C Content (mg/100g)	Anti-Oxidant Activity Towards (DPPH)	Ash Content (%)	Water Content (%)
S1	89.64	25.78%	1.80	1.97
S2	81.55	23.77%	1.86	1.85
S3	96.92	28.67%	1.89	1.79
Mean	89.37 ± 7.68	26.07 % ± 2,46	1.85 ± 0.04	1.87 ± 0.09