

Find messages, documents, photos or people



← Back ↶ ↷ → Archive Move Delete Spam ...

Re: new article submission Yahoo/Inbox



gavriloae ionel claudiu <ionelclaudiu@yahoo.com>
To: Benny Pratasik

Sat, Feb 22, 2020 at 1:45 AM

Dear Professor Pratasik,

I received and already performed the preliminary evaluation of the paper. The editing style was a disaster... So I have done a lot of work doing it properly. However, before sending the paper to reviewers for the fast track review, kindly solve the few problems in the References section. Thank you very much!

Cordially yours,
Claudiu G.

On Tuesday, February 18, 2020, 7:20:08 AM GMT+2, Benny Pratasik <spjong07@yahoo.com> wrote:

Dear Dr. Gavriloae,
I herewith send you a new article "Faviidae Coral Colonization Living and Growing on Agricultural Waste-Materialized Artificial Substrate" (attached). The authors wish to have fast service for the next issue. Thank you for your cooperation.

Sincerely Yours,
Silvester B. Pratasik

[Download all attachments as a zip file](#)

Faviidae Coral Colonization Living and Growing on Agricultural Waste-Materialized Artificial Substrate

Laurentius T. X. Lalamentik, Rene Ch. Kepel, Lawrence J. L. Lumingas, Unstain N. W. J. Rembet, Silvester B. Pratasik, Desy M. H. Mantiri

Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia

Corresponding Author: Laurentius T. X. Lalamentik
email : ottylalamentik@yahoo.co.id

Abstract. A study on colonization of Faviidae corals on the agricultural waste-materialized artificial substrate was conducted in Selat Besar, Rataotok district, southeast Minahasa regency, North Sulawesi. Nine artificial substrates moduls made of mixture of cement, sand, padi husk, and bamboo were placed for about 5 years on the sea bottom of Selat Besar waters. All corals of family Faviidae found on the artificial substrate were collected. Results showed that Faviidae corals could live and develop on those substrates. Fifteen species of 5 genera of Faviidae, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii* *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennetae*, *Echinopora gemmacea*, *E. lammelosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. senensis* were recorded in the present study. Mean number of colonies of Faviidae corals was 3 col.mod⁻¹, while mean diameter of the corals attached on the artificial substrate was 7.45 cm long. The distribution pattern of Faviidae corals was clumped. The diversity of Faviidae corals on the artificial substrate was low ($H' = 0.677 < 1$). The dominance index showed no dominant species ($D = 0.23$). In addition, the artificial substrate modul in this study could become an alternative technique to rehabilitate the degraded coral reefs.

Keywords: diversity, dominance, distribution pattern, Selat Besar

Introduction

Background. Corals are animals belonging to phylum Cnidaria (Reid et al 2009), while coral reef is an ecosystem built by calcium carbonate-producing marine biota, especially coral animals, together with other biota living on the sea bottom or water column, such as mollusks, crustaceans, echinoderms, porifera, tunicates, and other biota free-living in surrounding waters including plankton and fishes (Giyanto et al 2017; Lalamentik 1995). According to Barus (2013), coral reef is one of the unique communities that are entirely formed from biological activities. It is one of the main coastal and marine ecosystems that possess the highest productivity and biodiversity so that it is often called as tropical marine forest.

Coral Triangle (CTI) is known as center of world marine biodiversity with the highest coral diversity, 76% of total coral species. The region inside the ecological boundary of Coral Triangle covers nearly 73,000 km² of coral reefs (29% of total world coral reef area) distributed in six countries, Indonesia, Malaysia, Filipina, Timor Leste, Papua New Guinea and Solomon Islands (Burke et al 2012). As part of the CTI, Indonesia is an archipelagic country with 16,056 islands, 1,922,570 km² terrestrial area and 3,257,483 km² water in the west Indo-Pacific (BIG 2016). Indonesia that is located in the coral triangle area is recognized as one of the countries with high biodiversity consisting of more

than 80 coral genera and 596 species, particularly in North Sulawesi waters that has been recorded more than 80 genera (Suharsono 2008).

Coral taxonomic and distribution studies in North Sulawesi were carried out by several scientists (Lalamentik 1998; Halidu et al 2016; Nasaru et al 2017; Suleman et al 2017; Lalamentik & Rembet 2018). So far, there is no study on corals of family Faviidae living and growing on the artificial substrate, especially in North Sulawesi waters. Hence, present study focuses on Faviidae coral colonies on the artificial substrate with the objectives of identifying the coral species, analysing the density and the diameter of the coral colony, the ecological index, and assessing the distribution patterns. This finding is expected to be one of the references in coral development studies, concerning Faviidae coral colonization on the artificial substrate. Besides, it could provide alternative artificial substrates in coral reef rehabilitation program.

Method

Study site. This work was accomplished in Selat Besar waters, Ratatotok district, southeast Minahasa regency (Figure 1), North Sulawesi. Selat Besar is flanked by Totok bay and Moluccas Sea. The study site was laid on $0^{\circ}51'34.945''S$ and $124^{\circ}44'14.416'' E$ (Figure 1) using Global Position System (GPS).




Figure 1. Sampling locality

Artificial substrate construction. In this study, 9 artificial substrate moduls were set on the sea bottom, approximately 7-8 m depth. Sheehy (1986) stated

that one of the major roles of artificial substrate placement in the water is to create new habitats similar to artificial reefs to be inhabited by target species. The use of artificial substrate as coral new habitat had been conducted in Singaporean waters in order to restore the coral reefs (Ng & Chou 2017). Each modul consists of 10 concrete blocks made of mixture of padi husk, sand, and cement. The block frames were made of bamboo. The use of bamboo in concrete construction has been done before as mentioned in Glenn (1950) and Mehra & Ghosh (1965). One of the properties that would make bamboo a good substitute to steel in reinforced concrete is its strength (Varma, 2017). The block had a dimension of 100 cm leng, 15 cm width, and 15 cm height. The blocks were set in such a manner to yield a modul. Each modul is estimated to have an area of 4.5 m². These 9 moduls were placed since June 2014. Sample collections were carried out in July 2019 (after 5 years of deployment). The period of 5 years is determined because the massive coral growth is slow. This duration is considered enough to see whether there is Faviidae coral capable of adhering and growing on the artificial substrate.

Sample collection and handling procedures. All massive coral species found to attach on the modul substrate were taken using a chisel and hammer. The samples were then put into labelled bag as the modul number. All corals were collected by using SCUBA equipments. All activities were documented by using an underwater camera.

Before species identification, the coral specimens were soaked in bleach-containing freshwater for 24 h in order to kill the coral and to prevent the presence of fungi. The corals were then cleaned under the running water to remove the attached dirt so that the clean coral and the bleaching coral skeleton were obtained. The corals were placed under direct sunlight to get dry to ease the identification.

Species identification. Coral identification was done following Veron (1986; 2000), Suharsono (2008), and Veron et al (1977). The specimens of Faviidae corals were examined through morphological observations under the SZ51 Olympus microscope on the characteristics of paliform lobe, septa cleats, coralite type, number of first septa, second septa, and third septa. The measurements of width of paliform lobe, coralite size, and calix size used a multitoyo digital caliper.

Data analysis. Faviidae coral colonization data were directed to ecological index analyses as follows:

Dominance index. This index was employed to know the dominant species in data sampling boundary. The present study applied Simpson dominance index (Krebs, 1989):

$$D = \sum_{i=1}^s \left[\frac{n_i}{N} \right]^2 \dots\dots\dots (1)$$

Where D = dominance index, n_i = number of *i* individuals, and N = total number of individuals

Dominance index ranges between 0 and 1, if D approaches to 0, it indicates no dominance and the community structure is in stable condition, and if D approaches to 1, it indicates the presence of species dominance.

Density. The population density was calculated using Krebs (1989) as follows

$$K = \frac{N_i}{A} \dots\dots\dots (2)$$

Where, K = density (ind.modul⁻¹), Ni = number of individuals, and A = number of moduls where samples were collected.

Diversity. The diversity index of genus *Favia* Oken 1815 was calculated using Shannon-Wiener equation (Krebs 1989):

$$H = \sum_{t=1}^s \frac{ni}{N} \left(\ln \frac{ni}{N} \right) \dots\dots\dots (3)$$

Where H = Shannon-Wiener diversity index, ni = number of individuals *i*, N = total number of individuals, and s = number of genera.

Distribution pattern. The distribution pattern of Faviidae corals were estimated using Morisita index (Krebs 1989):

$$Id = q \frac{\sum_{i=1}^q ni (ni-1)}{N (N-1)} \dots\dots\dots (4)$$

Where Id = Morisita index, ni = number individuals in each plot, n = total number of individuals in all plots, and N = number of plots, in which Iδ = 1 is categorized as random distribution, Iδ > 1 as clumped, and Iδ < 1 as uniform.

Results and Discussion

Water conditions of Selat Besar. Water quality is crucial for colonization of macroinvertebrates (Orwa et al 2018). Water temperature is one of the limiting factors for aquatic organisms to live, one of which is coral. Field measurements found that water temperature in te study site ranged from 29° - 30°C. This range is still suitable for coral development. According to Kurniawan (2011), corals, in general, can live at the temperature above 18°C, while growth, they require an ideal water temperature, between 25°-32°C. Most coral reefs exist where salinity is stable and where average salinity is that of normal seawater 34-36‰ (Sheppard et al., 2017). Water salinity in the study site was between the range of 30-31‰. Guntur (2011) found that water salinity for ideal coral growth in Indonesia ranged from 29 to 36‰. Therefore, water salinity in Selat Besar highly supports the coral to grow well. Light intensity does not influence larval settlement, and their response to light intensity may be species-specific (Mundy & Babcock 1998).


Corals of family Faviidae recorded on the artificial substrate. Physical and biological factors are important for recruitment success (Doropoulos et al 2016). As many as 15 coral species of 5 genera were recorded in this study. *Favia* and *Favites* had the highest number of species, each with 4 species, followed by genus *Goniastrea* with 3 species, then *Echinopora* and *Platygyra*, 2 species, respectively. Table 02 demonstrates the coral species of Faviidae encountered on the modul substrate with mean number of colonies of 3.00 per modul. Even though these numbers are less than those reported by Kilfoyle et al (2008), 13 col.mod.⁻¹, the colonies comprised several coral families. The coral genera of Faviidae inhabiting the artificial substrate apparently come the coral reef around Selat Besar waters. It is reasonable, since Faviidae is coral family generally found in Selat Besar waters (Lalamentik 1996;1997;1998).

Tabel 2.

Faviidae coral species encountered living on the artificial substrate modul.

No	Species
1	<i>Echinopora gemmacea</i> (Lamarck, 1816)
2	<i>E. lammelosa</i> (Esper, 1795)
3	<i>Favia laxa</i> (Klunzinger, 1879)
4	<i>F. matthai</i> (Vaughan, 1918)
5	<i>F. palida</i> (Dana, 1846)
6	<i>F. stelligera</i> (Dana, 1846)
7	<i>Favites bennetae</i> (First record for Central Indonesia)
8	<i>F. complanata</i> (Ehrenberg, 1734)
9	<i>F. pentagona</i> (Esper, 1794)
10	<i>F. russelli</i> (Wells, 1954)
11	<i>Goniastrea aspera</i> (Verrill, 1864)
12	<i>G. favulus</i> (Dana, 1846)
13	<i>G. pectinata</i> (Ehrenberg, 1834)
14	<i>Platygyra daedalea</i> (Elis & Solander, 1786)
15	<i>P. sinensis</i> (Edwards & Haime, 1857)

Favites bennetae recorded in the present study is the first record found in the coral reef of Central Indonesian waters. This species was previously mostly found in the northeast of Australian waters (Veron et al 1977). Moreover, genus *Favia* was found having the highest mean number of colonies, 1.11 col.mod⁻¹, followed by *Goniastrea*, 0.67 col.mod⁻¹, and the lowest was found in *Platygyra*, 0.22 col.mod⁻¹ (Figure 2). Variations in number of Faviidae colonies per modul are dependent upon the abundance of larvae that survive and settle on the substrate. Faviidae tends to have good adaptability to occupying the available artificial substrates. Bachtiar et al (2011) found that Faviidae is one of the coral families capable of surviving in the reef ball deployed on the sea bottom.

Figure 2. Mean no. colonies of *Faviidae* per modul.

Mean diameter of the coral colony on the artificial substrate was 7.45 cm with the largest recorded in *Favia laxa*, 10.5 cm and the smallest in *F. palida*, 3 cm. In comparison among genera, mean diameter of Faviidae corals attached

on the artificial substrate ranged from 5.88 to 8.50 cm. Genus *Platygyra* has the largest mean diameter, 8.50 cm, and *Echinopora* had the smallest one, 5.88 cm (Figure 2). Variations in colony size could result from different sequence of planular settlements. This condition, according to Bachtar et al (2011), indicates that coral colonization on the artificial substrate sustainably occurs.




Figure 3. Mean diameter of Faviidae corals living on the artificial substrate.

Ecological index. The present study measured dominance index (D) and diversity index (H') of Faviidae corals thriving on the artificial substrate. The data are given in Table 3.

Table 3.

Ecological index of Faviidae corals

No	Ecological Index	Value	Remarks
1	Dominance Index (D)	0.23	No dominance
2	Distribution pattern (I δ)	1.79	Clumped
3	Diversity Index (H')	3.64	high

The dominance index of Faviidae corals is 0.23 (Table 3). It means that no species outnumbers the others. Selat Besar waters is still capable of supporting the coral life, and there is no competition that could make certain species be dominant (Muqsit et al 2016). The diversity index is 3.64. This value is supported by low species dominance index, in which there were 15 Faviidae species found. These numbers are high enough due to small study area cover. This finding is in agreement with Rondo et al (2014) that high diversity will cause low species dominance.

Distribution pattern analysis found that Faviidae in the study site had clumped distribution (I δ = 1.79 > 1). This pattern could result from the fact that there were four moduls not occupied by Faviidae corals. In Indonesia, studies on coral colonization was also done by Razak (2008) using Eco-reef modul in Manado bay waters, North Sulawesi, and found that the artificial substrate used are, in fact, inhabited by hard corals. Hence, the artificial substrate could be recommended as an alternative in coral reef rehabilitation program.

Conclusions. Fifteen coral species were found settling on the agricultural waste-materialized artificial substrate, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennetae*, *Echinopora gemmacea*, *E. lammelosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. senensis*. They had mean 3.00 col.mod⁻¹ with mean diameter of 7.45 cm. The ecological index also indicated that the available artificial substrate could support various coral species to grow. In this study, Faviidae corals growing on the available artificial substrate have reflected clumped distribution. Thus, this type of modul could be used for coral reef rehabilitation.

References

- Bachtiar I., Prayogo W., Sukri A. 2011 Coral colonization and growth reef ball modules at the Benete Bay, Sumbawa Island, Indonesia. Manado. P. 31-40.
- Doropoulos C., Roff G., Bozec Y. M., Zupan M., Werminghausen J., Mumby P. J. 2016 Characterizing the ecological trade-offs throughout the early ontogeny of coral re-cruitment. Ecol. Monogr. 86.
- Guntur. 2011 [Coral ecology in the artificial reef]. Ghalia Indonesia. Semarang. 139 pp. [in Indonesian]
- Halidu, A., Lalamentik L. T. X., Rembet U. N. W. J. 2016 [Hard coral distribution in the reef flat of the south coast of Putus-Putus island, east Ratatotok, Ratatotok district, southeast Minahasa]. Jurnal Ilmiah Platax 4(1):19-30. [in Indonesian]
- Kilfoyle K., Rangel M. A., Dodge R. E., Spieler R. E. 2008 Coral Reef Restoration: Standardized Module Intervention and monitoring program in Mexico, *Preliminary Results*. Poster presentation material, 11 Int Coral Reef Sym Florida, Session number XXIV, p. 541
- Krebs C. J. 1989 Ecological Methodology. Harper Collins Publishers. New York. 654 pp.
- Lalamentik L. Th. X. 1995 [Study on coral reef potential in Tombasian district, Minahasa, North Sulawesi] Fakultas Perikanan dan Ilmu Kelautan UNSRAT. Manado. 28 pp. [in Indonesian] Unpublished
- Lalamentik L. T. X. 1996 Coral Reef conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1996). 23 pp. Unpublished.
- Lalamentik L. T. X. 1997 Coral Reef Conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1997). 22 pp. Unpublished.
- Lalamentik L. Th. X. 1998 Coral reef Condition Arround the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study for May 1998 Data in Ratatotok and Adjacent Waters, The District of Minahasa North Sulawesi. Fakultas Perikanan dan Ilmu Kelautan UNSRAT. Manado. 23 pp. Unpublished.
- Lalamentik, L. Th. X., Rembet, U. N. J. 2018 [Study on the Taxonomy of Genus *Favia* (Oken, 1815) at the Reef Flats of Kampung Ambong Village in Likupang Timur District, North Minahasa]. JURNAL ILMIAH PLATAX, 6(1), 188-193. [in Indonesian]

- Mehra S. R., Ghosh R. G. 1965 "Bamboo-reinforced soil-cement," Civil Engineering and Public Works Review, Vol. 60, no. 711, October; Vol. 60, no. 712. November 1965.
- Mundy, C., Babcock, R., 1998. Role of light intensity and spectral quality in coral settlement: implications for depth-dependent settlement? J. Exp. Mar. Biol. Ecol. 223: 235–255.
- Muqsit A., Purnama D., Ta'alidin Z. 2016 [The community structure of coral reef in Pulau Dua, Enggano district, North Bengkulu regency.] Jurnal Enggano Vol. 1, No. 1 (75-78). Ilmu Kelautan Fakultas Pertanian Universitas Bengkulu. Bengkulu. [in Indonesian]
- Nasaru J. H., Lalamentik L. Th. X., Rembet U. N. W. J. 2017 [Pocillopora verrucosa (Ellis and Solander, 1786) distribution in the reef flat of the south coast of Putus-Putus island, east Rataotok village, southeast Minahasa regency]. Jurnal Ilmiah Platax 5(1): 96-103 pp. Universitas Sam Ratulangi Manado. [in Indonesian]
- Ng C. S. L., Chou L. M. 2017. Coral reef restoration in Singapore-past, present and future. *Sustainability Matters: Environmental Management in the Anthropocene. Singapore: World Scientific*, 3-23 pp.
- Orwa P. O., Omondi R., Chemoiwa E. J. 2018 Colonization patterns of benthic macroinvertebrates in fertilized and non-fertilized earthen fish ponds. Int J Aquac Fish Sci 4(3): 022-026. DOI: <http://doi.org/10.17352/2455-8400.000039>.
- Razak T. 2008. The population of hard coral colonies growing on Ecoreef artificial modules on Manado Tua Island, Bunaken National Park, North Sulawesi, Indonesia. Oral presentation material, 11 Int Coral Reef Sym Floria, Session number XXIV, p. 223.
- Reid C., Marshall J., Logan D., Kleine D. 2009 [Coral reef and climate change]. CoralWatch, The University of Queensland. Australia. 256 pp. [in Indonesian]
- Ricardo G. F., Jones R. J., Nordborg M., Andrew P. Negri. 2017. Settlement patterns of the coral *Acropora millepora* on sediment-laden surfaces. Science of the Total Environment. 609:277-288.
- Rondo M., Tamanampo J. F. W. S. 2014 Principles of Aquatic Ecology. Fakultas Perikanan dan Ilmu Kelautan, Universitas Sam Ratulangi. Manado. 427 pp.
- Sheppard C., Davy S., Pilling G., Graham N. 2017 *The biology of coral reefs*. Oxford University Press. 365 pp.
- Sheehy D. J. 1986 New approaches in artificial reef design and applications. p. 256-263. In F. M. D. Trie (ed.) Artificial reefs: marine and freshwater applications. Lewis Publishers Inc., Michigan, USA.
- Suharsono. 2008 [Coral species in Indonesia]. Lembaga Ilmu Pengetahuan (LIPI) : COREMAP Program. Jakarta. 116 p. [in Indonesian]
- Suleman Y., Lalamentik L. Th. X., Rembet U. N. W. J. 2017 [Hard coral *Favites abdita* (Ellis and Solander, 1786) distribution in the reef flat of Malalayan 2, Malalayang district, Manado city]. Jurnal Ilmiah Platax 5(1): 104-111. Universitas Sam Ratulangi. Manado. [Indonesian] pp [Indonesian]
- Varma M. B. 2017 Properties of cement concrete reinforced with bamboo-strip-Mat. IOSR, Journal of Mechanical and Civil Engineering (IOSR-JMCE), eISSN, 2278-1684: 47-59.
- Veron J. E. N. 1986 Corals of Australia and the Indo-Pacific. Angus and Robertson Publishers. Australia. 644 pp.
- Veron J. E. N. 2000 Corals of the World Volume 1. Australian Institute of Marine Science and CRR Qld Pty Ltd. Australia 463 pp.

Veron J. E. N., Pichon M., Wijsman-Best M. 1977 Scleractinia of Eastern Australia. Canberra. Vol. 3 (*Part 2*). 233 pp.

Authors:

L.T.X. Lalamentik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: ottylalamentik@yahoo.co.id

R. Ch. Kepel, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: renecharleskepel65@gmail.com

L.J.L. Lumingas, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: ljlumingas@yahoo.com

U.N.W.J. Rembet Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: onlyrembet@unsrat.ac.id

S.B. Pratasik, Faculty of Fisheries and Marine Science, Sasm Ratulangi University, Manado-95115, email: spjong07@yahoo.com

D.M.H. Mantiri, Faculty of Fisheries and Marine Science, Sasm Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: dmh_mantiri@unsrat.ac.id

Find messages, documents, photos or people

← Back ↶ ↷ → Archive Move Delete Spam ...

Re: new article submission

Yahoo/Inbox



gavriloae ionel claudiu <ionelclaudiu@yahoo.com>
To: Benny Pratasik

Sat, Feb 22, 2020 at 1:45 AM

Dear Professor Pratasik,

I received and already performed the preliminary evaluation of the paper. The editing style was a disaster... So I have done a lot of work doing it properly. However, before sending the paper to reviewers for the fast track review, kindly solve the few problems in the References section. Thank you very much!

Cordially yours,
Claudiu G.

On Tuesday, February 18, 2020, 7:20:08 AM GMT+2, Benny Pratasik <spjong07@yahoo.com> wrote:

Dear Dr. Gavriloae,
I herewith send you a new article "Faviidae Coral Colonization Living and Growing on Agricultural Waste-Materialized Artificial Substrate" (attached). The authors wish to have fast service for the next issue. Thank you for your cooperation.

Sincerely Yours,
Silvester B. Pratasik

[Download all attachments as a zip file](#)

Find messages, documents, photos or people

← Back ↶ ↷ → Archive ↗ Move 🗑 Delete 🛡 Spam ⋮

• the review for the paper of Lamentik et al's (Faviidae coral colonization...) Yahoo/Inbox ☆

 **gavrioloie ionel claudiu** <ionelclaudiu@yahoo.com> Thu, Mar 19, 2020 at 9:05 PM ☆
To: Benny Pratasik

Dear Professor Pratasik,

I am sending you the complete review for this paper, which is excellent. However, there are some problems with the scientific names and with the references section.


So, ask the authors to work only in the attached paper and to highlight the changes they will do.

Regarding the processing fee payment (300 USD), you already have the variants.
Thank you very much!

Cordially yours,
Claudiu G.



archive (3).zip Lamentik et al - l...pdf



Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate

Laurentius T. X. Lalamentik, Rene Ch. Kepel, Lawrence J. L. Lumingas, Unstain N. W. J. Rembet, Silvester B. Pratasik, Desy M. H. Mantiri

Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia. Corresponding author: L. T. X. Lalamentik, ottylalamentik@yahoo.co.id

Abstract. A study on colonization of Faviidae corals on the agricultural waste-materialized artificial substrate was conducted in Selat Besar, Ratatotok district, southeast Minahasa regency, North Sulawesi. Nine artificial substrates moduls made of mixture of cement, sand, padi husk, and bamboo were placed for about 5 years on the sea bottom of Selat Besar waters. All corals of family Faviidae found on the artificial substrate were collected. Results showed that Faviidae corals could live and develop on those substrates. Fifteen species of 5 genera of Faviidae, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennetae*, *Echinopora gemmacea*, *E. lammelosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. senensis* were recorded in the present study. Mean number of colonies of Faviidae corals was 3 col.mod⁻¹, while mean diameter of the corals attached on the artificial substrate was 7.45 cm long. The distribution pattern of Faviidae corals was clumped. The diversity of Faviidae corals on the artificial substrate was low ($H' = 0.677 < 1$). The dominance index showed no dominant species ($D = 0.23$). In addition, the artificial substrate modul in this study could become an alternative technique to rehabilitate the degraded coral reefs.

Key Words: diversity, dominance, distribution pattern, Selat Besar.

Introduction. Corals are animals belonging to phylum Cnidaria (Reid et al 2009), while coral reef is an ecosystem built by calcium carbonate-producing marine biota, especially coral animals, together with other biota living on the sea bottom or water column, such as mollusks, crustaceans, echinoderms, porifera, tunicates, and other biota free-living in surrounding waters including plankton and fishes (Giyanto et al 2017; Lalamentik 1995). According to Barus (2013), coral reef is one of the unique communities that are entirely formed from biological activities. It is one of the main coastal and marine ecosystems that possess the highest productivity and biodiversity so that it is often called as tropical marine forest.

Coral Triangle (CTI) is known as center of world marine biodiversity with the highest coral diversity, 76% of total coral species. The region inside the ecological boundary of Coral Triangle covers nearly 73,000 km² of coral reefs (29% of total world coral reef area) distributed in six countries, Indonesia, Malaysia, Filipina, Timor Leste, Papua New Guinea and Solomon Islands (Burke et al 2012). As part of the CTI, Indonesia is an archipelagic country with 16,056 islands, 1,922,570 km² terrestrial area and 3,257,483 km² water in the west Indo-Pacific (BIG 2016). Indonesia that is located in the coral triangle area is recognized as one of the countries with high biodiversity consisting of more than 80 coral genera and 596 species, particularly in North Sulawesi waters that has been recorded more than 80 genera (Suharsono 2008).

Coral taxonomic and distribution studies in North Sulawesi were carried out by several scientists (Lalamentik 1998; Halidu et al 2016; Nasaru et al 2017; Suleman et al 2017; Lalamentik & Rembet 2018). So far, there is no study on corals of family Faviidae living and growing on the artificial substrate, especially in North Sulawesi waters. Hence, present study focuses on Faviidae coral colonies on the artificial substrate with the

objectives of identifying the coral species, analysing the density and the diameter of the coral colony, the ecological index, and assessing the distribution patterns. This finding is expected to be one of the references in coral development studies, concerning Faviidae coral colonization on the artificial substrate. Besides, it could provide alternative artificial substrates in coral reef rehabilitation program.

Material and Method

Study site. This work was accomplished in Selat Besar waters, Ratatotok district, southeast Minahasa regency (Figure 1), North Sulawesi. Selat Besar is flanked by Totok bay and Moluccas Sea. The study site was laid on 0°51'34.945''S and 124°44'14.416'' E (Figure 1) using Global Position System (GPS).




Figure 1. Sampling locality.

Artificial substrate construction. In this study, 9 artificial substrate moduls were set on the sea bottom, approximately 7-8 m depth. Sheehy (1986) stated that one of the major roles of artificial substrate placement in the water is to create new habitats similar to artificial reefs to be inhabited by target species. The use of artificial substrate as coral new habitat had been conducted in Singaporean waters in order to restore the coral reefs (Ng & Chou 2017). Each modul consists of 10 concrete blocks made of mixture of padi husk, sand, and cement. The block frames were made of bamboo. The use of bamboo in concrete construction has been done before as mentioned in Glenn (1950) and Mehra & Ghosh (1965). One of the properties that would make bamboo a good substitute to steel in reinforced concrete is its strength (Varma, 2017). The block had a dimension of 100 cm leng, 15 cm width, and 15 cm height. The blocks were set in such a manner to yield a modul. Each modul is estimated to have an area of 4.5 m². These 9 moduls were placed since June 2014. Sample collections were carried out in July 2019 (after 5 years of

deployment). The period of 5 years is determined because the massive coral growth is slow. This duration is considered enough to see whether there is Faviidae coral capable of adhering and growing on the artificial substrate.

Sample collection and handling procedures. All massive coral species found to attach on the modul substrate were taken using a chisel and hammer. The samples were then put into labelled bag as the modul number. All corals were collected by using SCUBA equipments. All activities were documented by using an underwater camera.

Before species identification, the coral specimens were soaked in bleach-containing freshwater for 24 h in order to kill the coral and to prevent the presence of fungi. The corals were then cleaned under the running water to remove the attached dirt so that the clean coral and the bleaching coral skeleton were obtained. The corals were placed under direct sunlight to get dry to ease the identification.

Species identification. Coral identification was done following Veron (1986; 2000), Suharsono (2008), and Veron et al (1977). The specimens of Faviidae corals were examined through morphological observations under the SZ51 Olympus microscope on the characteristics of paliform lobe, septa cleats, coralite type, number of first septa, second septa, and third septa. The measurements of width of paliform lobe, coralite size, and calix size used a multitoyo digital caliper.

Data analysis. Faviidae coral colonization data were directed to ecological index analyses as follows:

Dominance index. This index was employed to know the dominant species in data sampling boundary. The present study applied Simpson dominance index (Krebs, 1989):

$$D = \frac{\sum_{i=1}^s \left[\frac{n_i}{N} \right]^2}{\dots\dots\dots} \quad (1)$$

Where D = dominance index, n_i = number of i individuals, and N = total number of individuals.

Dominance index ranges between 0 and 1, if D approaches to 0, it indicates no dominance and the community structure is in stable condition, and if D approaches to 1, it indicates the presence of species dominance.

Density. The population density was calculated using Krebs (1989) as follows

$$K = \frac{N_i}{A} \quad \dots\dots\dots \quad (2)$$

Where, K = density (ind.modul^{-1}), N_i = number of individuals, and A = number of moduls where samples were collected.

Diversity. The diversity index of genus *Favia* Oken 1815 was calculated using Shannon-Wiener equation (Krebs 1989):

$$H = \sum_{i=1}^s \frac{n_i}{N} \left(\frac{\ln n_i}{N} \right) \quad \dots\dots\dots \quad (3)$$

Where H = Shannon-Wiener diversity index, n_i = number of individuals i , N = total number of individuals, and s = number of genera.

Distribution pattern. The distribution pattern of Faviidae corals were estimated using Morisita index (Krebs 1989):

$$I_d = \frac{\sum_{i=1}^q n_i (n_i - 1)}{N (N - 1)} \quad \dots\dots\dots \quad (4)$$

Where I_d = Morisita index, n_i = number individuals in each plot, n = total number of individuals in all plots, and N = number of plots, in which $I_d = 1$ is categorized as random distribution, $I_d > 1$ as clumped, and $I_d < 1$ as uniform.

Results and Discussion

Water conditions of Selat Besar. Water quality is crucial for colonization of macroinvertebrates (Orwa et al 2018). Water temperature is one of the limiting factors for aquatic organisms to live, one of which is coral. Field measurements found that water temperature in the study site ranged from 29° - 30°C. This range is still suitable for coral development. According to Kurniawan (2011), corals, in general, can live at the temperature above 18°C, while growth, they require an ideal water temperature, between 25°-32°C. Most coral reefs exist where salinity is stable and where average salinity is that of normal seawater 34-36‰ (Sheppard et al., 2017). Water salinity in the study site was between the range of 30-31‰. Guntur (2011) found that water salinity for ideal coral growth in Indonesia ranged from 29 to 36‰. Therefore, water salinity in Selat Besar highly supports the coral to grow well. Light intensity does not influence larval settlement, and their response to light intensity may be species-specific (Mundy & Babcock 1998).

Corals of family Faviidae recorded on the artificial substrate. Physical and biological factors are important for recruitment success (Doropoulos et al 2016), As many as 15 coral species of 5 genera were recorded in this study. *Favia* and *Favites* had the highest number of species, each with 4 species, followed by genus *Goniastrea* with 3 species, then *Echinopora* and *Platygyra*, 2 species, respectively. Table 1 demonstrates the coral species of Faviidae encountered on the modul substrate with mean number of colonies of 3.00 per modul. Even though these numbers are less than those reported by Kilfoyle et al (2008), 13 col.mod.⁻¹, the colonies comprised several coral families. The coral genera of Faviidae inhabiting the artificial substrate apparently come the coral reef around Selat Besar waters. It is reasonable, since Faviidae is coral family generally found in Selat Besar waters (Lalamentik 1996;1997;1998).

Table 1

Faviidae coral species encountered living on the artificial substrate modul

No	Species
1	<i>Echinopora gemmacea</i> (Lamarck, 1816)
2	<i>E. lammelosa</i> (Esper, 1795)
3	<i>Favia laxa</i> (Klunzinger, 1879)
4	<i>F. matthai</i> (Vaughan, 1918)
5	<i>F. palida</i> (Dana, 1846)
6	<i>F. stelligera</i> (Dana, 1846)
7	<i>Favites bennetae</i> (first record for Central Indonesia)
8	<i>F. complanata</i> (Ehrenberg, 1734)
9	<i>F. pentagona</i> (Esper, 1794)
10	<i>F. russelli</i> (Wells, 1954)
11	<i>Goniastrea aspera</i> (Verrill, 1864)
12	<i>G. favulus</i> (Dana, 1846)
13	<i>G. pectinata</i> (Ehrenberg, 1834)
14	<i>Platygyra daedalea</i> (Elis & Solander, 1786)
15	<i>P. sinensis</i> (Edwards & Haime, 1857)

Favites bennetae recorded in the present study is the first record found in the coral reef of Central Indonesian waters. This species was previously mostly found in the northeast of Australian waters (Veron et al 1977). Moreover, genus *Favia* was found having the highest mean number of colonies, 1.11 col.mod.⁻¹, followed by *Goniastrea*, 0.67 col.mod.⁻¹,

and the lowest was found in *Platygyra*, 0.22 col.mod⁻¹ (Figure 2). Variations in number of Faviidae colonies per modul are dependent upon the abundance of larvae that survive and settle on the substrate. Faviidae tends to have good adaptability to occupying the available artificial substrates. Bachtiar et al (2011) found that Faviidae is one of the coral families capable of surviving in the reef ball deployed on the sea bottom.




Figure 2. Mean no. colonies of *Faviidae* per modul.

Mean diameter of the coral colony on the artificial substrate was 7.45 cm with the largest recorded in *Favia laxa*, 10.5 cm and the smallest in *F. palida*, 3 cm. In comparison among genera, mean diameter of Faviidae corals attached on the artificial substrate ranged from 5.88 to 8.50 cm. Genus *Platygyra* has the largest mean diameter, 8.50 cm, and *Echinopora* had the smallest one, 5.88 cm (Figure 3). Variations in colony size could result from different sequence of planular settlements. This condition, according to Bachtiar et al (2011), indicates that coral colonization on the artificial substrate sustainably occurs.




Figure 3. Mean diameter of *Faviidae* corals living on the artificial substrate.

Ecological index. The present study measured dominance index (D) and diversity index (H') of Faviidae corals thriving on the artificial substrate. The data are given in Table 2.

Ecological index of Faviidae corals

No	Ecological index	Value	Remarks
1	Dominance Index (D)	0.23	No dominance
2	Distribution pattern (I δ)	1.79	Clumped
3	Diversity index (H')	3.64	high

The dominance index of Faviidae corals is 0.23 (Table 3). It means that no species outnumbers the others. Selat Besar waters is still capable of supporting the coral life, and there is no competition that could make certain species be dominant (Muqsit et al 2016). The diversity index is 3.64. This value is supported by low species dominance index, in which there were 15 Faviidae species found. These numbers are high enough due to small study area cover. This finding is in agreement with Rondo et al (2014) that high diversity will cause low species dominance.

Distribution pattern analysis found that Faviidae in the study site had clumped distribution (Id = 1.79 > 1). This pattern could result from the fact that there were four moduls not occupied by Faviidae corals. In Indonesia, studies on coral colonization was also done by Razak (2008) using Eco-reef modul in Manado bay waters, North Sulawesi, and found that the artificial substrate used are, in fact, inhabited by hard corals. Hence, the artificial substrate could be recommended as an alternative in coral reef rehabilitation program.

Conclusions. Fifteen coral species were found settling on the agricultural waste-materialized artificial substrate, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennetae*, *Echinopora gemmacea*, *E. lammelosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. senensis*. They had mean 3.00 col.mod⁻¹ with mean diameter of 7.45 cm. The ecological index also indicated that the available artificial substrate could support various coral species to grow. In this study, Faviidae corals growing on the available artificial substrate have reflected clumped distribution. Thus, this type of modul could be used for coral reef rehabilitation.

References [indra2]

- Bachtiar I., Prayogo W., Sukri A. 2011 Coral colonization and growth reef ball modules at the Benete Bay, Sumbawa Island, Indonesia. Manado. P. 31-40.
- Doropoulos C., Roff G., Bozec Y. M., Zupan M., Werminghausen J., Mumby P. J. 2016 Characterizing the ecological trade-offs throughout the early ontogeny of coral Error! Hyperlink reference not valid.cruitment. Ecol. Monogr. 86Error! Hyperlink reference not valid[indra3].
- Guntur. 2011 [Coral ecology in the artificial reef]. Ghalia Indonesia. Semarang. 139 pp. [in Indonesian]
- Halidu, A., Lalamentik L. T. X., Rembet U. N. W. J. 2016 [Hard coral distribution in the reef flat of the south coast of Putus-Putus island, east Ratatotok, Ratatotok district, southeast Minahasa]. Jurnal Ilmiah Platax 4(1):19-30. [in Indonesian]
- Kilfoyle K., Rangel M. A., Dodge R. E., Spieler R. E. 2008 Coral Reef Restoration: Standardized Module Intervention and monitoring program in Mexico, Preliminary Results. Poster presentation material, 11 Int Coral Reef Sym Florida, Session number XXIV, p. 541
- Krebs C. J. 1989 Ecological Methodology. Harper Collins Publishers. New York. 654 pp.
- Lalamentik L. Th. X. 1995 [Study on coral reef potential in Tombasian district, Minahasa, North Sulawesi] Fakultas Perikanan dan Ilmu Kelautan UNSRAT. Manado. 28 pp. [in Indonesian] Unpublished
- Lalamentik L. T. X. 1996 Coral Reef conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters,

- the District of Minahasa North Sulawesi. (Survey Dated October, 1996). 23 pp. Unpublished.
- Lalamentik L. T. X. 1997 Coral Reef Conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Rataotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1997). 22 pp. Unpublished.
- Lalamentik L. Th. X. 1998 Coral reef Condition Around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study for May 1998 Data in Rataotok and Adjacent Waters, The District of Minahasa North Sulawesi. Fakultas Perikanan dan Ilmu Kelautan USNRAT. Manado. 23 pp. Unpublished.
- Lalamentik, L. Th. X., Rembet, U. N. J. 2018 [Study on the Taxonomy of Genus *Favia* (Oken, 1815) at the Reef Flats of Kampung Ambong Village in Likupang Timur District, North Minahasa]. *JURNAL ILMIAH PLATAX*, 6(1), 188-193. [in Indonesian]
- Mehra S. R., Ghosh R. G. 1965 "Bamboo-reinforced soil-cement," *Civil Engineering and Public Works Review*, Vol. 60, no. 711, October; Vol. 60, no. 712. November 1965.
- Error! Hyperlink reference not valid[indra4].Role of light intensity and spectral quality in coral settleError! Hyperlink reference not valid.ment: implications for depth-dependent settlement? *J. Exp. Mar. Biol. Ecol.* 223: 235–25Error! Hyperlink reference not valid.[indra5]
- Muqsit A., Purnama D., Ta'alidin Z. 2016 [The community structure of coral reef in Pulau Dua, Enggano district, North Bengkulu regency.] *Jurnal Enggano* Vol. 1, No. 1 (75-78). Ilmu Kelautan Fakultas Pertanian Universitas Bengkulu. Bengkulu. [in Indonesian]
- Nasaru J. H., Lalamentik L. Th. X., Rembet U. N. W. J. 2017 [*Pocillopora verrucosa* (Ellis and Solander, 1786) distribution in the reef flat of the south coast of Putus-Putus island, east Rataotok village, southeast Minahasa regency]. *Jurnal Ilmiah Platax* 5(1): 96-103 pp. Universitas Sam Ratulangi Manado. [in Indonesian]
- Ng C. S. L., Chou L. M. 2017. Coral reef restoration in Singapore-past, present and future. *Sustainability Matters: Environmental Management in the Anthropocene*. Singapore: World Scientific, 3-23 pp.
- Orwa P. O., Omondi R., Chemoiwa E. J. 2018 Colonization patterns of benthic macroinvertebrates in fertilized and non-fertilized earthen fish ponds. *Int J Aquac Fish Sci* 4(3): 022-026.
- Razak T. 2008. The population of hard coral colonies growing on Ecoreef artificial modules on Manado Tua Island, Bunaken National Park, North Sulawesi, Indonesia. Oral presentation material, 11 Int Coral Reef Sym Floria, Session number XXIV, p. 223.
- Reid C., Marshall J., Logan D., Kleine D. 2009 [Coral reef and climate change]. *CoralWatch*, The University of Queensland. Australia. 256 pp. [in Indonesian]
- Ricardo G. F., Jones R. J., Nordborg M., Andrew P. Negri. 2017. Settlement patterns of the coral *Acropora millepora* on sediment-laden surfaces. *Science of the Total Environment*. 609:277-288.
- Rondo M., Tamanampo J. F. W. S. 2014 *Principles of Aquatic Ecology*. Fakultas Perikanan dan Ilmu Kelautan, Universitas Sam Ratulangi. Manado. 427 pp.
- Sheppard C., Davy S., Pilling G., Graham N. 2017 *The biology of coral reefs*. Oxford University Press. 365 pp.
- Sheehy D. J. 1986 New approaches in artificial reef design and applications. p. 256-263. In F. M. D. Trie (ed.) *Artificial reefs: marine and freshwater applications*. Lewis Publishers Inc., Michigan, USA.
- Suharsono. 2008 [Coral species in Indonesia]. Lembaga Ilmu Pengetahuan (LIPI) : COREMAP Program. Jakarta. 116 p. [in Indonesian]
- Suleman Y., Lalamentik L. Th. X., Rembet U. N. W. J. 2017 [Hard coral *Favites abdita* (Ellis and Solander, 1786) distribution in the reef flat of Malalayan 2, Malalayang district, Manado city]. *Jurnal Ilmiah Platax* 5(1): 104-111. Universitas Sam Ratulangi. Manado. [Indonesian] pp [Indonesian]

- Varma M. B. 2017 Properties of cement concrete reinforced with bamboo-strip-Mat. IOSR, Journal of Mechanical and Civil Engineering (IOSR-JMCE), eISSN, 2278-1684: 47-59.
- Veron J. E. N. 1986 Corals of Australia and the Indo-Pacific. Angus and Robertson Publishers. Australia. 644 pp.
- Veron J. E. N. 2000 Corals of the World Volume 1. Australian Institute of Marine Science and CRR Qld Pty Ltd. Australia 463 pp.
- Veron J. E. N., Pichon M., Wijsman-Best M. 1977 Scleractinia of Eastern Australia. Canberra. Vol. 3 (Part 2). 233 pp.

Authors:

L.T.X. Lalamentik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: ottylalamentik@yahoo.co.id

R. Ch. Kepel, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: renecharleskepel65@gmail.com

L.J.L. Lumingas, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: ljllumingas@yahoo.com

U.N.W.J. Rembet Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: onlyrembet@unsrat.ac.id

S.B. Pratasik, Faculty of Fisheries and Marine Science, Sasm Ratulangi University, Manado-95115, email: spjong07@yahoo.com


D.M.H. Mantiri, Faculty of Fisheries and Marine Science, Sasm Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: dmh_mantiri@unsrat.ac.id

Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate

Laurentius T. X. Lalamentik, Rene Ch. Kepel, Lawrence J. L. Lumingas, Unstain N. W. J. Rembet, Silvester B. Pratasik, Desy M. H. Mantiri

Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia. Corresponding author: L. T. X. Lalamentik, ottylalamentik@yahoo.co.id

OPEN ACCESS



Bioflux (publishing house)
54 Ceahlau Street,
Cluj-Napoca 400488,
Romania, European Union

Certificate/Letter of acceptance

This certificate shows that your manuscript entitled:

Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate

Authors:

Laurentius T. X. Lalamentik, Rene Ch. Kepel, Lawrence J. L. Lumingas, Unstain N. W. J. Rembet, Silvester B. Pratasik, Desy M. H. Mantiri

was accepted for publication in vol 13, issue 2 (2020) of the scientific/academic journal:
Aquaculture, Aquarium, Conservation & Legislation –
International Journal of the Bioflux Society.

Thank you for publishing with us!

Cordially yours,

Editor-in-chief:
Claudiu Gavriiloaie, PhD



Find messages, documents, photos or people

Back Archive Move Delete Spam

article'spayment Yahoo/Sent

B Yahoo <spjong07@yahoo.com> To: gavriloaie ionel claudiu Fri, Mar 27, 2020 at 7:12 PM

Dear Dr. Gavrioloaie,
I herewith send you the bank receipt for publication fee of Lamentik et al. The paper is being revised. Thank you.

Wishes,
Silvester B. Pratasik



Bank receipt....pdf
2.2MB

Navigation icons

Find messages, documents, photos or people

Back Archive Move Delete Spam

Re: article'spayment

Yahoo/Inbox



gavriloaie ionel claudiu <ionelclaudiu@yahoo.com>
To: Yahoo

Sat, Mar 28, 2020 at 4:04 AM

Dear Professor Pratasik,

I have received the payment document. But I am still waiting for the revised version of the paper, because I did not receive it yet. At least I was not able to find it in my Inbox folder. So, kindly check the messages to me and see if you've sent it. Thank you very much!

Cordially yours,
Claudiu G.

On Friday, March 27, 2020, 1:12:46 PM GMT+2, Yahoo <spjong07@yahoo.com> wrote:

Dear Dr. Gavrioloaie,
I herewith send you the bank receipt for publication fee of Lalamantik et al. The paper is being revised. Thank you.

Wishes,
Silvester B. Pratasik


On Saturday, March 28, 2020, 4:22:25 AM GMT+2, Yahoo <spjong07@yahoo.com> wrote:

Dear Dr. Gavrioloie,

I herewith send you the revised paper of Lalamentik et al and Rondonuwu et al (enclosed). Those articles have been corrected and edited following the reviewers' comments. I also provide some arguments in Lalamentik et al's paper to clarify the issue after discussing it with the corresponding author, particularly concerning the coral species identification. As far as I know, Lalamentik is one of the best coral taxonomists in Indonesia who focuses on morphological characteristics, and therefore, he has strongly followed Veron's coral description. All corrections are given in yellow. Also, the payment for Rondonuwu et al's publication fee will be done on next Monday. Thank you

Sincerely Yours,

Silvester B. Pratasik



Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate

Laurentius T. X. Lalamentik, Rene C. Kepel, Lawrence J. L. Lumingas, Unstain N. W. J. Rembet, Silvester B. Pratasik, Desy M. H. Mantiri

Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia. Corresponding author: L. T. X. Lalamentik, ottylalamentik@yahoo.co.id

Abstract. A study on colonization of Faviidae corals on the agricultural waste-materialized artificial substrate was conducted in Selat Besar, Ratatotok district, southeast Minahasa regency, North Sulawesi. Nine artificial substrates modules made of mixture of cement, sand, padi husk, and bamboo were placed for about 5 years on the sea bottom of Selat Besar waters. All corals of family Faviidae found on the artificial substrate were collected. Results showed that Faviidae corals could live and develop on those substrates. Fifteen species of 5 genera of family Faviidae, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennettiae*, *Echinopora gemmacea*, *E. lamellosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. sinensis* were recorded in the present study. Mean number of colonies of Faviidae corals was 3 col mod⁻¹, while mean diameter of the corals attached on the artificial substrate was 7.45 cm long. The distribution pattern of Faviidae corals was clumped. The diversity of Faviidae corals on the artificial substrate was low ($H' = 0.677 < 1$). The dominance index showed no dominant species ($D = 0.23$). In addition, the artificial substrate module in this study could become an alternative technique to rehabilitate the degraded coral reefs.

Key Words: diversity, dominance, distribution pattern, Selat Besar.

Introduction. Corals are animals belonging to phylum Cnidaria (Reid et al 2009), while coral reef is an ecosystem built by calcium carbonate-producing marine biota, especially coral animals, together with other biota living on the sea bottom or water column, such as mollusks, crustaceans, echinoderms, porifera, tunicates, and other biota free-living in surrounding waters including plankton and fishes (Giyanto et al 2017; Lalamentik 1995). According to Barus (2018), coral reef is one of the unique communities that are entirely formed from biological activities. It is one of the main coastal and marine ecosystems that possess the highest productivity and biodiversity so that it is often called as tropical marine forest.

Coral Triangle (CTI) is known as center of world marine biodiversity with the highest coral diversity, 76% of total coral species. The region inside the ecological boundary of CTI covers nearly 73,000 km² of coral reefs (29% of total world coral reef area) distributed in six countries, Indonesia, Malaysia, Philippines, Timor Leste, Papua New Guinea and Solomon Islands (Burke et al 2012). As part of the CTI, Indonesia is an archipelagic country with 16,056 islands, 1,922,570 km² terrestrial area and 3,257,483 km² water in the west Indo-Pacific (Briney 2020). Indonesia that is located in the CTI area is recognized as one of the countries with high biodiversity consisting of more than 80 coral genera and 596 species, particularly in North Sulawesi waters where have been recorded more than 80 genera (Suharsono 2008).

Coral taxonomic and distribution studies in North Sulawesi were carried out by several scientists (Lalamentik 1998; Halidu et al 2016; Nasaru et al 2017; Suleman et al 2017; Lalamentik & Rembet 2018). So far, there is no study on corals of family Faviidae living and growing on the artificial substrate, especially in North Sulawesi waters. Hence, the present study focuses on Faviidae coral colonies on the artificial substrate with the

Commented [indra1]: see the comment on Table 1

Commented [indra2]: same as above

objectives of identifying the coral species, analysing the density and the diameter of the coral colony, the ecological index, and assessing the distribution patterns. This finding is expected to be one of the references in coral development studies, concerning Faviidae coral colonization on the artificial substrate. Besides, it could provide alternative artificial substrates in coral reef rehabilitation program.

Material and Method

Study site. This work was accomplished in Selat Besar waters, Ratatotok district, southeast Minahasa regency (Figure 1), North Sulawesi. Selat Besar is flanked by Totok bay and Moluccas Sea. The study site was laid on 0°51'34.945"S and 124°44'14.416" E (Figure 1) using Global Position System (GPS).




Figure 1. Sampling locality.

Artificial substrate construction. In this study, 9 artificial substrate modules were set on the sea bottom, approximately 7-8 m depth (Figure 2). Sheehy (1986) stated that one of the major roles of artificial substrate placement in the water is to create new habitats similar to natural reefs to be inhabited by target species. The use of artificial substrate as coral new habitat had been conducted in Singaporean waters in order to restore the coral reefs (Ng & Chou 2017). Each module consists of 10 concrete blocks made of mixture of padi husk, sand, and cement. The block frames were made of bamboo. The use of bamboo in concrete construction has been done before as mentioned in Glenn (1950) and Mehra & Ghosh (1965). One of the properties that would make bamboo a good substitute to steel in reinforced concrete is its strength (Varma 2017). The block had a dimension of 100 cm length, 15 cm width, and 15 cm height. The blocks were set in such a manner to yield a module. Each module is estimated to have an area of 4.5 m². These 9 modules were placed since June 2014. Sample collections were carried out in July 2019 (after 5

years of deployment). The period of 5 years is determined because the massive coral growth is slow. This duration is considered enough to see whether there is Faviidae coral capable of adhering and growing on the artificial substrate.



Figure 2. Artificial substrate modules. Left: new module; Right: colonized module.

Sample collection and handling procedures. All massive coral species found to attach on the module substrate were taken using a chisel and hammer. The samples were then put into labelled bag as the module number. All corals were collected by using SCUBA equipments. All activities were documented by using an underwater camera.

Before species identification, the coral specimens were soaked in bleach-containing freshwater for 24 h in order to kill the coral and to prevent the presence of fungi. The corals were then cleaned under the running water to remove the attached dirt so that the clean coral and the bleaching coral skeleton were obtained. The corals were placed under direct sunlight to get dry to ease the identification.

Species identification. Coral identification was done following Veron (1986, 2000), Suharsono (2008), and Veron et al (1977). The specimens of Faviidae corals were examined through morphological observations under the SZ51 Olympus microscope on the characteristics of paliform lobe, septa cleats, coralite type, number of first septa, second septa, and third septa. The measurements of width of paliform lobe, coralite size, and calix size used an [LCD electronic digital gauge stainless vernier caliper 150 mm.](#)

Data analysis. Faviidae coral colonization data were directed to ecological index analyses as follows:

Dominance index. This index was employed to know the dominant species in data sampling boundary. The present study applied Simpson dominance index (Krebs 1989):

$$D = \frac{\sum_{i=1}^s [n_i]^2}{N^2} \dots\dots\dots (1)$$

Where D = dominance index, ni = number of i individuals, and N = total number of individuals.

Dominance index ranges between 0 and 1, if D approaches to 0, it indicates no dominance and the community structure is in stable condition, and if D approaches to 1, it indicates the presence of species dominance.

Density. The population density was calculated using Krebs (1989) as follows

$$K = \frac{Ni}{A} \dots\dots\dots (2)$$

Where, $K = \text{density (ind module}^{-1}\text{)}$, $N_i = \text{number of individuals}$, and $A = \text{number of modules where samples were collected}$.

Diversity. The diversity index of genus *Favia* Oken, 1815 was calculated using Shannon-Wiener equation (Krebs 1989):

$$H = \sum_{i=1}^s \frac{n_i}{N} \left(\frac{\ln n_i}{N} \right) \dots \dots \dots (3)$$

Where $H = \text{Shannon-Wiener diversity index}$, $n_i = \text{number of individuals } i$, $N = \text{total number of individuals}$, and $s = \text{number of genera}$.

Distribution pattern. The distribution pattern of Faviidae corals was estimated using Morisita index (Krebs 1989):

$$I_d = q \frac{\sum_{i=1}^q n_i (n_i - 1)}{N (N - 1)} \dots \dots \dots (4)$$

Where $I_d = \text{Morisita index}$, $n_i = \text{number individuals in each plot}$, and $N = \text{total number of individuals in all plots}$, in which $I_d = 1$ is categorized as random distribution, $I_d > 1$ as clumped, and $I_d < 1$ as uniform.

Results and Discussion

Water conditions of Selat Besar. Water quality is crucial for colonization of macroinvertebrates (Orwa et al 2018). Water temperature is one of the limiting factors for aquatic organisms to live, one of which is coral. Field measurements found that water temperature in the study site ranged from 29 to 30°C. This range is still suitable for coral development. According to Reid (2009), corals, in general, can live at water temperature from 18°-30°C, while for growth, they require an ideal water temperature from 27 to 29°C (Giyanto et al 2017). Most coral reefs exist where salinity is stable and where average salinity is that of normal seawater 34-36‰ (Sheppard et al 2017). Water salinity in the study site was in the range of 30-31‰. Guntur (2011) found that water salinity for ideal coral growth in Indonesia ranged from 29 to 36‰. Therefore, water salinity in Selat Besar highly supports the coral to grow well. Light intensity does not influence larval settlement, and their response to light intensity may be species-specific (Mundy & Babcock 1998).

Corals of family Faviidae recorded on the artificial substrate. Physical and biological factors are important for recruitment success (Doropoulos et al 2016). As many as 15 coral species of 5 genera were recorded in this study (Table 1). *Favia* and *Favites* had the highest number of species, each with 4 species, followed by genus *Goniastrea* with 3 species, then *Echinopora* and *Platygyra* with 2 species each. Table 1 demonstrates the coral species of Faviidae encountered on the module substrate with mean number of colonies of 3.00 per module. Even though these numbers are less than those reported by Kilfoyle et al (2008), 13 col mod⁻¹, the colonies comprised several coral families. The coral genera of Faviidae inhabiting the artificial substrate apparently come from the coral reef around Selat Besar waters. It is reasonable, since Faviidae is coral family generally found in Selat Besar waters (Lalamentik 1996, 1997, 1998).

Table 1

Faviidae coral species encountered living on the artificial substrate modul

No	Species
1	<i>Echinopora gemmacea</i> (Lamarck, 1816)
2	<i>E. lamellosa</i> (Esper, 1795)
3	<i>Favia laxa</i> (Klunzinger, 1879)
4	<i>F. matthai</i> Vaughan, 1918

Commented [indra3]: this might change if you will take into account the green comment on table 1

Commented [indra4]: when searching for the validity of the scientific names we discovered that some of the species in *Favia* genus are actually in *Goniastrea* or *Dipsastrea*. Actually, there are mismatches for almost all the five genera in the table. So, check again all the names by using reliable and updated databases (you could use <http://www.marinespecies.org>).

After checking and updating the species and genera names, you need to make all the corresponding updates and corrections both in text and figures/tables

Commented [C5R4]: All identifications were done using the standard identification book as mentioned in the method. *Favia* looks like *Goniastrea*, but they are quite different when we observe in more detail. In this study, as standard identification procedure, we observed the number of septa in primary septa, secondary septa, and tertiary septa. We also measured the coralite diameter and observe the paliform lobe.

Dipsastrea is probably not a coral name

Commented [indra6]: we removed the parentheses since this is the correct form (<http://www.marinespecies.org/aphia.php?p=taxdetails&id=207437>). We did the necessary corrections for all the authors names

Ok

5	<i>F. pallida</i> (Dana, 1846)
6	<i>F. stelligera</i> (Dana, 1846)
7	* <i>Favites bennettiae</i> Veron, Pichon & Wijsman-Best, 1977
8	<i>F. complanata</i> (Ehrenberg, 1834)
9	<i>F. pentagona</i> (Esper, 1795)
10	<i>F. russelli</i> (Wells, 1954)
11	<i>Goniastrea aspera</i> Verrill, 1866
12	<i>G. favulus</i> (Dana, 1846)
13	<i>G. pectinata</i> (Ehrenberg, 1834)
14	<i>Platygyra daedalea</i> (Elis & Solander, 1786)
15	<i>P. sinensis</i> (Milne Edwards & Haime, 1849)

* The first record for Central Indonesia.

Favites bennettiae recorded in the present study is the first record found in the coral reef of Central Indonesian waters. This species was previously mostly found in the northeast of Australian waters (Veron et al 1977). Moreover, genus *Favia* was found having the highest mean number of colonies, 1.11 col mod⁻¹, followed by *Goniastrea*, 0.67 col mod⁻¹, and the lowest was found in *Platygyra*, 0.22 col mod⁻¹ (Figure 3). Variations in number of Faviidae colonies per module are dependent upon the abundance of larvae that survive and settle on the substrate. Faviidae tends to have good adaptability to occupying the available artificial substrates. Bachtiar et al (2011) found that Faviidae is one of the coral families capable of surviving in the reef ball deployed on the sea bottom.




Figure 3. Mean number of colonies of Faviidae per module.

Mean diameter of the coral colony on the artificial substrate was 7.45 cm with the largest recorded in *Favia laxa*, 10.5 cm, and the smallest in *F. pallida*, 3 cm. In comparison among genera, mean diameter of Faviidae corals attached on the artificial substrate ranged from 5.88 to 8.50 cm. Genus *Platygyra* has the largest mean diameter, 8.50 cm, and *Echinopora* had the smallest one, 5.88 cm (Figure 4). Variations in colony size could result from different sequence of planular settlements. This condition, according to Bachtiar et al (2011), indicates that coral colonization on the artificial substrate sustainably occurs.

Commented [indra7]: when writing the numbers (both on the left side and on the top of the columns), follow the English language rules. So, instead of 1,20 you should have 1.20; 1,00 - 1.00; 0,80 - 0.80 and so on.
 The comment goes for the figure 3 as well.
 Then, *Echiopora* should be written as *Echinopora*

Commented [C8R7]: Corrected




Figure 4. Mean diameter of Faviidae corals living on the artificial substrate.

Ecological index. The present study measured dominance index (D) and diversity index (H') of Faviidae corals thriving on the artificial substrate. The data are given in Table 2.

Commented [indra9]: exactly the same comments as for figure 2

Commented [C10R9]: Corrected

Table 2

Ecological index of Faviidae corals

No	Ecological index	Value	Remarks
1	Dominance Index (D)	0.23	No dominance
2	Distribution pattern (Id)	1.79	Clumped
3	Diversity index (H')	3.64	high

The dominance index of Faviidae corals is 0.23 (Table 2). It means that no species outnumbers the others. Selat Besar waters is still capable of supporting the coral life, and there is no competition that could make certain species be dominant (Muqsit et al 2016). The diversity index is 3.64. This value is supported by low species dominance index, in which there were 15 Faviidae species found. These numbers are high enough due to small study area cover. This finding is in agreement with Rondo et al (2014) that high diversity will cause low species dominance.

Distribution pattern analysis found that Faviidae in the study site had clumped distribution ($Id = 1.79 > 1$). This pattern could result from the fact that there were four modules not occupied by Faviidae corals. In Indonesia, studies on coral colonization was also done by Razak (2008) using eco-reef module in Manado bay waters, North Sulawesi, and found that the artificial substrate used are, in fact, inhabited by hard corals. Hence, the artificial substrate could be recommended as an alternative in coral reef rehabilitation program.

Conclusions. Fifteen coral species were found settling on the agricultural waste-materialized artificial substrate, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennettiae*, *Echinopora gemmacea*, *E. lamellosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. sinensis*. They had mean 3.00 col mod⁻¹ with mean diameter of 7.45 cm. The ecological index also indicated that the available artificial substrate could support various coral species to grow. In this study, Faviidae corals growing on the available artificial substrate have reflected clumped distribution. Thus, this type of module could be used for coral reef rehabilitation.

References

Commented [indra11]: edit the list following the journal's format

Commented [C12R11]: corrected

- Bachtiar I., Prayogo W., Sukri A., 2011 Coral colonization and growth reef ball modules at the Benete Bay, Sumbawa Island, Indonesia. Manado. P. 31-40.
- Barus, B. S., Prartono T., Soedarma D., 2018 Environmental impact on the lifeform of coral reefs in Lampung bay waters] *Jurnal Ilmu dan Tehnologi Kelautan Tropis* 10(3): 699-709 [in Indonesian]
- Briney A., 2020 Geography of Indonesia: Learn About the World's Largest Archipelago Nation. ThoughtCo. Accessed from <https://www.thoughtco.com/geography-of-indonesia-1435052>
- Burke L., Reyntar K., Spalding M., Perry A., 2012 [Looking back the threatened coral reefs in the coral triangle]. World Resources Institute. USA. 76 p. [in Indonesian]
- Doropoulos C., Roff G., Bozec Y. M., Zupan M., Werminghausen J., Mumby P. J., 2016 Characterizing the ecological trade-offs throughout the early ontogeny of coral recruitment. *Ecol. Monogr.* 86: 20-44.
- Giyanto, Abrar M., Hadi T. A., Budianto A., Hafizt M., Salatahpoly A., Iswari M. Y., 2017 [The status of Indonesian coral reefs 2017]. Penelitian Oceanografi LIPI Jakarta. 30 p. [in Indonesian]
- Glenn H. E., 1950 Bamboo-reinforcement in Portland cement concrete. Engineering Experiment Station, Clamson Agricultural College, Clamson, South Caroline, Bulletin no. 4. May 1950. 171 p.
- Guntur, 2011 [Coral ecology in the artificial reef]. Ghalia Indonesia. Semarang. 139 pp. [in Indonesian]
- Halidu A., Lalamentik L. T. X., Rembet U. N. W. J., 2016 [Hard coral distribution in the reef flat of the south coast of Putus-Putus island, east Ratatotok, Ratatotok district, southeast Minahasa]. *Jurnal Ilmiah Platax* 4(1):19-30. [in Indonesian]
- Kilfoyle K., Rangel M. A., Dodge R. E., Spieler R. E., 2008 Coral Reef Restoration: Standardized Module Intervention and monitoring program in Mexico, Preliminary Results. Poster presentation material, 11 Int Coral Reef Sym Florida, Session number XXIV, p. 541
- Krebs C. J., 1989 *Ecological Methodology*. Harper Collins Publishers. New York. 654 pp.
- Lalamentik L. Th. X., 1995 [Study on coral reef potential in Tombasian district, Minahasa, North Sulawesi] Fakultas Perikanan dan Ilmu Kelautan UNSRAT. Manado. 28 pp. [in Indonesian] Unpublished
- Lalamentik L. T. X., 1996 Coral Reef conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1996). 23 pp. Unpublished.
- Lalamentik L. T. X., 1997 Coral Reef Conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1997). 22 pp. Unpublished.
- Lalamentik L. Th. X., 1998 Coral reef Condition Arround the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study for May 1998 Data in Ratatotok and Adjacent Waters, The District of Minahasa North Sulawesi. Fakultas Perikanan dan Ilmu Kelautan UNSRAT. Manado. 23 pp. Unpublished.
- Lalamentik L. Th. X., Rembet U. N. J., 2018 [Study on the Taxonomy of Genus Favia (Oken, 1815) at the Reef Flats of Kampung Ambong Village in Likupang Timur District, North Minahasa]. *JURNAL ILMIAH PLATAX*, 6(1), 188-193. [in Indonesian]
- Mehra S. R., Ghosh R. G., 1965 "Bamboo-reinforced soil-cement," *Civil Engineering and Public Works Review*, Vol. 60, no. 711, October; Vol. 60, no. 712. November 1965.
- Mundy C. N., Babcock R., 1998 Role of light intensity and spectral quality in coral settleError! Hyperlink reference not valid.ment: implications for depth-dependent settlement? *J. Exp. Mar. Biol. Ecol.* 223 (2): 235-255.
- Muqsit A., Purnama D., Ta'alidin Z., 2016 [The community structure of coral reef in Pulau Dua, Enggano district, North Bengkulu regency.] *Jurnal Enggano* Vol. 1, No. 1 (75-78). Ilmu Kelautan Fakultas Pertanian Universitas Bengkulu. Bengkulu. [in Indonesian]

- Nasaru J. H., Lalamentik L. Th. X., Rembet U. N. W. J., 2017 [*Pocillopora verrucosa* (Ellis and Solander, 1786) distribution in the reef flat of the south coast of Putus-Putus island, east Rataatok village, southeast Minahasa regency]. *Jurnal Ilmiah Platax* 5(1): 96-103 pp. Universitas Sam Ratulangi Manado. [in Indonesian]
- Ng C. S. L., Chou L. M., 2017 Coral reef restoration in Singapore-past, present and future. *Sustainability Matters: Environmental Management in the Anthropocene*. Singapore: World Scientific, 3-23 pp.
- Orwa P. O., Omondi R., Chemoiwa E. J., 2018 Colonization patterns of benthic macroinvertebrates in fertilized and non-fertilized earthen fish ponds. *Int J Aquac Fish Sci* 4(3): 022-026.
- Razak T., 2008 The population of hard coral colonies growing on Ecoreef artificial modules on Manado Tua Island, Bunaken National Park, North Sulawesi, Indonesia. Oral presentation material, 11 Int Coral Reef Sym Floria, Session number XXIV, p. 223.
- Reid C., Marshall J., Logan D., Kleine D., 2009 [Coral reef and climate change]. *CoralWatch*, The University of Queensland. Australia. 256 pp. [in Indonesian]
- Rondo M., Tamanampo J. F. W. S., 2014 *Principles of Aquatic Ecology*. Fakultas Perikanan dan Ilmu Kelautan, Universitas Sam Ratulangi. Manado. 427 pp.
- Sheppard C., Davy S., Pilling G., Graham N., 2017 *The biology of coral reefs*. Oxford University Press. 365 pp.
- Sheehy D. J., 1986 New approaches in artificial reef design and applications. p. 256-263. In F. M. D. Trie (ed.) *Artificial reefs: marine and freshwater applications*. Lewis Publishers Inc., Michigan, USA.
- Suharsono, 2008 [Coral species in Indonesia]. Lembaga Ilmu Pengetahuan (LIPI) : COREMAP Program. Jakarta. 116 p. [in Indonesian]
- Suleman Y., Lalamentik L. Th. X., Rembet U. N. W. J., 2017 [Hard coral *Favites abdita* (Ellis and Solander, 1786) distribution in the reef flat of Malalayan 2, Malalayang district, Manado city]. *Jurnal Ilmiah Platax* 5(1): 104-111. Universitas Sam Ratulangi. Manado. [Indonesian]
- Varma M. B., 2017 Properties of cement concrete reinforced with bamboo-strip-Mat. *IOSR, Journal of Mechanical and Civil Engineering (IOSR-JMCE)*, eISSN, 2278-1684: 47-59.
- Veron J. E. N., 1986 *Corals of Australia and the Indo-Pacific*. Angus and Robertson Publishers. Australia. 644 pp.
- Veron J. E. N., 2000 *Corals of the World Volume 1*. Australian Institute of Marine Science and CRR Qld Pty Ltd. Australia 463 pp.
- Veron J. E. N., Pichon M., Wijsman-Best M., 1977 *Scleractinia of Eastern Australia*. Canberra. Vol. 3 (Part 2). 233 pp.

Received: 18 February 2020. Accepted: 17 March 2020. Published online: xx March 2020.

Authors:

Laurentius T. X. Lalamentik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: ottylalamentik@yahoo.co.id
 R. Ch. Kepel, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: renecharleskepel65@gmail.com
 Lawrence J. L. Lumingas, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: ljllumingas@yahoo.com
 Unstain N. W. J. Rembet, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: onlyrembet@unsrat.ac.id
 Silvester B. Pratasik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: spjong07@yahoo.com

Desy M. H. Mantiri, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: dmh_mantiri@unsrat.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Lalamentik L. T. X., Kepel R. C., Lumingas L. J. L., Rembet U. N. W. J., Pratasik S. B., Mantiri D. M. H., 2020 Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate. AACL Bioflux 13(2):xxx-xxx.

Re: revised paper
Yahoo/Inbox

gavriloaie ionel claudiu <ionelclaudiu@yahoo.com>

To: Yahoo

Tue, Mar 31, 2020 at 12:14 AM

Dear Professor Pratasik,

Regarding the comments on the paper of professor Lalamentik, both reviewers and myself had to search on the Internet for the validity of coral Latin names. We used marinespecies.org this time. For example, as response to our comments, professor Lalamentik said that *Dipsastraea* is probably not a coral genus (we said that *Favia pallida* is actually classified as *Dipsastraea pallida* - see this: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=758233>, and this: Hoeksema, B. W.; Cairns, S. (2020). World List of Scleractinia. *Dipsastraea pallida* (Dana, 1846). Accessed through: World Register of Marine Species at: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=758233> on 2020-03-30.). So, we have no doubt that professor Lalamentik is a really great scientist and an expert in the coral species. However, we cannot know 100% a scientist is aware about the latest changes in the taxonomy. The books of dr. Veron were published in 2000. How many changes occurred in this field in the last 20 years?

And thus we are at the same point: how much can we trust the databasis?

For example, for fish species I know many Romanian and foreigner scientist and I often ask them via the phone or e-mail about one aspect or another regarding the taxonomy and systematics. But I do know anybody in Romania who knows much about the corals, since we only have few species in the Black Sea.

So, this is a day by day challenge for us here on Bioflux. All the advices are much appreciated.

I apologise for sending you such extended message. Thank you very much for your time and consideration.

Cordially yours,

Claudiu G.


On Saturday, March 28, 2020, 4:22:25 AM GMT+2, Yahoo <spjong07@yahoo.com> wrote:

Dear Dr. Gavriolaie,

I herewith send you the revised paper of Lalamentik et al and Rondonuwu et al (enclosed). Those articles have been corrected and edited following the reviewers' comments. I also provide some arguments in Lalamentik et al's paper to clarify the issue after discussing it with the corresponding author, particularly concerning the coral species identification. As far as I know, Lalamentik is one of the best coral taxonomists in Indonesia who focuses on morphological characteristics, and therefore, he has strongly followed Veron's coral description. All corrections are given in yellow. Also, the payment for Rondonuwu et al's publication fee will be done on next Monday. Thank you

Sincerely Yours,

Silvester B. Pratasik



Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate

Laurentius T. X. Lalamentik, Rene C. Kepel, Lawrence J. L. Lumingas, Unstain N. W. J. Rembet, Silvester B. Pratasik, Desy M. H. Mantiri

Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia. Corresponding author: L. T. X. Lalamentik, ottylalamentik@yahoo.co.id

Abstract. A study on colonization of Faviidae corals on the agricultural waste-materialized artificial substrate was conducted in Selat Besar, Ratatotok district, southeast Minahasa regency, North Sulawesi. Nine artificial substrates modules made of mixture of cement, sand, padi husk, and bamboo were placed for about 5 years on the sea bottom of Selat Besar waters. All corals of family Faviidae found on the artificial substrate were collected. Results showed that Faviidae corals could live and develop on those substrates. Fifteen species of 5 genera of Faviidae, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennettiae*, *Echinopora gemmacea*, *E. lamellosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. sinensis* were recorded in the present study. Mean number of colonies of Faviidae corals was 3 col mod⁻¹, while mean diameter of the corals attached on the artificial substrate was 7.45 cm long. The distribution pattern of Faviidae corals was clumped. The diversity of Faviidae corals on the artificial substrate was low ($H' = 0.677 < 1$). The dominance index showed no dominant species ($D = 0.23$). In addition, the artificial substrate module in this study could become an alternative technique to rehabilitate the degraded coral reefs.

Key Words: diversity, dominance, distribution pattern, Selat Besar.

Introduction. Corals are animals belonging to phylum Cnidaria (Reid et al 2009), while coral reef is an ecosystem built by calcium carbonate-producing marine biota, especially coral animals, together with other biota living on the sea bottom or water column, such as mollusks, crustaceans, echinoderms, porifera, tunicates, and other biota free-living in surrounding waters including plankton and fishes (Giyanto et al 2017; Lalamentik 1995). According to Barus (2013), coral reef is one of the unique communities that are entirely formed from biological activities. It is one of the main coastal and marine ecosystems that possess the highest productivity and biodiversity so that it is often called as tropical marine forest.

Coral Triangle (CTI) is known as center of world marine biodiversity with the highest coral diversity, 76% of total coral species. The region inside the ecological boundary of CTI covers nearly 73,000 km² of coral reefs (29% of total world coral reef area) distributed in six countries, Indonesia, Malaysia, Philippines, Timor Leste, Papua New Guinea and Solomon Islands (Burke et al 2012). As part of the CTI, Indonesia is an archipelagic country with 16,056 islands, 1,922,570 km² terrestrial area and 3,257,483 km² water in the west Indo-Pacific (BIG 2016). Indonesia that is located in the CTI area is recognized as one of the countries with high biodiversity consisting of more than 80 coral genera and 596 species, particularly in North Sulawesi waters where have been recorded more than 80 genera (Suharsono 2008).

Coral taxonomic and distribution studies in North Sulawesi were carried out by several scientists (Lalamentik 1998; Halidu et al 2016; Nasaru et al 2017; Suleman et al 2017; Lalamentik & Rembet 2018). So far, there is no study on corals of family Faviidae living and growing on the artificial substrate, especially in North Sulawesi waters. Hence, the present study focuses on Faviidae coral colonies on the artificial substrate with the

Commented [indra1]: see the comment on Table 1

Commented [indra2]: same as above

Commented [indra3]: it is missing from the list

Commented [indra4]: to be added to the list

Commented [indra5]: add to list

Commented [indra6]: add to list

objectives of identifying the coral species, analysing the density and the diameter of the coral colony, the ecological index, and assessing the distribution patterns. This finding is expected to be one of the references in coral development studies, concerning Faviidae coral colonization on the artificial substrate. Besides, it could provide alternative artificial substrates in coral reef rehabilitation program.

Material and Method

Study site. This work was accomplished in Selat Besar waters, Ratatotok district, southeast Minahasa regency (Figure 1), North Sulawesi. Selat Besar is flanked by Totok bay and Moluccas Sea. The study site was laid on 0°51'34.945''S and 124°44'14.416'' E (Figure 1) using Global Position System (GPS).




Figure 1. Sampling locality.

Artificial substrate construction. In this study, 9 artificial substrate modules were set on the sea bottom, approximately 7-8 m depth. Sheehy (1986) stated that one of the major roles of artificial substrate placement in the water is to create new habitats similar to artificial reefs to be inhabited by target species. The use of artificial substrate as coral new habitat had been conducted in Singaporean waters in order to restore the coral reefs (Ng & Chou 2017). Each modul consists of 10 concrete blocks made of mixture of padi husk, sand, and cement. The block frames were made of bamboo. The use of bamboo in concrete construction has been done before as mentioned in Glenn (1950) and Mehra & Ghosh (1965). One of the properties that would make bamboo a good substitute to steel in reinforced concrete is its strength (Varma 2017). The block had a dimension of 100 cm length, 15 cm width, and 15 cm height. The blocks were set in such a manner to yield a module. Each module is estimated to have an area of 4.5 m². These 9 modules were placed since June 2014. Sample collections were carried out in July 2019 (after 5 years of

Commented [indra7]: it would be great to add some photo or drawing depicting one modul

Commented [indra8]: may be natural?

Commented [indra9]: add to list

deployment). The period of 5 years is determined because the massive coral growth is slow. This duration is considered enough to see whether there is Faviidae coral capable of adhering and growing on the artificial substrate.

Sample collection and handling procedures. All massive coral species found to attach on the module substrate were taken using a chisel and hammer. The samples were then put into labelled bag as the module number. All corals were collected by using SCUBA equipments. All activities were documented by using an underwater camera.

Before species identification, the coral specimens were soaked in bleach-containing freshwater for 24 h in order to kill the coral and to prevent the presence of fungi. The corals were then cleaned under the running water to remove the attached dirt so that the clean coral and the bleaching coral skeleton were obtained. The corals were placed under direct sunlight to get dry to ease the identification.

Species identification. Coral identification was done following Veron (1986, 2000), Suharsono (2008), and Veron et al (1977). The specimens of Faviidae corals were examined through morphological observations under the SZ51 Olympus microscope on the characteristics of paliform lobe, septa cleats, coralite type, number of first septa, second septa, and third septa. The measurements of width of paliform lobe, coralite size, and calix size used a multitoyo digital caliper.

Data analysis. Faviidae coral colonization data were directed to ecological index analyses as follows:

Dominance index. This index was employed to know the dominant species in data sampling boundary. The present study applied Simpson dominance index (Krebs 1989):

$$D = \frac{\sum_{i=1}^s [n_i]^2}{N^2} \dots\dots\dots (1)$$

Where D = dominance index, n_i = number of i individuals, and N = total number of individuals.

Dominance index ranges between 0 and 1, if D approaches to 0, it indicates no dominance and the community structure is in stable condition, and if D approaches to 1, it indicates the presence of species dominance.

Density. The population density was calculated using Krebs (1989) as follows

$$K = \frac{N_i}{A} \dots\dots\dots (2)$$

Where, K = density (ind module⁻¹), N_i = number of individuals, and A = number of modules where samples were collected.

Diversity. The diversity index of genus *Favia* Oken, 1815 was calculated using Shannon-Wiener equation (Krebs 1989):

$$H = -\sum_{i=1}^s \frac{n_i}{N} \left(\frac{n_i}{N} \right) \dots\dots\dots (3)$$

Where H = Shannon-Wiener diversity index, n_i = number of individuals i , N = total number of individuals, and s = number of genera.

Distribution pattern. The distribution pattern of Faviidae corals was estimated using Morisita index (Krebs 1989):

$$I_d = \frac{\sum_{i=1}^q n_i (n_i - 1)}{N (N - 1)} \dots\dots\dots (4)$$

Commented [indra10]: ???

Where I_d = Morisita index, n_i = number individuals in each plot, n = total number of individuals in all plots, and N = number of plots, in which $I_d = 1$ is categorized as random distribution, $I_d > 1$ as clumped, and $I_d < 1$ as uniform.

- Commented [indra11]: we see no "n" in the equation
- Commented [indra12]:
- Commented [indra13R12]: is this I_d actually?

Results and Discussion

Water conditions of Selat Besar. Water quality is crucial for colonization of macroinvertebrates (Orwa et al 2018). Water temperature is one of the limiting factors for aquatic organisms to live, one of which is coral. Field measurements found that water temperature in the study site ranged from 29 to 30°C. This range is still suitable for coral development. According to Kurniawan (2011), corals, in general, can live at the temperature above 18°C, while growth, they require an ideal water temperature, between 25 and 32°C. Most coral reefs exist where salinity is stable and where average salinity is that of normal seawater 34-36‰ (Sheppard et al 2017). Water salinity in the study site was in the range of 30-31‰. Guntur (2011) found that water salinity for ideal coral growth in Indonesia ranged from 29 to 36‰. Therefore, water salinity in Selat Besar highly supports the coral to grow well. Light intensity does not influence larval settlement, and their response to light intensity may be species-specific (Mundy & Babcock 1998).

- Commented [indra14]: add to list

Corals of family Faviidae recorded on the artificial substrate. Physical and biological factors are important for recruitment success (Doropoulos et al 2016). As many as 15 coral species of 5 genera were recorded in this study. *Favia* and *Favites* had the highest number of species, each with 4 species, followed by genus *Goniastrea* with 3 species, then *Echinopora* and *Platygyra* with 2 species each. Table 1 demonstrates the coral species of Faviidae encountered on the module substrate with mean number of colonies of 3.00 per module. Even though these numbers are less than those reported by Kilfoyle et al (2008), 13 col mod⁻¹, the colonies comprised several coral families. The coral genera of Faviidae inhabiting the artificial substrate apparently come the coral reef around Selat Besar waters. It is reasonable, since Faviidae is coral family generally found in Selat Besar waters (Lalamentik 1996, 1997, 1998).

- Commented [indra15]: this might change if you will take into account the green comment on table 1

- Commented [indra16]: it is Table 1, not 02, since this is the first table in the paper

- Commented [indra17]: it has no meaning

Table 1
Faviidae coral species encountered living on the artificial substrate modul

No	Species
1	<i>Echinopora gemmacea</i> (Lamarck, 1816)
2	<i>E. lamellosa</i> (Esper, 1795)
3	<i>Favia laxa</i> (Klunzinger, 1879)
4	<i>F. matthai</i> (Vaughan, 1918)
5	<i>F. pallida</i> (Dana, 1846)
6	<i>F. stelligera</i> (Dana, 1846)
7	* <i>Favites bennettiae</i> Veron, Pichon & Wijsman-Best, 1977
8	<i>F. complanata</i> (Ehrenberg, 1834)
9	<i>F. pentagona</i> (Esper, 1795)
10	<i>F. russelli</i> (Wells, 1954)
11	<i>Goniastrea aspera</i> Verrill, 1866
12	<i>G. favulus</i> (Dana, 1846)
13	<i>G. pectinata</i> (Ehrenberg, 1834)
14	<i>Platygyra daedalea</i> (Elis & Solander, 1786)
15	<i>P. sinensis</i> (Milne Edwards & Haime, 1849)

- Commented [indra18]: when searching for the validity of the scientific names we discovered that some of the species in *Favia* genus are actually in *Goniastrea* or *Dipsastrea*. Actually, there are mismatches for almost all the five genera in the table. So, check again all the names by using reliable and updated databases (you could use <http://www.marinespecies.org>).

- After checking and updating the species and genera names, you need to make all the corresponding updates and corrections both in text and figures/tables

- Commented [indra19]: we removed the parentheses since this is the correct form (<http://www.marinespecies.org/aphia.php?p=taxdetails&id=207437>). We did the necessary corrections for all the authors names

* The first record for Central Indonesia.

Favites bennettiae recorded in the present study is the first record found in the coral reef of Central Indonesian waters. This species was previously mostly found in the northeast of Australian waters (Veron et al 1977). Moreover, genus *Favia* was found having the

highest mean number of colonies, 1.11 col mod⁻¹, followed by *Goniastrea*, 0.67 col mod⁻¹, and the lowest was found in *Platygyra*, 0.22 col mod⁻¹ (Figure 2). Variations in number of Faviidae colonies per module are dependent upon the abundance of larvae that survive and settle on the substrate. Faviidae tends to have good adaptability to occupying the available artificial substrates. Bachtiar et al (2011) found that Faviidae is one of the coral families capable of surviving in the reef ball deployed on the sea bottom.




Figure 2. Mean no. colonies of Faviidae per modul.

Mean diameter of the coral colony on the artificial substrate was 7.45 cm with the largest recorded in *Favia laxa*, 10.5 cm and the smallest in *F. pallida*, 3 cm. In comparison among genera, mean diameter of Faviidae corals attached on the artificial substrate ranged from 5.88 to 8.50 cm. Genus *Platygyra* has the largest mean diameter, 8.50 cm, and *Echinopora* had the smallest one, 5.88 cm (Figure 3). Variations in colony size could result from different sequence of planular settlements. This condition, according to Bachtiar et al (2011), indicates that coral colonization on the artificial substrate sustainably occurs.




Figure 3. Mean diameter of Faviidae corals living on the artificial substrate.

Ecological index. The present study measured dominance index (D) and diversity index (H') of Faviidae corals thriving on the artificial substrate. The data are given in Table 2.

Commented [indra20]: when writing the numbers (both on the left side and on the top of the columns), follow the English language rules. So, instead of 1,20 you should have 1.20; 1,00 - 1.00; 0,80 - 0.80 and so on.

The comment goes for the figure 3 as well.

Then, *Echiopora* should be written as *Echinopora*

Commented [indra21]: exactly the same comments as for figure 2

Table 2

Ecological index of Faviidae corals

No	Ecological index	Value	Remarks
1	Dominance Index (D)	0.23	No dominance
2	Distribution pattern (Id)	1.79	Clumped
3	Diversity index (H')	3.64	high

The dominance index of Faviidae corals is 0.23 (Table 2). It means that no species outnumbers the others. Selat Besar waters is still capable of supporting the coral life, and there is no competition that could make certain species be dominant (Muqsit et al 2016). The diversity index is 3.64. This value is supported by low species dominance index, in which there were 15 Faviidae species found. These numbers are high enough due to small study area cover. This finding is in agreement with Rondo et al (2014) that high diversity will cause low species dominance.

Distribution pattern analysis found that Faviidae in the study site had clumped distribution (Id = 1.79 > 1). This pattern could result from the fact that there were four modules not occupied by Faviidae corals. In Indonesia, studies on coral colonization was also done by Razak (2008) using eco-reef module in Manado bay waters, North Sulawesi, and found that the artificial substrate used are, in fact, inhabited by hard corals. Hence, the artificial substrate could be recommended as an alternative in coral reef rehabilitation program.

Conclusions. Fifteen coral species were found settling on the agricultural waste-materialized artificial substrate, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennettae*, *Echinopora gemmacea*, *E. lamellosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. sinensis*. They had mean 3.00 col mod⁻¹ with mean diameter of 7.45 cm. The ecological index also indicated that the available artificial substrate could support various coral species to grow. In this study, Faviidae corals growing on the available artificial substrate have reflected clumped distribution. Thus, this type of module could be used for coral reef rehabilitation.

References

- Bachtiar I., Prayogo W., Sukri A. 2011 Coral colonization and growth reef ball modules at the Benete Bay, Sumbawa Island, Indonesia. Manado. P. 31-40.
- Doropoulos C., Roff G., Bozec Y. M., Zupan M., Werminghausen J., Mumby P. J., 2016 Characterizing the ecological trade-offs throughout the early ontogeny of coral recruitment. Ecol. Monogr. 86: 20-44.
- Guntur. 2011 [Coral ecology in the artificial reef]. Ghalia Indonesia. Semarang. 139 pp. [in Indonesian]
- Halidu, A., Lalamentik L. T. X., Rembet U. N. W. J. 2016 [Hard coral distribution in the reef flat of the south coast of Putus-Putus island, east Ratatotok, Ratatotok district, southeast Minahasa]. Jurnal Ilmiah Platax 4(1):19-30. [in Indonesian]
- Kilfoyle K., Rangel M. A., Dodge R. E., Spieler R. E. 2008 Coral Reef Restoration: Standardized Module Intervention and monitoring program in Mexico, Preliminary Results. Poster presentation material, 11 Int Coral Reef Sym Florida, Session number XXIV, p. 541
- Krebs C. J. 1989 Ecological Methodology. Harper Collins Publishers. New York. 654 pp.
- Lalamentik L. Th. X. 1995 [Study on coral reef potential in Tombasian district, Minahasa, North Sulawesi] Fakultas Perikanan dan Ilmu Kelautan UNSRAT. Manado. 28 pp. [in Indonesian] Unpublished
- Lalamentik L. T. X. 1996 Coral Reef conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1996). 23 pp. Unpublished.

Commented [indra22]: edit the list following the journal's format

- Lalamentik L. T. X. 1997 Coral Reef Conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1997). 22 pp. Unpublished.
- Lalamentik L. Th. X. 1998 Coral reef Condition Arround the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study for May 1998 Data in Ratatotok and Adjacent Waters, The District of Minahasa North Sulawesi. Fakultas Perikanan dan Ilmu Kelautan USNRAT. Manado. 23 pp. Unpublished.
- Lalamentik, L. Th. X., Rembet, U. N. J. 2018 [Study on the Taxonomy of Genus Favia (Oken, 1815) at the Reef Flats of Kampung Ambong Village in Likupang Timur District, North Minahasa]. JURNAL ILMIAH PLATAX, 6(1), 188-193. [in Indonesian]
- Mehra S. R., Ghosh R. G. 1965 "Bamboo-reinforced soil-cement," Civil Engineering and Public Works Review, Vol. 60, no. 711, October; Vol. 60, no. 712. November 1965.
- Mundy C. N., Babcock R., 1998 Role of light intensity and spectral quality in coral settlement? J. Exp. Mar. Biol. Ecol. 223 (2): 235-255.
- Muqsit A., Purnama D., Ta'alidin Z. 2016 [The community structure of coral reef in Pulau Dua, Enggano district, North Bengkulu regency.] Jurnal Enggano Vol. 1, No. 1 (75-78). Ilmu Kelautan Fakultas Pertanian Universitas Bengkulu. Bengkulu. [in Indonesian]
- Nasaru J. H., Lalamentik L. Th. X., Rembet U. N. W. J. 2017 [Pocillopora verrucosa (Ellis and Solander, 1786) distribution in the reef flat of the south coast of Putus-Putus island, east Ratatotok village, southeast Minahasa regency]. Jurnal Ilmiah Platax 5(1): 96-103 pp. Universitas Sam Ratulangi Manado. [in Indonesian]
- Ng C. S. L., Chou L. M. 2017. Coral reef restoration in Singapore-past, present and future. Sustainability Matters: Environmental Management in the Anthropocene. Singapore: World Scientific, 3-23 pp.
- Orwa P. O., Omondi R., Chemoiwa E. J. 2018 Colonization patterns of benthic macroinvertebrates in fertilized and non-fertilized earthen fish ponds. Int J Aquac Fish Sci 4(3): 022-026.
- Razak T. 2008. The population of hard coral colonies growing on Ecoreef artificial modules on Manado Tua Island, Bunaken National Park, North Sulawesi, Indonesia. Oral presentation material, 11 Int Coral Reef Sym Floria, Session number XXIV, p. 223.
- Reid C., Marshall J., Logan D., Kleine D. 2009 [Coral reef and climate change]. CoralWatch, The University of Queensland. Australia. 256 pp. [in Indonesian]
- Ricardo G. F., Jones R. J., Nordborg M., Andrew P. Negri. 2017. Settlement patterns of the coral *Acropora millepora* on sediment-laden surfaces. Science of the Total Environment. 609:277-288.
- Rondo M., Tamanampo J. F. W. S. 2014 Principles of Aquatic Ecology. Fakultas Perikanan dan Ilmu Kelautan, Universitas Sam Ratulangi. Manado. 427 pp.
- Sheppard C., Davy S., Pilling G., Graham N. 2017 The biology of coral reefs. Oxford University Press. 365 pp.
- Sheehy D. J. 1986 New approaches in artificial reef design and applications. p. 256-263. In F. M. D. Trie (ed.) Artificial reefs: marine and freshwater applications. Lewis Publishers Inc., Michigan, USA.
- Suharsono. 2008 [Coral species in Indonesia]. Lembaga Ilmu Pengetahuan (LIPI) : COREMAP Program. Jakarta. 116 p. [in Indonesian]
- Suleman Y., Lalamentik L. Th. X., Rembet U. N. W. J. 2017 [Hard coral *Favites abdita* (Ellis and Solander, 1786) distribution in the reef flat of Malalayan 2, Malalayang district, Manado city]. Jurnal Ilmiah Platax 5(1): 104-111. Universitas Sam Ratulangi. Manado. [Indonesian] pp [Indonesian]
- Varma M. B. 2017 Properties of cement concrete reinforced with bamboo-strip-Mat. IOSR, Journal of Mechanical and Civil Engineering (IOSR-JMCE), eISSN, 2278-1684: 47-59.
- Veron J. E. N. 1986 Corals of Australia and the Indo-Pacific. Angus and Robertson Publishers. Australia. 644 pp.

Commented [indra23]: to be quoted in text or to be removed from the list

Veron J. E. N. 2000 Corals of the World Volume 1. Australian Institute of Marine Science and CRR Qld Pty Ltd. Australia 463 pp.
Veron J. E. N., Pichon M., Wijsman-Best M. 1977 Scleractinia of Eastern Australia. Canberra. Vol. 3 (Part 2). 233 pp.

Received: 18 February 2020. Accepted: 17 March 2020. Published online: xx March 2020.

Authors:

Laurentius T. X. Lalamentik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: ottylalamentik@yahoo.co.id

R. Ch. Kepel, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: renecharleskepel65@gmail.com

Lawrence J. L. Lumingas, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: ljllumingas@yahoo.com

Unstain N. W. J. Rembet, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: onlyrembet@unsrat.ac.id

Silvester B. Pratasik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: spjong07@yahoo.com

Desy M. H. Mantiri, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: dmh_mantiri@unsrat.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Lalamentik L. T. X., Kepel R. C., Lumingas L. J. L., Rembet U. N. W. J., Pratasik S. B., Mantiri D. M. H., 2020 Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate. AACL Bioflux 13(2):xxx-xxx.

Browser tabs: (180 unread) - spjong07@yahoo.com

URL: //mail.yahoo.com/d/folders/1/messages/AKNiM31Q8bRPXoOGygTtuCflrRE?reason=invalid_crumb

Navigation: NEWS FINANCE SPORTS ENTERTAINMENT LIFE SHOPPING YAHOO PLUS MORE...

Search: Find messages, documents, photos or people

Actions: Back, Archive, Move, Delete, Spam

Subject: Re: revised paper (Yahoo/Inbox)

From: **gavriloaie ionel claudiu** <ionelclaudiu@yahoo.com> To: Yahoo

Date: Wed, Apr 1, 2020 at 2:07 AM

Dear Professor Pratasik,

Thank you very much for your message. We do not know any greater expert in the coral field than Professor Lalamentik. So, of course we will take into account his comments.

However, my question regarding the online databasis is still available. How much trust to put on them?

I am working on editing the final form of Professor Lalamentik's paper. As soon as I have it done I will send it to you for the approval.
Thank you very much!

Cordially yours,
Claudiu G.

On Tuesday, March 31, 2020, 5:31:29 AM GMT+3, Yahoo <spjong07@yahoo.com> wrote:

Dear Dr. Gavrioloaie,
I just talked to Mr. Lalamentik about the new name of corals. He greatly appreciate your comment, but he has not got the new book concerning it and hope to have it someday. Therefore, if you/reviewers think that the name should be changed, he does not mind. He himself is not confident to change it because of the reference constraints. Thank you

Taskbar: search, File Explorer, Firefox, VLC, Mail, Chrome, Word

006-urmad1 - s33ng07@yahoo.com

https://mail.yahoo.com/0/0kden/2/messages/ANkwaR0NF4E/3cP1EAGyBTJEU7gvoe_nf8mei-w4R0kHM6Uy5383ouZ29uZ2xLmN46538gura_referrer_sig=AD4AAAEF0Z8k...

HOME MAIL NEWS FINANCE SPORTS ENTERTAINMENT LIFE SHOPPING YAHOO PLUS MORE... y!mail+ Upgrade Now

yahoo/mail Find messages, documents, photos or people

Compose

Intro 2/3
Unread
Starred
Drafts
Sent
Archive
Spam
Trash
▲ Last
Views show
Folders hide
+ New folder
junk

← Back → Archive Move Delete Spam ...

Re: revised paper Yahoo/Sent

B Yahoo <s33ng07@yahoo.com>
To: gavinloze jonel claudu Wed, Apr 1, 2022 at 9:33 AM

Dear Dr. Gavriloski,
I am very sorry for the mistakes. I have checked WORMS and I am working now on all the corrections. I also need to rewrite the paper and will send it back to you today. Thank you.

Sincerely Yours,
Sivester B. Pratasik

On Wednesday, April 1, 2022, 02:07:36 AM GMT+8, gavriloski jonel claudu <jonelclaudu@yahoo.com> wrote:

Dear Professor Pratasik,
Thank you very much for your message. We do not know any greater expert in the coral field than Professor Lalamentik. So, of course we will take into account his comments.


However, my question regarding the online databasis is still available. How much trust to put on them?

I am working on editing the final form of Professor Lalamentik's paper. As soon as I have it done I will send it to you for the approval.
Thank you very much!

一键畅游全球网络, 上网必用

Type here to search

28°C Hujan ringan 6:42 PM 4/14/2022



Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate

Laurentius T. X. Lalamentik, Rene Ch. Kepel, Lawrence J. L. Lumingas, Unstain N. W. J. Rembet, Silvester B. Pratasik, Desy M. H. Mantiri

Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia. Corresponding author: L. T. X. Lalamentik, ottylalamentik@yahoo.co.id

Abstract. A study on colonization of Faviidae corals on the agricultural waste-materialized artificial substrate was conducted in Selat Besar, Ratatotok district, southeast Minahasa regency, North Sulawesi. Nine artificial substrates moduls made of mixture of cement, sand, padi husk, and bamboo were placed for about 5 years on the sea bottom of Selat Besar waters. All corals of family Faviidae found on the artificial substrate were collected. Results showed that Faviidae corals could live and develop on those substrates. Fifteen species of 5 genera of Faviidae, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennetae*, *Echinopora gemmacea*, *E. lammelosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. senensis* were recorded in the present study. Mean number of colonies of Faviidae corals was 3 col.mod⁻¹, while mean diameter of the corals attached on the artificial substrate was 7.45 cm long. The distribution pattern of Faviidae corals was clumped. The diversity of Faviidae corals on the artificial substrate was low ($H' = 0.677 < 1$). The dominance index showed no dominant species ($D = 0.23$). In addition, the artificial substrate modul in this study could become an alternative technique to rehabilitate the degraded coral reefs.

Key Words: diversity, dominance, distribution pattern, Selat Besar.

Introduction. Corals are animals belonging to phylum Cnidaria (Reid et al 2009), while coral reef is an ecosystem built by calcium carbonate-producing marine biota, especially coral animals, together with other biota living on the sea bottom or water column, such as mollusks, crustaceans, echinoderms, porifera, tunicates, and other biota free-living in surrounding waters including plankton and fishes (Giyanto et al 2017; Lalamentik 1995). According to Barus (2013), coral reef is one of the unique communities that are entirely formed from biological activities. It is one of the main coastal and marine ecosystems that possess the highest productivity and biodiversity so that it is often called as tropical marine forest.

Coral Triangle (CTI) is known as center of world marine biodiversity with the highest coral diversity, 76% of total coral species. The region inside the ecological boundary of Coral Triangle covers nearly 73,000 km² of coral reefs (29% of total world coral reef area) distributed in six countries, Indonesia, Malaysia, Filipina, Timor Leste, Papua New Guinea and Solomon Islands (Burke et al 2012). As part of the CTI, Indonesia is an archipelagic country with 16,056 islands, 1,922,570 km² terrestrial area and 3,257,483 km² water in the west Indo-Pacific (BIG 2016). Indonesia that is located in the coral triangle area is recognized as one of the countries with high biodiversity consisting of more than 80 coral genera and 596 species, particularly in North Sulawesi waters that has been recorded more than 80 genera (Suharsono 2008).

Coral taxonomic and distribution studies in North Sulawesi were carried out by several scientists (Lalamentik 1998; Halidu et al 2016; Nasaru et al 2017; Suleman et al 2017; Lalamentik & Rembet 2018). So far, there is no study on corals of family Faviidae living and growing on the artificial substrate, especially in North Sulawesi waters. Hence, present study focuses on Faviidae coral colonies on the artificial substrate with the

objectives of identifying the coral species, analysing the density and the diameter of the coral colony, the ecological index, and assessing the distribution patterns. This finding is expected to be one of the references in coral development studies, concerning Faviidae coral colonization on the artificial substrate. Besides, it could provide alternative artificial substrates in coral reef rehabilitation program.

Material and Method

Study site. This work was accomplished in Selat Besar waters, Ratatotok district, southeast Minahasa regency (Figure 1), North Sulawesi. Selat Besar is flanked by Totok bay and Moluccas Sea. The study site was laid on 0°51'34.945''S and 124°44'14.416'' E (Figure 1) using Global Position System (GPS).




Figure 1. Sampling locality.

Artificial substrate construction. In this study, 9 artificial substrate moduls were set on the sea bottom, approximately 7-8 m depth. Sheehy (1986) stated that one of the major roles of artificial substrate placement in the water is to create new habitats similar to artificial reefs to be inhabited by target species. The use of artificial substrate as coral new habitat had been conducted in Singaporean waters in order to restore the coral reefs (Ng & Chou 2017). Each modul consists of 10 concrete blocks made of mixture of padi husk, sand, and cement. The block frames were made of bamboo. The use of bamboo in concrete construction has been done before as mentioned in Glenn (1950) and Mehra & Ghosh (1965). One of the properties that would make bamboo a good substitute to steel in reinforced concrete is its strength (Varma, 2017). The block had a dimension of 100 cm leng, 15 cm width, and 15 cm height. The blocks were set in such a manner to yield a modul. Each modul is estimated to have an area of 4.5 m². These 9 moduls were placed since June 2014. Sample collections were carried out in July 2019 (after 5 years of

deployment). The period of 5 years is determined because the massive coral growth is slow. This duration is considered enough to see whether there is Faviidae coral capable of adhering and growing on the artificial substrate.

Sample collection and handling procedures. All massive coral species found to attach on the modul substrate were taken using a chisel and hammer. The samples were then put into labelled bag as the modul number. All corals were collected by using SCUBA equipments. All activities were documented by using an underwater camera.

Before species identification, the coral specimens were soaked in bleach-containing freshwater for 24 h in order to kill the coral and to prevent the presence of fungi. The corals were then cleaned under the running water to remove the attached dirt so that the clean coral and the bleaching coral skeleton were obtained. The corals were placed under direct sunlight to get dry to ease the identification.

Species identification. Coral identification was done following Veron (1986; 2000), Suharsono (2008), and Veron et al (1977). The specimens of Faviidae corals were examined through morphological observations under the SZ51 Olympus microscope on the characteristics of paliform lobe, septa cleats, coralite type, number of first septa, second septa, and third septa. The measurements of width of paliform lobe, coralite size, and calix size used a multitoyo digital caliper.

Data analysis. Faviidae coral colonization data were directed to ecological index analyses as follows:

Dominance index. This index was employed to know the dominant species in data sampling boundary. The present study applied Simpson dominance index (Krebs, 1989):

$$D = \frac{\sum_{i=1}^s [n_i]^2}{[N]^2} \dots\dots\dots (1)$$

Where D = dominance index, n_i = number of i individuals, and N = total number of individuals.

Dominance index ranges between 0 and 1, if D approaches to 0, it indicates no dominance and the community structure is in stable condition, and if D approaches to 1, it indicates the presence of species dominance.

Density. The population density was calculated using Krebs (1989) as follows

$$K = \frac{N_i}{A} \dots\dots\dots (2)$$

Where, K = density (ind.modul⁻¹), N_i = number of individuals, and A = number of moduls where samples were collected.

Diversity. The diversity index of genus *Favia* Oken 1815 was calculated using Shannon-Wiener equation (Krebs 1989):

$$H = -\sum_{i=1}^s \frac{n_i}{N} \left(\frac{n_i}{N} \right) \dots\dots\dots (3)$$

Where H = Shannon-Wiener diversity index, n_i = number of individuals i , N = total number of individuals, and s = number of genera.

Distribution pattern. The distribution pattern of Faviidae corals were estimated using Morisita index (Krebs 1989):

$$I_d = \frac{\sum_{i=1}^q n_i (n_i - 1)}{N (N - 1)} \dots\dots\dots (4)$$

Where I_d = Morisita index, n_i = number individuals in each plot, n = total number of individuals in all plots, and N = number of plots, in which $I\delta = 1$ is categorized as random distribution, $I\delta > 1$ as clumped, and $I\delta < 1$ as uniform.

Results and Discussion

Water conditions of Selat Besar. Water quality is crucial for colonization of macroinvertebrates (Orwa et al 2018). Water temperature is one of the limiting factors for aquatic organisms to live, one of which is coral. Field measurements found that water temperature in the study site ranged from 29° - 30°C. This range is still suitable for coral development. According to Kurniawan (2011), corals, in general, can live at the temperature above 18°C, while growth, they require an ideal water temperature, between 25°-32°C. Most coral reefs exist where salinity is stable and where average salinity is that of normal seawater 34-36‰ (Sheppard et al., 2017). Water salinity in the study site was between the range of 30-31‰. Guntur (2011) found that water salinity for ideal coral growth in Indonesia ranged from 29 to 36‰. Therefore, water salinity in Selat Besar highly supports the coral to grow well. Light intensity does not influence larval settlement, and their response to light intensity may be species-specific (Mundy & Babcock 1998). This study was done in shallow water, about 8 m deep, so that light intensity is not a limiting factor for coral settlement. The study did not also focus on the coral settlement in relation with the light intensity.

Corals of family Faviidae recorded on the artificial substrate. Physical and biological factors are important for recruitment success (Doropoulos et al 2016), As many as 15 coral species of 5 genera were recorded in this study. *Favia* and *Favites* had the highest number of species, each with 4 species, followed by genus *Goniastrea* with 3 species, then *Echinopora* and *Platygyra*, 2 species, respectively. Table 1 demonstrates the coral species of Faviidae encountered on the modul substrate with mean number of colonies of 3.00 per modul. Even though these numbers are less than those reported by Kilfoyle et al (2008), 13 col.mod.⁻¹, the colonies comprised several coral families. The coral genera of Faviidae inhabiting the artificial substrate apparently come the coral reef around Selat Besar waters. It is reasonable, since Faviidae is coral family generally found in Selat Besar waters (Lalamentik 1996;1997;1998).

Commented [indra1]: it is Table 1, not 02, since this is the first table in the paper

Commented [C2R1]: yes, table 1

Table 1

Faviidae coral species encountered living on the artificial substrate modul

No	Species
1	<i>Echinopora gemmacea</i> (Lamarck, 1816)
2	<i>E. lammelosa</i> (Esper, 1795)
3	<i>Favia laxa</i> (Klunzinger, 1879)
4	<i>F. matthai</i> (Vaughan, 1918)
5	<i>F. palida</i> (Dana, 1846)
6	<i>F. stelligera</i> (Dana, 1846)
7	<i>Favites bennetiae</i> (first record for Central Indonesia)
8	<i>F. complanata</i> (Ehrenberg, 1734)
9	<i>F. pentagona</i> (Esper, 1794)
10	<i>F. russelli</i> (Wells, 1954)
11	<i>Goniastrea aspera</i> (Verrill, 1864)
12	<i>G. favulus</i> (Dana, 1846)
13	<i>G. pectinata</i> (Ehrenberg, 1834)
14	<i>Platygyra daedalea</i> (Elis & Solander, 1786)
15	<i>P. sinensis</i> (Edwards & Haime, 1857)

Favites bennetiae recorded in the present study is the first record found in the coral reef of Central Indonesian waters. This species was previously mostly found in the northeast of Australian waters (Veron et al 1977). Moreover, genus *Favia* was found having the highest mean number of colonies, 1.11 col.mod⁻¹, followed by *Goniastrea*, 0.67 col.mod⁻¹, and the lowest was found in *Platygyra*, 0.22 col.mod⁻¹ (Figure 2). Variations in number of Faviidae colonies per modul are dependent upon the abundance of larvae that survive and settle on the substrate. Faviidae tends to have good adaptability to occupying the available artificial substrates. Bachtiar et al (2011) found that Faviidae is one of the coral families capable of surviving in the reef ball deployed on the sea bottom.




Figure 2. Mean no. colonies of *Faviidae* per modul.

Mean diameter of the coral colony on the artificial substrate was 7.45 cm with the largest recorded in *Favia laxa*, 10.5 cm and the smallest in *F. palida*, 3 cm. In comparison among genera, mean diameter of Faviidae corals attached on the artificial substrate ranged from 5.88 to 8.50 cm. Genus *Platygyra* has the largest mean diameter, 8.50 cm, and *Echinopora* had the smallest one, 5.88 cm (Figure 3). Variations in colony size could result from different sequence of planular settlements. This condition, according to Bachtiar et al (2011), indicates that coral colonization on the artificial substrate sustainably occurs.




Figure 3. Mean diameter of *Faviidae* corals living on the artificial substrate.

Ecological index. The present study measured dominance index (D) and diversity index (H') of Faviidae corals thriving on the artificial substrate. The data are given in Table 2.

Table 2

Ecological index of Faviidae corals

No	Ecological index	Value	Remarks
1	Dominance Index (D)	0.23	No dominance
2	Distribution pattern (I δ)	1.79	Clumped
3	Diversity index (H')	3.64	high

The dominance index of Faviidae corals is 0.23 (Table 3). It means that no species outnumbers the others. Selat Besar waters is still capable of supporting the coral life, and there is no competition that could make certain species be dominant (Muqsit et al 2016). The diversity index is 3.64. This value is supported by low species dominance index, in which there were 15 Faviidae species found. These numbers are high enough due to small study area cover. This finding is in agreement with Rondo et al (2014) that high diversity will cause low species dominance.

Distribution pattern analysis found that Faviidae in the study site had clumped distribution (Id = 1.79 > 1). This pattern could result from the fact that there were four moduls not occupied by Faviidae corals. In Indonesia, studies on coral colonization was also done by Razak (2008) using Eco-reef modul in Manado bay waters, North Sulawesi, and found that the artificial substrate used are, in fact, inhabited by hard corals. Hence, the artificial substrate could be recommended as an alternative in coral reef rehabilitation program.

Conclusions. Fifteen coral species were found settling on the agricultural waste-materialized artificial substrate, *Favia pallida*, *F. laxa*, *F. stelligera*, *F. matthaii*, *Favites pentagona*, *F. russelli*, *F. complanata*, *F. bennetiae*, *Echinopora gemmacea*, *E. lammelosa*, *Goniastrea aspera*, *G. favulus*, *G. pectinata*, *Platygyra daedalea*, and *P. senensis*. They had mean 3.00 col.mod⁻¹ with mean diameter of 7.45 cm. The ecological index also indicated that the available artificial substrate could support various coral species to grow. In this study, Faviidae corals growing on the available artificial substrate have reflected clumped distribution. Thus, this type of modul could be used for coral reef rehabilitation.

References

- Bachtiar I., Prayogo W., Sukri A., 2011 Coral colonization and growth reef ball modules at the Benete Bay, Sumbawa Island, Indonesia. Manado. P. 31-40.
- Doropoulos C., Roff G., Bozec Y. M., Zupan M., Werminghausen J., Mumby P. J., 2016 Characterizing the ecological trade-offs throughout the early ontogeny of coral recruitment. *Ecol. Monogr.* 86: 20-44.
- Guntur, 2011 [Coral ecology in the artificial reef]. Ghalia Indonesia. Semarang. 139 pp. [in Indonesian]
- Halidu, A., Lalamentik L. T. X., Rembet U. N. W. J., 2016 [Hard coral distribution in the reef flat of the south coast of Putus-Putus island, east Ratatotok, Ratatotok district, southeast Minahasa]. *Jurnal Ilmiah Platax* 4(1):19-30. [in Indonesian]
- Kilfoyle K., Rangel M. A., Dodge R. E., Spieler R. E., 2008 Coral Reef Restoration: Standardized Module Intervention and monitoring program in Mexico, Preliminary Results. Poster presentation material, 11 Int Coral Reef Sym Florida, Session number XXIV, p. 541
- Krebs C. J., 1989 Ecological Methodology. Harper Collins Publishers. New York. 654 pp.
- Lalamentik L. Th. X., 1995 [Study on coral reef potential in Tombasian district, Minahasa, North Sulawesi] Fakultas Perikanan dan Ilmu Kelautan UNSRAT. Manado. 28 pp. [in Indonesian] Unpublished

Commented [indra3]: edit the list following the journal's format

- Lalamentik L. T. X., 1996 Coral Reef conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1996). 23 pp. Unpublished.
- Lalamentik L. T. X., 1997 Coral Reef Conditions around the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study in Ratatotok and Adjacent Waters, the District of Minahasa North Sulawesi. (Survey Dated October, 1997). 22 pp. Unpublished.
- Lalamentik L. Th. X., 1998 Coral reef Condition Arround the Gold Mining Area of PT. Newmont Minahasa Raya: A Monitoring Study for May 1998 Data in Ratatotok and Adjacent Waters, The District of Minahasa North Sulawesi. Fakultas Perikanan dan Ilmu Kelautan USNRAT. Manado. 23 pp. Unpublished.
- Lalamentik, L. Th. X., Rembet, U. N. J., 2018 [Study on the Taxonomy of Genus Favia (Oken, 1815) at the Reef Flats of Kampung Ambong Village in Likupang Timur District, North Minahasa]. JURNAL ILMIAH PLATAX, 6(1), 188-193. [in Indonesian]
- Mehra S. R., Ghosh R. G., 1965 "Bamboo-reinforced soil-cement," Civil Engineering and Public Works Review, Vol. 60, no. 711, October; Vol. 60, no. 712. November 1965.
- Mundy C. N., Babcock R., 1998 Role of light intensity and spectral quality in coral settleError! Hyperlink reference not valid.ment: implications for depth-dependent settlement? J. Exp. Mar. Biol. Ecol. 223 (2): 235-255.
- Muqsit A., Purnama D., Ta'alidin Z., 2016 [The community structure of coral reef in Pulau Dua, Enggano district, North Bengkulu regency.] Jurnal Enggano Vol. 1, No. 1 (75-78). Ilmu Kelautan Fakultas Pertanian Universitas Bengkulu. Bengkulu. [in Indonesian]
- Nasaru J. H., Lalamentik L. Th. X., Rembet U. N. W. J., 2017 [Pocillopora verrucosa (Ellis and Solander, 1786) distribution in the reef flat of the south coast of Putus-Putus island, east Ratatotok village, southeast Minahasa regency]. Jurnal Ilmiah Platax 5(1): 96-103 pp. Universitas Sam Ratulangi Manado. [in Indonesian]
- Ng C. S. L., Chou L. M., 2017 Coral reef restoration in Singapore-past, present and future. Sustainability Matters: Environmental Management in the Anthropocene. Singapore: World Scientific, 3-23 pp.
- Orwa P. O., Omondi R., Chemoiwa E. J., 2018 Colonization patterns of benthic macroinvertebrates in fertilized and non-fertilized earthen fish ponds. Int J Aquac Fish Sci 4(3): 022-026.
- Razak T., 2008 The population of hard coral colonies growing on Ecoreef artificial modules on Manado Tua Island, Bunaken National Park, North Sulawesi, Indonesia. Oral presentation material, 11 Int Coral Reef Sym Floria, Session number XXIV, p. 223.
- Reid C., Marshall J., Logan D., Kleine D., 2009 [Coral reef and climate change]. CoralWatch, The University of Queensland. Australia. 256 pp. [in Indonesian]
- Ricardo G. F., Jones R. J., Nordborg M., Negri A. P., 2017 Settlement patterns of the coral Acropora millepora on sediment-laden surfaces. Science of the Total Environment. 609:277-288.
- Rondo M., Tamanampo J. F. W. S., 2014 Principles of Aquatic Ecology. Fakultas Perikanan dan Ilmu Kelautan, Universitas Sam Ratulangi. Manado. 427 pp.
- Sheppard C., Davy S., Pilling G., Graham N., 2017 The biology of coral reefs. Oxford University Press. 365 pp.
- Sheehy D. J., 1986 New approaches in artificial reef design and applications. p. 256-263. In F. M. D. Trie (ed.) Artificial reefs: marine and freshwater applications. Lewis Publishers Inc., Michigan, USA.
- Suharsono, 2008 [Coral species in Indonesia]. Lembaga Ilmu Pengetahuan (LIPI) : COREMAP Program. Jakarta. 116 p. [in Indonesian]
- Suleman Y., Lalamentik L. Th. X., Rembet U. N. W. J., 2017 [Hard coral Favites abdita (Ellis and Solander, 1786) distribution in the reef flat of Malalayan 2, Malalayang district, Manado city]. Jurnal Ilmiah Platax 5(1): 104-111. Universitas Sam Ratulangi. Manado. [Indonesian] pp [Indonesian]

- Varma M. B., 2017 Properties of cement concrete reinforced with bamboo-strip-Mat. IOSR, Journal of Mechanical and Civil Engineering (IOSR-JMCE), eISSN, 2278-1684: 47-59.
- Veron J. E. N., 1986 Corals of Australia and the Indo-Pacific. Angus and Robertson Publishers. Australia. 644 pp.
- Veron J. E. N., 2000 Corals of the World Volume 1. Australian Institute of Marine Science and CRR Qld Pty Ltd. Australia 463 pp.
- Veron J. E. N., Pichon M., Wijsman-Best M., 1977 Scleractinia of Eastern Australia. Canberra. Vol. 3 (Part 2). 233 pp.

Authors:

- L.T.X. Lalamentik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: ottylalamentik@yahoo.co.id
- R. Ch. Kepel, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: renecharleskepel65@gmail.com
- L.J.L. Lumingas, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: ljlumingas@yahoo.com
- U.N.W.J. Rembet Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: onlyrembet@unsrat.ac.id
- S.B. Pratasik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, email: spjong07@yahoo.com
- D.M.H. Mantiri, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado-95115, North Sulawesi, Indonesia, email: dmh_mantiri@unsrat.ac.id

Find messages, documents, photos or people

← Back ↶ ↷ → Archive Move Delete Spam ...

Re: revised paper Yahoo/Inbox

 **gavriloaie ionel claudiu** <ionelclaudiu@yahoo.com> To: Yahoo Thu, Apr 2, 2020 at 6:06 PM

Dear Professor Pratasik,

I received the revised version of the paper. More soon.
Thank you!

Sincerely yours,
Claudiu G.

On Thursday, April 2, 2020, 5:38:13 AM GMT+3, Yahoo <spjong07@yahoo.com> wrote:

Dear Dr. Gavriloaie,
I herewith send you the revised paper of Lalamentik et al. I put already WoRMs for name validation, so that the number of genera becomes 8. Some data are also added in Table 1. Data in Figure also change. These make the whole document has to be written. I put all changes in yellow. I hope this version has already met your requirements.

Sincerely Yours,
Silvester B. Pratasik


On Wednesday, April 1, 2020, 11:50:22 PM GMT+8, gavriloaie ionel claudiu <ionelclaudiu@yahoo.com> wrote:

Dear Professor Pratasik,

Find messages, documents, photos or people

← Back ↶ ↷ → Archive Move Delete Spam

Re: the final form of the paper of Lalamentik et al's - Faviidae coral colonization ... Yahoo/Inbox

 **gavriloaie ionel claudiu** <ionelclaudiu@yahoo.com>
To: Yahoo Sun, Apr 19, 2020 at 7:12 PM

Dear Professor Pratasik,

I have uploaded the paper on the journal's website: <http://www.bioflux.com.ro/home/volume-13-2-2020/>.
Thank you very much for another kind and good cooperation with you!

Sincerely yours,
Claudiu G.

On Sunday, April 19, 2020, 1:48:17 AM GMT+3, Yahoo <spjong07@yahoo.com> wrote:

Dear Dr. Gavriiloaie,
I agree with you to have only English translation. Therefore, I would greatly appreciate if you could erase the original Indonesian titles in the references list.

Sincerely Yours,
Silvester B. Pratasik

2020.910-918.doc



Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate

Laurentius T. X. Lalamentik, Rene C. Kepel, Lawrence J. L. Lumingas, Unstain N. W. J. Rembet, Silvester B. Pratasik, Desy M. H. Mantiri

Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia. Corresponding author: L. T. X. Lalamentik, ottylalamentik@yahoo.co.id

Abstract. A study on colonization of Faviidae corals on the agricultural waste-materialized artificial substrate was conducted in Selat Besar, Rataotok district, southeast Minahasa regency, North Sulawesi. Nine artificial substrates modules made of mixture of cement, sand, padi husk, and bamboo were placed for about 5 years on the sea bottom of Selat Besar waters. All corals of family Faviidae found on the artificial substrate were collected. Results showed that Faviidae corals could live and develop on those substrates. Fifteen species of 8 genera of family Faviidae were recorded in the present study, *Dipsastraea pallida*, *D. laxa*, *D. matthaii*, *Favites pentagona*, *F. complanata*, *Paragoniastrea russelli*, *Oulophyllia bennettiae*, *Echinopora gemmacea*, *E. lamellosa*, *Goniastrea stelligera*, *G. favulus*, *G. pectinata*, *Coelastrea aspera*, *Platygyra daedalea*, and *P. sinensis*. Mean number of colonies of Faviidae corals was 3 col mod⁻¹, while mean diameter of the corals attached on the artificial substrate was 2.35 cm long. The distribution pattern of Faviidae corals was uniform. The diversity of Faviidae corals on the artificial substrate was low ($H' = 2.568$). The dominance index showed no dominant species ($D = 0.089$). In addition, the artificial substrate module in this study could become an alternative technique to rehabilitate the degraded coral reefs.

Key Words: diversity, dominance, distribution pattern, Selat Besar.

Introduction. Corals are animals belonging to phylum Cnidaria (Reid et al 2009), while coral reef is an ecosystem built by calcium carbonate-producing marine biota, especially coral animals, together with other biota living on the sea bottom or water column, such as mollusks, crustaceans, echinoderms, porifera, tunicates, and other biota free-living in surrounding waters including plankton and fishes (Lalamentik 1995; Giyanto et al 2017). According to Barus et al (2018), coral reef is one of the unique communities that are entirely formed from biological activities. It is one of the main coastal and marine ecosystems that possess the highest productivity and biodiversity so that it is often called as tropical marine forest.

Coral Triangle (CTI) is known as center of world marine biodiversity with the highest coral diversity, 76% of total coral species. The region inside the ecological boundary of CTI covers nearly 73,000 km² of coral reefs (29% of total world coral reef area) distributed in six countries, Indonesia, Malaysia, Philippines, Timor Leste, Papua New Guinea and Solomon Islands (Burke et al 2012). As part of the CTI, Indonesia is an archipelagic country with 16,056 islands, 1,922,570 km² terrestrial area and 3,257,483 km² water in the west Indo-Pacific (Briney 2020). Indonesia that is located in the CTI area is recognized as one of the countries with high biodiversity consisting of more than 80 coral genera and 596 species, particularly in North Sulawesi waters where have been recorded more than 80 genera (Suharsono 2008).

Coral taxonomic and distribution studies in North Sulawesi were carried out by several scientists (Lalamentik 1998; Halidu et al 2016; Nasaru et al 2017; Suleman et al 2017; Reskiwati et al 2018). So far, there is no study on corals of family Faviidae living and growing on the artificial substrate, especially in North Sulawesi waters. Hence, the

present study focuses on Faviidae coral colonies on the artificial substrate with the objectives of identifying the coral species, analysing the density and the diameter of the coral colony, the ecological index, and assessing the distribution patterns. This finding is expected to be one of the references in coral development studies, concerning Faviidae coral colonization on the artificial substrate. Besides, it could provide alternative artificial substrates in coral reef rehabilitation program.

Material and Method

Study site. This work was accomplished in Selat Besar waters, Ratatotok district, southeast Minahasa regency (Figure 1), North Sulawesi. Selat Besar is flanked by Totok bay and Moluccas Sea. The study site was laid on $0^{\circ}51'34.945''\text{S}$ and $124^{\circ}44'14.416''\text{E}$ (Figure 1) using Global Position System (GPS).




Figure 1. Sampling locality.

Artificial substrate construction. In this study, 9 artificial substrate modules were set on the sea bottom, approximately 7-8 m depth (Figure 2). Sheehy (1986) stated that one of the major roles of artificial substrate placement in the water is to create new habitats similar to natural reefs to be inhabited by target species. The use of artificial substrate as coral new habitat had been conducted in Singaporean waters in order to restore the coral reefs (Ng & Chou 2017). Each module consists of 10 concrete blocks made of mixture of padi husk, sand, and cement. The block frames were made of bamboo. The use of bamboo in concrete construction has been done before as mentioned in Glenn (1950) and Mehra & Ghosh (1965). One of the properties that would make bamboo a good substitute to steel in reinforced concrete is its strength (Varma 2017). The block had a dimension of 100 cm length, 15 cm width, and 15 cm height. The blocks were set in such a manner to yield a module. Each module is estimated to have an area of 4.5 m^2 . These 9 modules

were placed since June 2014. Sample collections were carried out in July 2019 (after 5 years of deployment). The period of 5 years is determined because the massive coral growth is slow. This duration is considered enough to see whether there is Faviidae coral capable of adhering and growing on the artificial substrate.




Figure 2. Artificial substrate modules. Left: new module; Right: colonized module.

Sample collection and handling procedures. All massive coral species found to attach on the module substrate were taken using a chisel and hammer. The samples were then put into labelled bag as the module number. All corals were collected by using SCUBA equipments. All activities were documented by using an underwater camera.

Before species identification, the coral specimens were soaked in bleach-containing freshwater for 24 h in order to kill the coral and to prevent the presence of fungi. The corals were then cleaned under the running water to remove the attached dirt so that the clean coral and the bleaching coral skeleton were obtained. The corals were placed under direct sunlight to get dry to ease the identification.

Species identification. Coral identification was done following Veron (1986), Veron & Stafford-Smith (2000), Suharsono (2008), and Veron et al (1977). The specimens of Faviidae corals were examined through morphological observations under the SZ51 Olympus microscope on the characteristics of paliform lobe, septa cleats, coralite type, number of first septa, second septa, and third septa. The measurements of width of paliform lobe, coralite size, and calix size used an LCD electronic digital gauge stainless Vernier caliper 150 mm. World Register database of Marine Species (WoRMS) was utilized for scientific name validation (Hoeksema & Cairns 2020).

Data analysis. Faviidae coral colonization data were directed to ecological index analyses as follows:

Dominance index. This index was employed to know the dominant species in data sampling boundary. The present study applied Simpson dominance index (Krebs 1989):

$$D = \frac{\sum_{i=1}^s \left[\frac{n_i}{N} \right]^2}{\dots \dots \dots} \dots \dots \dots (1)$$

Where D = dominance index, n_i = number of i individuals, and N = total number of individuals.

Dominance index ranges between 0 and 1, if D approaches to 0, it indicates no dominance and the community structure is in stable condition, and if D approaches to 1, it indicates the presence of species dominance.

Density. The population density was calculated using Krebs (1989) as follows:

$$K = \frac{\sum Ni}{A} \dots\dots\dots (2)$$

Where, K = density (ind module⁻¹), Ni = number of individuals, and A = number of modules where samples were collected.

Diversity. The diversity index of genus *Favia* Oken, 1815 was calculated using Shannon-Wiener equation (Krebs 1989):

$$H = \sum_{i=1}^s \frac{ni}{N} \left(\frac{\ln ni}{N} \right) \dots\dots\dots (3)$$

Where H = Shannon-Wiener diversity index, ni = number of individuals *i*, N = total number of individuals, and s = number of genera.

Distribution pattern. The distribution pattern of Faviidae corals was estimated using Morisita index (Krebs 1989):

$$Id = \frac{\sum_{i=1}^q ni (ni - 1)}{N(N - 1)} \dots\dots\dots (4)$$

Where Id = Morisita index, ni = number of individuals in each plot, and N = total number of individuals in all plots, in which Id = 1 is categorized as random distribution, Id > 1 as clumped, and Id < 1 as uniform.

Results and Discussion

Water conditions of Selat Besar. Water quality is crucial for colonization of macroinvertebrates (Orwa et al 2018). Water temperature is one of the limiting factors for aquatic organisms to live, one of which is coral. Field measurements found that water temperature in the study site ranged from 29 to 30°C. This range is still suitable for coral development. According to Reid et al (2009), corals, in general, can live at water temperature from 18 to 30°C, while for growth, they require an ideal water temperature from 27 to 29°C (Giyanto et al 2017). Most coral reefs exist where salinity is stable and where average salinity is that of normal seawater 34-36‰ (Sheppard et al 2017). Water salinity in the study site was in the range of 30-31‰. Guntur (2011) found that water salinity for ideal coral growth in Indonesia ranged from 29 to 36‰. Therefore, water salinity in Selat Besar highly supports the coral to grow well. Light intensity does not influence larval settlement, and their response to light intensity may be species-specific (Mundy & Babcock 1998).

Corals of family Faviidae recorded on the artificial substrate. Physical and biological factors are important for recruitment success (Doropoulos et al 2016). As many as 15 coral species of 8 genera were recorded in this study (Table 1). *Dipsastraea* and *Goniastrea* had the highest number of species, each with 3 species, followed by genus *Favites*, *Echinopora*, and *Platygyra* with 2 species each, while *Oulophyllia*, *Paragoniastrea*, and *Coelastrea* were only represented by one species. Mean density of Faviidae corals on the module substrate was 3.00 col mod⁻¹. Even though these numbers are less than those reported by Kilfoyle et al (2008), 13 col mod⁻¹, the colonies comprised several coral genera. The coral genera of Faviidae inhabiting the artificial substrate apparently come from the coral reef around Selat Besar waters. It is reasonable, since Faviidae is coral family generally found in Selat Besar waters (Lalamentik 1996, 1997, 1998).

Oulophyllia bennettiae recorded in the present study is the first record found in the coral reef of Central Indonesian waters. This species was previously mostly found in the northeast of Australian waters (Veron et al 1977). Moreover, genus *Dipsastraea* was found having the highest mean number of colonies, 0.889 col mod⁻¹, followed by

Goniastrea, 0.667 col mod⁻¹, and the lowest was found in *Oulophyllia* and *Paragoniastrea*, 0.111 col mod⁻¹ (Figure 3).

Table 1

Faviidae coral species encountered living on the artificial substrate module

No	Species	Mean diameter (cm)
1	<i>Echinopora gemmacea</i> (Lamarck, 1816)	1.361
2	<i>E. lamellosa</i> (Esper, 1795)	0.556
3	<i>Dipsastraea laxa</i> (Klunzinger, 1879)	2.389
4	<i>D. matthaii</i> (Vaughan, 1918)	1.056
5	<i>D. pallida</i> (Dana, 1846)	2.000
6	* <i>Oulophyllia bennettiae</i> (Veron, Pichon & Wijsman-Best, 1977)	0.778
7	<i>Favites complanata</i> (Ehrenberg, 1834)	1.056
8	<i>F. pentagona</i> (Esper, 1795)	0.694
9	<i>Paragoniastrea russelli</i> (Wells, 1954)	0.944
10	<i>Coelastrea aspera</i> (Verrill, 1866)	1.556
11	<i>Goniastrea favulus</i> (Dana, 1846)	0.556
12	<i>G. pectinata</i> (Ehrenberg, 1834)	0.972
13	<i>G. stelligera</i> (Dana, 1846)	0.667
14	<i>Platygyra daedalea</i> (Elis & Solander, 1786)	1.222
15	<i>P. sinensis</i> (Milne Edwards & Haime, 1849)	0.667

* The first record for Central Indonesia.

Variations in number of Faviidae colonies per module are dependent upon the abundance of larvae that survive and settle on the substrate. Faviidae tends to have good adaptability to occupying the available artificial substrates. Bachtiar & Prayogo (2010) found that Faviidae is one of the coral families capable of surviving in the reef ball deployed on the sea bottom as well.




Figure 3. Mean number of colonies of *Faviidae* per module.

Mean diameter of the coral colony on the artificial substrate was 2.35 cm with the largest recorded in *Dipsastraea*, 5.444 cm, and the smallest in *Oulophyllia*, 0.778 cm (Figure 4). In species level, the largest mean diameter was recorded in *D. laxa* and the lowest in *E. lamellosa* and *G. favulus*, respectively (Table 1). Variations in colony size could result from different sequence of planula settlements and available space on the module. This

condition, according to Bachtiar & Prayogo (2010), indicates that coral colonization on the artificial substrate sustainably occurs.




Figure 4. Mean diameter of Faviidae corals living on the artificial substrate.

Ecological index. The present study measured dominance index (D) and diversity index (H') of Faviidae corals thriving on the artificial substrate. The data are given in Table 2.

Table 2

Ecological index of Faviidae corals

No	Ecological index	Value	Remarks
1	Dominance Index (D)	0.089	No dominance
2	Distribution pattern (Id)	0.49	uniform
3	Diversity index (H')	2.568	high

The dominance index of Faviidae corals is 0.089 (Table 2). It means that no species outnumbers the others. Selat Besar waters is still capable of supporting the coral life, and there is no competition that could make certain species be dominant (Muqsit et al 2016). The diversity index is 2.568. This value is higher than that in all study sites of natural coral reefs reported by Akbar et al (2020) in Bangka Belitung Islands reflecting that Selat Besar waters is potential for coral development. This high diversity index is also supported by low species dominance index, in which there were 15 Faviidae species found. These numbers are high enough due to small study area cover. This finding is in agreement with Rondo et al (2014) that high diversity will cause low species dominance.

Distribution pattern analysis found that Faviidae in the study site had uniform distribution ($Id = 0.49 > 1$). This pattern could result from the fact that there were four modules not occupied by Faviidae corals. In Indonesia, studies on coral colonization was also done by Razak (2008) using eco-reef module in Manado bay waters, North Sulawesi, and found that the artificial substrate used are, in fact, inhabited by hard corals. Hence, the artificial substrate could be recommended as an alternative in coral reef rehabilitation program.

Conclusions. Fifteen coral species were found settling on the agricultural waste-materialized artificial substrate, *Dipsastraea pallida*, *D. laxa*, *D. matthaii*, *Favites*

pentagona, *F. complanata*, *Oulophyllia bennettiae*, *Echinopora gemmacea*, *E. lamellosa*, *Coelastrea aspera*, *Goniastrea stelligera* *G. favulus*, *G. pectinata*, *Paragoniastrea russelli*, *Platygyra daedalea*, and *P. sinensis*. They had mean 3.00 col mod⁻¹ with mean diameter of 2.35 cm. The ecological index also indicated that the available artificial substrate could support various coral species to grow. In this study, Faviidae corals growing on the available artificial substrate have reflected uniform distribution. Thus, this type of module could be used for coral reef rehabilitation. In addition, water conditions in Selat Besar are important factors supporting the coral development.

References

- Akbar H., Syari I. A., Suyatna I., Putriningtias A., Bahri S., Destilawaty, Putra S. A., 2020 Condition of coral reefs in East Belitung, Bangka Belitung Islands, Indonesia. *AACL Bioflux* 13(1):64-70.
- Bachtiar I., Prayogo W., 2010 Coral recruitment on reef ball™ modules at the Benete Bay, Sumbawa Island, Indonesia. *Journal of Coastal Management* 13(2):119-125.
- Barus B. S., Prartono T., Soedarma D., 2018 Environmental effect on coral reefs life form in the Lampung Bay]. *Jurnal Ilmu dan Teknologi Kelautan Tropis* 10(3):699-709. [in Indonesian]
- Briney A., 2020 Geography of Indonesia: learn about the World's largest archipelago nation. Available at: <https://www.thoughtco.com/geography-of-indonesia-1435052>. Accessed: January, 2020.
- Burke L., Reyntar K., Spalding M., Perry A., 2012 Looking back the threatened coral reefs in the Coral Triangle. World Resources Institute, USA, 76 pp.
- Doropoulos C., Roff G., Bozec Y. M., Zupan M., Werminghausen J., Mumby P. J., 2016 Characterizing the ecological trade-offs throughout the early ontogeny of coral recruitment. *Ecological Monographs* 86(1):20-44.
- Giyanto, Abrar M., Hadi T. A., Budianto A., Hafizt M., Salatahpoly A., Iswari M. Y., 2017 [The status of Indonesian coral reefs 2017]. *Penelitian Oceanografi LIPI*, Jakarta, 30 pp. [in Indonesian]
- Glenn H. E., 1950 Bamboo-reinforcement in Portland cement concrete. Engineering Experiment Station, Clamson Agricultural College, Clamson, South Caroline, Bulletin no. 4, 171 pp.
- Guntur, 2011 [Coral ecology in the artificial reef]. *Ghalia Indonesia*, Semarang, 139 pp. [in Indonesian]
- Halidu A., Lalamentik L. T. X., Rembet U. N. W. J., 2016 [Distribution of coral reefs stone at the reef flat of the South coast of Putus-Putus Island, East Ratatotok, Ratatotok district, Southeast Minahasa Regency]. *Jurnal Ilmiah Platax* 4(1):19-30. [in Indonesian]
- Hoeksema B. W., Cairns S., 2020 World List of Scleractinia. *Favia matthaii* Vaughan, 1918. Accessed through: World Register of Marine Species at: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=207437> on 2020-04-01.
- Kilfoyle K., Rangel Avalos M. A., Dodge R. E., Spieler R. E., 2008 Coral reef restoration: standardized module intervention and monitoring program in Mexico, preliminary results. *Marine & Environmental Sciences Faculty Proceedings*, Presentation at the 11th International Coral Reef Symposium Florida, Lectures 395, Session number XXIV, 541 pp.
- Krebs C. J., 1989 *Ecological methodology*. Harper & Row Publishers, New York, 654 pp.
- Lalamentik L. T. X., 1995 [Study on coral reef potential in Tombasian district, Minahasa, North Sulawesi]. *Fakultas Perikanan dan Ilmu Kelautan UNSRAT*, Manado, 28 pp. [in Indonesian]
- Lalamentik L. T. X., 1996 [Coral reef conditions around the gold mining area of PT. Newmont Minahasa Raya: a monitoring study in Ratatotok and adjacent waters, the District of Minahasa North Sulawesi]. *Fakultas Perikanan dan Ilmu Kelautan. Universitas Sam Ratulangi*, Manado, Survey dated October, 1996, 23 pp. [in Indonesian]

- Lalamentik L. T. X., 1997 [Coral reef conditions around the gold mining area of PT. Newmont Minahasa Raya: a monitoring study in Ratatotok and adjacent waters, the District of Minahasa North Sulawesi]. Fakultas Perikanan dan Ilmu Kelautan. Universitas Sam Ratulangi, Manado, Survey Dated October, 1997, 22 pp. [in Indonesian]
- Lalamentik L. T. X., 1998 [Coral reef condition around the gold mining area of PT. Newmont Minahasa Raya: a monitoring study for May 1998 data in Ratatotok and adjacent waters, the District of Minahasa North Sulawesi]. Fakultas Perikanan dan Ilmu Kelautan USNRAT, Manado, 42 pp. [in Indonesian]
- Mehra S. R., Ghosh R. G., 1965 Bamboo-reinforced soil-cement. *Civil Engineering and Public Works Review*, Vol. 60, no. 711, October 1965; vol. 60, no. 712. November 1965.
- Mundy C. N., Babcock R., 1998 Role of light intensity and spectral quality in coral settlement: implications for depth-dependent settlement? *Journal of Experimental Marine Biology and Ecology* 223(2):235-255.
- Muqsit A., Purnama D., Ta'alidin Z., 2016 [The community structure of coral reef in Pulau Dua, Enggano district, North Bengkulu regency]. *Jurnal Enggano* 1(1):75-87. [in Indonesian]
- Nasaru J. H., Lalamentik L. T. X., Rembet U. N. W. J., 2017 [Distribution of *Pocillopora verrucosa* (Ellis and Solander, 1786) at the reef flat of the south coast of Putus-Putus Island East Ratatotok, Ratatotok District, Southeast Minahasa regency]. *Jurnal Ilmiah Platax* 5(1):61-68. [in Indonesian]
- Ng C. S. L., Chou L. M., 2017 Coral reef restoration in Singapore - past, present and future. *Sustainability Matters: Environmental Management in the Anthropocene* 3, 23 pp.
- Orwa P. O., Omondi R., Chemoiwa E. J., 2018 Colonization patterns of benthic macroinvertebrates in fertilized and non-fertilized earthen fish ponds. *International Journal of Aquaculture and Fishery Sciences* 4(3):022-026.
- Razak T., 2008 The population of hard coral colonies growing on Ecoreef artificial modules on Manado Tua Island, Bunaken National Park, North Sulawesi, Indonesia. Oral presentation material, 11th International Coral Reef Symposium Florida, Session number XXIV, pp. 223.
- Reid C., Marshall J., Logan D., Kleine D., 2009 Coral reef and climate change: the guide for education and awareness. CoralWatch, University of Queensland, Australia, 256 pp.
- Reskiwati, Lalamentik L. T. X., Rembet U. N. J., 2018 [Study on the taxonomy of genus *Favia* (Oken, 1815) at the reef flats of Kampung Ambong Village in Likupang Timur District, North Minahasa]. *Jurnal Ilmiah Platax* 6(1):188-193. [in Indonesian]
- Rondo M., Tamanampo J. F. W. S., 2014 Principles of aquatic ecology. Fakultas Perikanan dan Ilmu Kelautan, Universitas Sam Ratulangi, Manado, 427 pp. [in Indonesian]
- Sheppard C., Davy S., Pilling G., Graham N., 2017 The biology of coral reefs. Oxford University Press, 384 pp.
- Sheehy D. J., 1986 New approaches in artificial reef design and applications. In: Artificial reefs: marine and freshwater applications. Trie F. M. D. (ed), Lewis Publishers Inc., Michigan, USA, pp. 253-263.
- Suharsono, 2008 [Coral species in Indonesia]. Lembaga Ilmu Pengetahuan (LIPI): COREMAP Program, Jakarta, 116 pp. [in Indonesian]
- Suleman Y., Lalamentik L. T. X., Rembet U. N. W. J., 2017 [The Distribution of *Favites abdita* coral reef (Ellis and Solander, 1786) in the land of coral coast village of Malalayan Dua, Malalayang sub-district, Manado city]. *Jurnal Ilmiah Platax* 5(1):69-76. [in Indonesian]
- Varma M. B., 2017 Properties of cement concrete reinforced with bamboo-strip-mat. *IOSR, Journal of Mechanical and Civil Engineering (IOSR-JMCE)* 14(1):47-59.
- Veron J. E. N., 1986 Corals of Australia and the Indo-Pacific. Angus and Robertson Publishers, Australia, 644 pp.
- Veron J. E. N., Stafford-Smith M., 2000 Corals of the world. Volume 1. Australian Institute of Marine Science, Australia, 463 pp.

Veron J. E. N., Pichon M., Wijsman-Best M., 1977 Scleractinia of Eastern Australia - Part 2. Families Faviidae, Trachyphylliidae. Australian Institute of Marine Science Monograph, Series 3: 233 pp.

Received: 18 February 2020. Accepted: 17 March 2020. Published online: 16 April 2020.

Authors:

Laurentius T. X. Lalamentik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: ottylalamentik@yahoo.co.id

R. Ch. Kepel, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: renecharleskepel65@gmail.com

Lawrence J. L. Lumingas, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: ljllumingas@yahoo.com

Unstain N. W. J. Rembet, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: onlyrembet@unsrat.ac.id

Silvester B. Pratasik, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: spjong07@yahoo.com

Desy M. H. Mantiri, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Jl. Kampus Bahu, Manado-95115, North Sulawesi, Indonesia, e-mail: dmh_mantiri@unsrat.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Lalamentik L. T. X., Kepel R. C., Lumingas L. J. L., Rembet U. N. W. J., Pratasik S. B., Mantiri D. M. H., 2020 Faviidae coral colonization living and growing on agricultural waste-materialized artificial substrate. AACL Bioflux 13(2): 910-918.