

Submissions

Biodiversitas Journal of Biological Diversity

Tasks 2

English View Site titlje

OJS
OPEN JOURNAL SYSTEMS

Submissions

Submissions

My Queue 1 Archives 1

Help

My Assigned

Search New Submission

12522 Ransaleleh
Growth of Pteropus alecto Bats (Chiroptera: Pteropodidae) in cages
Revisions have been requested. 1/1

Platform & workflow by OJS / PKP

Type here to search

31°C Berawan 12:00 17/10/2022

Tiltje Ransaleleh, Growth of Pter...

12522 / Ransaleleh / Growth of Pteropus alecto Bats (Chiroptera: Pteropodidae) in cages

Upload File Library

Workflow Publication

Submission Review Copyediting Production

Round 1

Round 1 Status
Revisions have been requested.

Notifications

[biodiv] Editor Decision 2022-10-16 06:14 PM

Type here to search

31°C Berawan 12:06 17/10/2022

Tiltje Ransaleleh, Growth of Pter...

Notifications

[biodiv] Editor Decision 2022-10-16 06:14 PM

Reviewer's Attachments

Search

1063931-1, 12522-Article Text-1063485-1-4-20221010.doc October 16, 2022

Revisions

Search Upload File

No Files

Review Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
No Items				

Type here to search

31°C Berawan 12:06 17/10/2022

Notifications

[biodiv] Editor Decision

2022-10-16 06:14 PM

Tiltje Ransaleleh:

We have reached a decision regarding your submission to Biodiversitas Journal of Biological Diversity, "Growth of Pteropus alecto Bats (Chiroptera: Pteropodidae) in cages".

Our decision is: Revisions Required

[Biodiversitas Journal of Biological Diversity](#)

12619 / Ransaleleh / Growth of Pteropus alecto Bats (Chiroptera: Pteropodidae) in cages

Workflow Publication

Submission Review Copyediting Production

Submission Files

File Name	Date	Type
1064168-1 tiltje, Ransaleleh et al. 2022. Revision accord. reviewer' 1 notes.doc	October 19, 2022	Article Text

[Download All Files](#)

Pre-Review Discussions

Name	From	Last Reply	Replies	Closed
Comments for the Editor	tiltje	-	0	<input type="checkbox"/>
	2022-10-19 03:06 AM			

Growth of *Pteropus alecto* bats (Chiroptera: Pteropodidae) in cages

Abstract. The **Black Flying Fox** (*Pteropus alecto*), known as *Paniki yaki*, is widely consumed by the residents of Minahasa, North Sulawesi, Indonesia. It is imported from outside the provinces of Gorontalo, Central Sulawesi, and South Sulawesi. The peak of selling bats in Sulawesi is during Thanksgiving Day and religious holidays such as Easter, Christmas, and New Year. Even though the conservation status of this species, according to the *International Union for Conservation of Nature* (IUCN), is of **Least Concern**, this species will one day become extinct due to continuous exploration without control. Conservation or cultivation is one of the efforts to preserve this type of bat. Cultivation will be successful when the growth of this mammal is outside the habitat. Therefore, research has been conducted on breeding this animal in cages using five bats. The fruits given as food are papaya, banana, and mango, and the variables measured were the amount of consumption, body weight gain, and morphometry. Furthermore, the data obtained were tabulated and narrated descriptively. The results showed that consumption of *P. alecto* 1,2,3,4, and 5 fruits were 110.64, 147.86, 192.61, 249.18, and 331.61 g/day, with body weight gain of 0.50, 0.53, 0.96, 1, 17, 1, and 50 g/day. The morphometric growth of *P. alecto* 1 was total body length 0.10, forearm 0.06, tibia 0.03, ear 0.01 and wingspan 0.08 mm/day. Sequentially, the morphometric growth of *P. alecto* 2 was 0.11,0.07,0.05, 0.01 and 0.12 mm/day, while *P. alecto* 3 was 0.29, 0.13, 0.06, 0.03, and 0.89 mm/day. The growth of *P. alecto* 4 was 0.19, 0.02, 0.02,0.01 and 0.89 mm/day, while *P. alecto* 5 was 0.20, 0.01,0.01,0.01 and 0.85 mm/day. In conclusion, the 1st and 2nd *P. alecto* were in early bone growth, while the 3rd *P. alecto* was in bone and meat growth, meanwhile the 4th and 5th *P. alecto* were in meat and slow bone growth, which could be seen from morphometric measurement, body weight gain, and total consumption.

Key words: cage, consumption, morphometry, growth, *P. alecto*.

Running title: Growth of bats in cages

INTRODUCTION

Bats are mammals classified in the kingdom Animalia, subphylum Vertebrata, class Mammalia, and order Chiroptera. The order of Chiroptera is divided into two suborders, namely the *Megachiroptera* and the *Microchiroptera* (...). The suborder *Megachiroptera* has one family, Pteropodidae, containing *P. alecto* (Nowak 1994). Bats have a vital function in regulating the balance of ecosystems as agents of fruit pollination (Stewart and Dudash 2016; Thavry et al. 2017; Aziz et al. 2017; Lim et al. 2018; Rodriguez et al.2019; Sheherazade et al. 2019; Ng et al. 2020; Baqi et al. 2021) and seed dispersal, which plays a vital role in forest regeneration and maintenance (Sarmiento et al. 2014; Oleksy et al. 2017; Awind and Jayakumar 2019; Shah et al. 2021). However, bat meat is also used as a source of food by some people (Mildenstein et al. 2016; Suwannarong et al.2016; Ransaleleh et al. 2020)

Wild animal meat may be eaten as halal food by the Minahasa people of North Sulawesi, Indonesia, particularly Christians. Types of wild animal meat consumed include bats (Ransaleleh et al. 2013; Ransaleleh et al. 2020), rats (Laatung et al. 2021), pigs forest (*Sus scrofa*) and Reticulated python (*Malayopython reticulatus* (Latinne et al. 2020)), in Manado used to call rice field snake. The types of bats consumed include *Pteropus alecto*, known as *Paniki yaki*. *P. alecto* meat can be found in traditional markets and supermarkets, especially during certain seasons such as thanksgiving, and religious holidays, namely Easter, Christmas, and New Year. Furthermore, *P. alecto* traded in traditional markets, and supermarkets are imported from neighboring provinces, namely Gorontalo, Central Sulawesi, South Sulawesi, and Southeast Sulawesi. Until 2012, bats were imported by suppliers to North Sulawesi to be sold alive and put in cages made of woven bamboo, then transported using open vehicles for a distance of two to three days. Therefore, many bats died and experienced weight loss due to the stress of the journey. From 2012 to 2022, the suppliers brought bats to North Sulawesi for consumption in frozen form and packaged in styrofoam.

The body weight of traded bats varies between 515-679 g (Ransaleleh et al., 2013), and the survey in September 2021 showed that the weight of traded *P. alecto* bats varied from 300-500 g with the same ratio of males and females (unpublished). The number of bats imported at Christmas and New Year is 500,000 kg (Sheherazade and Tsang, 2015) between 1,000,000 to 1,500,000 when 1 kilogram consists of 2 to 3 individuals. Latinne et al. (2020) reported that the number of bats traded in Sulawesi is estimated at more than one million individuals annually. There is no information on the age of bats from variations in body weight that are hunted and traded. Furthermore, hunters do not consider bats' reproductive and growth status during hunting activities.

Even though the conservation status of the *P. alecto* species is of the least concern (Roberts et al. 2017), the population trend will decline, and sustainability will be threatened when the bats continue to be hunted. It is vital to think

Comment [REV1]: Include the scientific name on the first mention

Comment [REV2]: Is of Least Concern

Comment [REV3]: Where were the bats sourced from and how old are they?

Comment [REV4]: Some of the key words are already included in the title. Remove any key words that are in the title and use new terms to increase paper discoverability

Comment [REV5]: Include a citation here

Comment [REV6]: There is no comma required after the first author as per the Biodiversitas citation style. Please check the author guidelines and revise citations accordingly

Comment [REV7]: Check spacing here

Comment [REV8]: Include the scientific names here

Comment [REV9]: Are there any citations to support here?

53 about actions and solutions for preserving bats following the impact of ecological function. One of the actions or solutions
54 to be considered is the legal status and socialization of bats, conservation (Frick et al. 2019), and cultivation (Ransaleleh
55 et al. 2021). For bat cultivation, many factors should be considered, namely behavior, feed, reproduction and growth. The
56 growth of *P. alecto* in nature and captivity has not been scientifically informed in Indonesia. Therefore, research has been
57 carried out on the growth of *P. alecto* at various ages in cages. The benefit is providing information on the harvesting age
58 of *P. alecto* to facilitate the management of maintenance and utilization.

59 MATERIALS AND METHODS

60 Study area

61 This research was conducted in the Polii-Ransaleleh family captivity administratively located in the V neighborhood of
62 Wanea sub-district, Wanea District, Manado City, North Sulawesi, Indonesia, at coordinates 1°27'39" N and 124°50'33" E
63 (Figure 1). The research was conducted for 6 months, from March to August 2022.



64
65 Figure 1. Research location for Environment V, Wanea District, Manado City, North Sulawesi Province, Indonesia. At coordinates
66 1°27'39" N and 124°50'33" E. Source: Ransaleleh et al. 2021
67

68 Procedures

69 This research were used five pups *P. alecto*. They were progeny from adult males and females *P. alecto* that have been kept in cages
70 since 2011, without growth records (body weight gain, size morphometry, and fruit consumption). *P. alecto* was obtained
71 from different ages and gender according to different times of birth. Sex, age, and initial body weight of the *P. alectos*
72 used in this experiment were 1st female 58 days, 205 g *P. alecto*, 2nd male 79 days, 275 g *P. alecto*, 3rd female 133 days,
73 370g *P. alecto*, 4th male 447 days, 415 g *P. alecto* and 5th male 533 days 470 g *P. alecto* respectively. Furthermore, *P.*
74 *alecto* 1 and 2 were still suckling on their mothers, while *P. alecto* 3 suckles occasionally. *P. alecto* 4 and 5 are no longer
75 breastfeeding, and the separation of *P. alecto* 1 and 2 was based on when the bats could pick fruit and feed themselves in
76 the cage. *P. alecto* 5 was individually housed in a cage of wood than wrapped with ram and isolated with ram into five
77 units. Each cage unit measures 75x50x50 cm (Length x Width x Height) and is equipped with a drinking container. The
78 fruits given as food for *P. alecto* were ripe papaya (*Carica papaya*), banana (*Musa paradisiaca*), and mango (*Mangifera*
79 *indica*). The bats consumed papaya daily, while bananas and mango were given occasionally (Ransaleleh et al. 2021;
80 Ransaleleh et al. 2022).

81 The research procedure is that the *P. alecto* was separated from the group cage, body weight was weighed, and
82 morphometry was measured. In addition, *P. alecto* was weighed and measured morphometrically and placed in a cage unit.
83 The research was conducted for one and five months of pre-study and data collection. Fruits as food ingredients are given
84 daily in the afternoon and evening on an *ad libitum* basis. Before the fruit is given, it is cut into pieces, weighed, placed in
85 its container, and put into the cage unit. The rest of the fruit is weighed in the morning (Ransaleleh et al. 2022). Body
86 weight and morphometric measurements were carried out once a week. The observed variables were the consumption of
87 fruits per day, calculated from the number of fruits given minus those not consumed, and body weight gain (g), calculated
88 from the initial and the final body weight. The morphometry (mm) assessment was performed once a week by weighing
89 the body weight of the head and length, measured from the tip of the snout to the base of the tail. The forearm length was

Comment [REV10]: Use the word sex instead as gender is a social (human) construct.

Comment [REV11]: this isn't very clear. Please explain how they were sourced first.

Comment [REV12]: Feeding on milk

Comment [REV13]: Was only one individual kept in each cage at all times? What was the temperature and humidity that the animals were exposed to? What cage furnishings were available?

90 calculated from the tip of the elbow to the folded wings. The tibia length and ear were measured from the knee joint to the
 91 ankle and from the auditory meatus to the pinna (Wiantoro et al. 2016). Additionally, the wingspan was measured from the
 92 right to the left **wingtips**.

Comment [REV14]: What6 were the cages made of? Were they mesh? Bamboo? Can you provide a supporting image?

93 **Data analysis**

94 The fruit consumption data obtained were tabulated weekly and averaged daily. Data on weight gain were obtained
 95 from the final minus the initial body weight. Furthermore, the morphometric size was tabulated from the final minus the
 96 initial, and all data were presented in tabular form. The data obtained were narrated descriptively for the growth rate using
 97 a regression model.

98 **RESULTS AND DISCUSSION**

99 **Consumption of *P. alecto* bat fruit in cages during the research**

100 Consumption of papaya, banana, and mango of *P. alecto* bats in cages during the research can be seen in Table 1.
 101 The types of fruit consumed by the *P. alecto* five bats were papaya, followed by bananas and mangoes. *P. alecto* 1-5 bats
 102 widely consume papaya fruit containing 88.32% water. In contrast, bananas and mangoes contain less water at 65.16 and
 103 77.5%.

104 **Table1. Consumption of *P. alecto* bat fruit per individual in a cage during the research (grams).**

Fruit type	<i>P. alecto</i> 1	<i>P. alecto</i> 2	<i>P. alecto</i> 3	<i>P. alecto</i> 4	<i>P. alecto</i> 5
Total consumption of papaya fruit	8015.00	13130.00	15180.00	28710.00	24970.00
Average consumption/week	400.75	656.50	759.00	1435.50	1248.50
Average consumption/day	57.25	93.78	108.43	205.07	178.36
Total consumption of bananas	4460.00	4295.00	8800.00	3135.00	17100.00
Average consumption/week	223.00	214.75	440.00	156.75	855.00
Average consumption/day	31.86	30.68	62.85	22.39	178.34
Total consumption of mango fruit	3015.00	3250.00	2985.00	3040.00	4355.00
Average consumption/week	150.75	162.50	149.25	152.00	217.75
Average consumption/day	21.53	23.21	21.32	21.71	31.11
Total consumption of papaya, banana, and mango	15490.00	20675.00	26965.00	34885.00	46425.00
Average consumption/week	774.50	1033.75	1348.25	1744.25	2321.25
Average consumption/day	110.64	147.68	192.61	249.18	331.61

Comment [REV15]: S this measured in grams? Be specific here

106 Based on observations during the research, the bats took the papaya fruit when the fruit was given in the cage. They
 107 sniffed it for a few seconds, pick up the fruit with their claws and put it in their mouth, then chewed and swallowed the
 108 juice after the septum removed. The bats were picked the fruit up again and again if the fruits still available in the cage
 109 until they full enough. The leftover fruits are mostly bananas and mangoes. Since bats are born and kept in cages, the fruit
 110 often given daily as food is papaya. Therefore, they become accustomed to consuming this fruit, which is easily obtained
 111 and available. Bananas and mango are only combined with papaya fruit or occasionally given.

Comment [REV16]: their claws

Comment [REV17]: wording doesn't make sense here. Can you rephrase?

112 The total consumption of papaya, banana, and mango per individual per day of *P. alecto* 1-5 was 110.64 g, 147.68 g,
 113 192.61 g, 249.61 g, and 331.61 g. The difference in the amount of fruit consumption per individual per day was due to
 114 differences in the initial body weight, where the initials were 205.00 g, 255.00 g, 370.00 g, 415.00 g, and 470 g. There
 115 have been no reports on the amount of consumption of papaya, banana, and mango in *P. alecto* per individual per day in
 116 their habitat and captivity. However, the research on fruit preferences consumed by *P. alecto* has been reported by
 117 Ransaleleh et al. (2022). Weber et al. (2015); Win and Mya (2015) stated that the types of fruits that are food for bats of
 118 the genus *Pteropus* include mangoes, papayas, and bananas.

Comment [REV18]: ItALICISE HERE

Comment [REV19]: Do they eat anything else in the wild?

119 The result is concerned with the habit of fruit consumption and the selection of the type consumed by bats daily.
 120 Additionally, body weight is closely related to the amount of fruit consumed in the cage *P. alecto* with a low initial body
 121 weight consumes less fruit. However, there are no scientific reports on the amount of fruit consumption at different body
 122 weights of bats. The results are basic information for selecting types and quantities of fruit in the conservation and
 123 management of *P. alecto*.

124 **Body weight gain of *P. alecto* bats in cages during the research**

125 The weight gain of *P. alecto* in cages can be seen in Table 2. Table 2 shows that the body weight gain of each bat was
 126 different during the research. This was caused by differences in age and body weight of the *P. alecto* 5.

127
128
129
130
131


132 Table 2. Body weight gain of *P. alecto* in cages during the research

	Weight gain (grams)				
	<i>P. alecto</i> 1	<i>P. alecto</i> 2	<i>P. alecto</i> 3	<i>P. alecto</i> 4	<i>P. alecto</i> 5
Initial body weight	205.00	255.00	370.00	415.00	470.00
Final body weight	275.00	345.00	505.00	580.00	685.00
Total body weight gain	70.00	90.00	135.00	165.00	215.00
Average weight gain/week	3.50	3.75	6.75	8.25	10.75
Average body weight gain/day	0.50	0.53	0.96	1.17	1.50

Comment [REV20]: Bat?

Comment [REV21]: Bat?

133 The age of *P. alecto* 1-5 bats at the beginning of the research were 58 days (1 month 28 days = 1.9 months), 79 days (2
 134 months 19 days = 2.6 months), 133 days (4 months 16 days). = 4.5 months), 447 days (1 year 4 months 11 days = 16.3
 135 months), and 533 days (1 year 5 months 19 days = 17.6 months) with initial body weight of 205 g, 255 g, 370 g, 415 g and
 136 470 g. After the research, the ages of *P. alecto* 1, *P. alecto* 1-5, were 189 days (6 months 18 days = 6.6 months), 219 days
 137 (7 months 8 days = 7.3 months), 273 days (9 months 3 days = 9.1 months), 587 (1 year 7 months 17 days = 19.5 months),
 138 and 673 (1 year 10 months 8 days = 22.3 months) with body weight at 275 g, 345 g, 505 g, 580 g, and 685 g. The growth
 139 rate of *P. alecto* 1-5 bats was closely related to the amount of fruit consumption described in the regression model. The
 140 result showed that the growth rate of *Pteropus alecto* 1-5 bats was closely related to the amount of consumption (Figure 2).
 141



143 Figure 2. Growth model, the relationship between total consumption and growth week 1 to 20 of *P. alecto* 1-5 in cages.

Comment [REV22]: Useful information here but make sure that the x and y axis labels are in english

144 Generally, the amount of consumption is closely related to the growth of bats. This is indicated by the high value of R²
 145 (index of determination) (close to the value of 1), ranging from 0.81 to 0.97. The value of R² also shows that the model
 146 obtained is feasible to use, and 81% - 97% changes in growth can be explained by the amount of consumption. The
 147 resulting model shows the value of increasing growth for every 1 unit of consumption (variable x; gram). In *P. alecto* 1, 2,
 148 3, 4, and 5 with the model $y = 82.71 + 0.20x$, $y = 62.44 + 0.22x$, $y = 180.77 + 0.18x$, $y = 152.45 + 0.20x$, and $y = -546.45$
 149 $+ 0.48x$, the increase was 0.20, 0.22, 0.18, 0.20, and 0.48 units. The *P. alecto* 5 regression model has a negative constant
 150 (a) value and is quite large. Therefore, large amount of consumption ($x > 1000$) can achieve a positive (+) growth value.
 151 The high level of *P. alecto* 5 consumption was due to having the largest body weight of 450 grams. The standard error
 152 ranges from 3.66 to 15.31 from the mean value. The error level in obtaining data is quite small due to a relatively high
 153 accuracy. Based on the growth curve in Figure 2, it can be explained that growth is still ongoing until 22.3 months.
 154 Meanwhile, the age the weight gain begins to decrease is unknown. Further research is needed to determine the sigmoid
 155 curve and the growth of bone, meat, and fat components of bats kept in cages.
 156

157 Judging from the age, *P. alecto* 1 and 2 were still slow compared to the growth of 3 at 4.5 months of age weighing
 158 370 g (initial weight). *P. alecto* 2 and 1 bats attained a body weight of 345 g and 275 g at the age of 7.3 and 6.6 months.
 159 The slow growth was because *P. alecto* 1 was 1.9 months old when separated from its mother, and *P. alecto* 2 was 2.6
 160 months old and in lactation. Therefore, the growth became slow because the nutritional needs were not met initially. Todd
 161 et al. (2018) stated that the early growth of animals was focused on bone. The growth of animals under the availability of
 162 fewer nutrients will be affected. Vardon and Tidemann (1998) reported that the lactation period of *P. alecto* bats is 3-4
 163 months.
 164
 165

Comment [REV23]: The word slow is more appropriate

166 The total body weight gain of *P. alecto* 3, 4, and 5 was 135g or 6.75 g per week or 0.96 g per day, 165 g or 8.25 g per
 167 week or 1.17 g per day, and 215 g or 10.75 g per week or 1.50 grams per day, respectively. This research provides
 168 information and illustrates that at 9.1-22.3 months, *P. alecto* bats kept in cages can achieve the same body weight as the
 169 species taken in the wild. Meanwhile, bats under nine months of age cannot achieve weight loss. The bodies of bats
 170 marketed for consumption are the same as those taken in nature/habitat, especially when the infants are separated from
 171 their mothers under the age of weaning. Ransaleleh et al. (2013) reported that the body weight of *P. alecto*, which is
 172 hunted in the wild and traded for consumption, ranges from 508-679 g. However, the results reported by Ransaleleh et al.
 173 (2013) did not provide information on the bats' age. To obtain maximum growth, young *P. alecto* bats cannot be separated
 174 from their mothers under three months. During this period, young bats still need their mother's milk which is rich in
 175 nutrients and needed in early growth.

Comment [REV24]: Good point here

177 Morphometry of *P. alecto* bats in cages during the research

178 Measurement of morphometric characters is useful in determining the unique characteristics possessed by an animal,
 179 including bats. The characteristics of *P. alecto* bats include head, forearm, tibia, ear, and wingspan length. The
 180 morphometric characteristics in the research can be seen in Table 3.

183 Table 3. The increase in the morphometric size of *P. alecto* in the cage during the research

Morphometric increment (mm)	<i>P. alecto</i> 1	<i>P. alecto</i> 2	<i>P. alecto</i> 3	<i>P. alecto</i> 4	<i>P. alecto</i> 5
Body length (head and body) at the beginning of the research	163.00	179.00	220.00	238.00	248.00
Body length at the end of the research	177.00	195.00	261.00	265.00	276.00
Increase in body length during the research	14.00	16.00	41.00	27.00	28.00
Average increase in body length/week	0.70	0.80	2.05	1.35	1.40
Average increase in body length/day	0.10	0.11	0.29	0.19	0.20
The length of the forearm at the beginning of the research	115.00	135.00	158.00	178.00	183.00
The length of the forearm at the end of the research	124.00	152.00	177.00	182.00	184.00
The length of the forearm during the research	9.00	11.00	19.00	4.00	1.00
The average increase in the length of the forearm/week	0.45	0.55	1.05	0.20	0.05
The average increase in the forearm/day	0.06	0.07	0.13	0.02	0.01
Ear length at the beginning of the research	26.00	26.00	27.00	32.00	33.00
Ear length at the end of the research	28.00	28.00	31.00	33.00	33.00
Increase in ear length during the research	2.00	2.00	4.00	1.00	1.00
Average increase in ear length/week	0.10	0.10	0.20	0.05	0.05
Average increase in ear length/day	0.01	0.01	0.03	0.01	0.01
Tibia length at the beginning of the research	52.00	58.00	60.00	70.00	75.00
Tibia length at the end of the research	56.00	65.00	69.00	74.00	79.00
Tibia length increase during the research	4.00	7.00	9.00	4.00	8.00
Average increase in tibia length /week,	0.20	0.35	0.45	0.20	0.20
Average increase in tibia length increase/day	0.03	0.05	0.06	0.02	0.01
Width of the initial wingspan of the research	790.00	850.00	990.00	1120.00	1130.00
Width of the final wingspan of the research	801.00	867.00	1115.00	1240.00	1250.00
Increase in the width of the wingspan during the research	11.00	17.00	125.00	120.00	120.00
Average increase in wingspan/week	0.55	0.85	6.25	6.00	6.00
Average increase in wingspan/day	0.08	0.12	0.89	0.85	0.85

Comment [REV25]: No need for CaPs here

184 The morphometric characteristics provide information that *P. alecto* 1 bats aged 1.9 months have a total body length
 185 (body + head), forearm, ear, tibia, and wingspan length of 163.00 mm, 115.00 mm, 26.00 mm, 52.00 mm, and 790.00 mm,
 186 respectively. After being reared separately from the mother for 144 days, the 6.6-month-old brood had a total body,
 187 forearm, tibia, ear, and wingspan length of 177.00 mm, 124.00 mm, 56.00 mm, 28.00 mm, and 801.00 mm. Likewise, *P.*
 188 *alecto* 2, separated from its mother at the age of 2.6 months, had a body, forearm, tibia, ears and wingspan length of
 189 179.00 mm, 135.00 mm, 58.00 mm, 26.00 mm, and 850.00 mm. After 7.3 months of age, it has a total body, forearm, tibia,
 190 ears, and wingspan length of 195.00 mm, 152.00 mm, 65.00 mm, 28.00 mm, and 867.00 mm.

192 The morphometric characteristics of *P. alecto* 1 and 2 bats are very much different from *P. alecto* 3, which have an
 193 age difference of 2.6 months and 1.9 months. At the beginning of the research, 4.5 months old *P. alecto* 3, had 220.00 mm
 194 body length, 158.00 mm forearm length, 60.00 mm tibia length, 27.00 mm ear length, and 990.00 mm wingspan. After 9.1
 195 months of age, it has a total body length of 261.00 mm, forearm length of wings of 177.00 mm, tibia length of 69.00 mm,
 196 length of ears of 31.00 mm, and a wingspan of 1115.00 mm. The difference in the morphometric characteristics of *P.*
 197 *alecto* 1, 2, and 3 was used as research material, where *P. alecto* 3 had passed the breastfeeding period. Therefore, the
 198 growth of bone size was faster because the nutritional needs were met. Meanwhile, *P. alecto* 1 and 2 were still in the
 199 suckling period, which still needed nutrients from the mother's milk, hence their growth was slow.

200 Differences in morphometric characteristics were also seen in *P. alecto* 4 and 5 when compared with 3. At the age of
 201 16.4 months, *P. alecto* 4 had a total body length of 238.00 mm, forearm length of wings 178.00 mm, tibia length of 70.00
 202 mm, ear length of 32.00 mm, and wingspan 1120.00 mm. After age 19.5 months, there is a total body length of 265.00

Comment [REV26]: Check wording here

203 mm, forearm length of the wings of 182.00 mm, tibia length of 74.00 mm, ear length of 33.00 mm, and wingspan for the
204 forearm of 1270.00 mm. Similarly, *P. alecto* 5, at 17.6 months, had a total body length of 248.00 mm, forearm length of
205 183 mm, tibia length of 75 mm, ear length of 32 mm, and a wingspan of 1130 mm. At 22.3 months, the bat had a total
206 body length of 276.00, forearm length of 184 mm, tibia length of 79.00 mm, ear length of 33 mm, and wingspan of 1250
207 mm. The age difference between *P. alecto* 4 and 3 was 7.3 months, while that of *P. alecto* 5 and 3 was 8.5. However, the
208 difference in total body length was only 23-33 mm higher than *P. alecto* 3, while the forearm, wingspan, tibia, and ear
209 length is only 1.00 mm lower, with a difference in wingspan at 130.00 mm.

210 The growth of *P. alecto* 1, and 2 morphometric characteristics was still slow, while *P. alecto* 3 grew fast, and started
211 slowly for *P. alecto* 4 and 5. Therefore, the bat separated from its mother at the age of 58-79 days, will experience slow
212 growth. The rapid growth of morphometric characteristics occurs in bats that are no longer suckling at the age of 4.5-9
213 months, but slowed down at 16.4-22.3 months. This is identical to the measurement of the body skeleton, and in the animal
214 theory, post-natal growth begins with bone, followed by meat and fat gain. These results inform that *P. alecto* 1 and 2 bats
215 are in the bone growth stage, while *P. alecto* 3 are in the bone and meat growth stage. *P. alecto* 4 and 5 are in the flesh
216 growth stage, with a slowed bone development.

217 The detailed and complete morphometric characteristics have not been reported scientifically. Some studies only
218 reported one of the morphometric characteristics, such as body length, forearm, ears, and tibia, but were not accompanied
219 by age and body weight information. On the contrary, the information submitted in scientific journals only mentions body
220 weight and does not inform the morphometric characteristics. Flannery (1995) reported that the body, forearm, and calf
221 length was 219-278 mm, 156-185 mm, and 68-75 mm, with unknown age and body weight. Vardon and Tidemann (1998)
222 also stated that the forearm length of 109, 136.6, 140.4, 141.7, 142.3, 143.4, 145.3, 149.5, 149.5, 152.6, 155.3, and 157.7
223 are for 0.7-0.8, 2.8-3.0, 3.5-3.6, 33.5-36, 3.7-3.9, 3.8-4.0, 4.5-4.6, 4.5-5.6, 5.7-5.8, 6.5-6.6, 5.7-5.8, 6.5-6.6, 7.5, and 8.4-
224 8.5 months. Ransaleleh et al. (2013) reported that the body weight of bats from nature/habitat sold for consumption ranged
225 from 508-679g, with a forearm length of wings 154.67-166.11 mm, calf 73.93-77.22 mm, and ear 32-32.53. However, the
226 age of the bats on body weight and morphometry are not known. These results are beneficial to complete information on
227 the morphometry, age, and body weight of *P. alecto* bats. The growth of morphometric variables, such as forearm wing
228 and tibia length, grew rapidly at 4.5-9.1 months. The separation was conducted at the age of four months and adult bats
229 can be used for meat at 16.4-22.3 months. In conclusion, the 1st and 2nd *P. alecto* were in early bone growth, while the 3rd
230 *P. alecto* was in bone and meat growth, meanwhile the 4th and 5th *P. alecto* were in meat and slow bone growth, which
231 could be seen from morphometric measurement, body weight gain, and total consumption.

232 ACKNOWLEDGEMENTS

233 The author is grateful to Sam Ratulangi University's Chancellor through the Institute for Research and Community
234 Service Chair, who has financed this research with the 2022 Unsrat Superior Basic Research (RDUU) research scheme
235 with contract number No. 211/UN12-13/LT/2022.

236 REFERENCES

- 237 Ashwin CP, Jayakumar S. 2019. Diet of indian flying fox *Pteropus giganteus* (Brunnich, 1782) in selected districts of Karala and Tamil Nadu India. *J.*
238 *Sci. Trans Environ Technov.* 13(1): 31-36.
- 239 Aziz SA, Clements GR, McConkey KR et al. 2017. Pollination by the locally endangered island flying fox (*Pteropus hypomelanus*) enhances fruit
240 production of the economically important durian (*Durio zibethinus*). *Ecol Evol.* 7(21):8670-8684. [Doi.org/10.1002/ece3.3213](https://doi.org/10.1002/ece3.3213).
- 241 Baqi A, Lim VC, Yazid H, Khan TAA, Lian CJ, Nelson BR, Seelan JSS, Appalasamy S, Mokhtar SI, Kumaran JV. 2021. A review of durian plant bat
242 pollinator interactions. *J Plant Interact.* 17(1):105-126. [Doi=10.1080/17429145.2021.2015466](https://doi.org/10.1080/17429145.2021.2015466)
- 243 Flannery T. 1995. Mammals of the South-West Pacific & Moluccoan Islands. Sydney, Australian Museum / Reed Book.
- 244 Frick WF, Kingston T, Flanders J. 2019. A review of the threat and challenges to global bat conservation. *Ann N Y Acad Sci* 1469(1):5-25.
245 [Doi:10.1111/nvas.14045](https://doi.org/10.1111/nvas.14045)
- 246 Laatung S, Fuah SM, Masyu'ud B, Sumantri C, Salundik. 2021. Spesies of white-tailed forest rats hunted and traded, their conservation status and habitat
247 characteristics, in North Sulawesi Indonesia. *Biodiversitas* 22(7): 2778-2784. [Doi.org/10.13057/biodiv/d220727](https://doi.org/10.13057/biodiv/d220727)
- 248 Latinne A, Saputro S, Kalengkongan J, Kowel CL, Gaghiwuc L, Ransaleleh TA, Nangoy MJ, Wahyuni I, Kusumaningrum T, Safari D, Feferholtz Y, Li
249 H, Hagan E, Miller M, Francisco L, Daszak P, Olival KJ, Pamungkas J. 2020. Characterizing and quantifying the wildlife trade network in
250 Sulawesi, Indonesia. *Glob Ecol Conserv* 21: 1-18. [Doi:10.1016/j.gecco.2019.e00887](https://doi.org/10.1016/j.gecco.2019.e00887)
- 251 Lim VC, Ramlil R, Bhasu S, Wilson J. 2018. Pollination implications of the diverse diet of tropical nectar feeding bats roosting in anurban cave. *Peerj.*
252 [6:e4572. Doi.org/10.7717/peerj.4572](https://doi.org/10.7717/peerj.4572).
- 253 Mildenstein T, Tanshi I, Racey PA. 2016. Exploitation of bats for bushmeat and medicine. In : C.C. Voigt and T. Kingstone (eds). The antropocene
254 conservation of bats in a change world. PP 325-375. DOI 10.1007/978-3-319-25220-9_12
- 255 Ng WS, Mohd-Azlan J, Wong SY. 2020. Floral biology and pollination strategy of durio (malvaceae) in Sarawak. Malaysian Borneo. *Biodiversitas.*
256 [21\(12\):5579-5594. Doi.org/10.13057/biodiv/d211203](https://doi.org/10.13057/biodiv/d211203)
- 257 Nowak RM. 1995. Bats of The World. The Johns Hopkins University Press, Baltimore London.
- 258 Oleksy R, Giuglioli L, McKetterick TJ, Racey PA, Jones G. 2017. Flying foxes create extensive seed shadows and enhance germination success of
259 pioneerplant species in deforested Madagascan landscapes. [Doi.org/10.1371/journal.pone.0184023](https://doi.org/10.1371/journal.pone.0184023)
- 260 Ransaleleh TA, Maheswari RRA, Sugita P, Manalu W. 2013. Identifikasi kelelawar pemakan buah asal Sulawesi berdasarkan morfometri. *Jurnal*
261 *Veteriner.* 14(4). 485-494 [Indonesia]

262 Ransaleleh TA, Nangoy MJ, Wahyuni I, Lomboan A, Koneri R, Saputro S, Pamungkas J, Latinne A. 2020. Identification of bats on traditional market
 263 in Dumoga district, North Sulawesi. IOP Conf.Ser: Earth Environ Sci 473 : 012067. [Doi:10.1088/1755-1315/473/1/012067](https://doi.org/10.1088/1755-1315/473/1/012067)
 264 Ransaleleh TA, Wahyuni I, Kawatu M, Nangoy MJ, Wiantoro S. 2021. Behavior of the back flying fox, *Pteropus alecto* (Chiroptera: Pteropodidae) in
 265 cages. Biodiversitas 22 (12) : 5636-5644. [Doi.org/10.13057/biodiv/d221262](https://doi.org/10.13057/biodiv/d221262)
 266 Ransaleleh TA, Kristi FA, Kawatu M, Nangoy MJ. 2022. Preferensi pakan kelelawar *Pteropus alecto* dipenangkaran/Ex-situ. Zootec 42(1): 52-58
 267 [Indonesia]
 268 Roberts B, Eby P, Tsang SM, Sheherazade. 2017. *Pteropus alecto*. The IUCN Red List of Threatened Species 2017:
 269 e.T18715A22080057. [Doi.org/10.2305/IUCN.UK.2017-2.RLTS.T18715A22080057.en](https://doi.org/10.2305/IUCN.UK.2017-2.RLTS.T18715A22080057.en). Accessed on 31 August 2022.
 270 Rodriguez PAA, Kromer T, Tschapka M, Franco JGG, Sarti JE, MacSwiney GMC. 2019. Bat pollination in bromeliaceae. Plant Ecol. Diversity.
 271 12(1):1-19. [Doi.org/10.1080/17550874.2019.1566409](https://doi.org/10.1080/17550874.2019.1566409)
 272 Sarmento R, Alves-Costa CP, Ayub A, Mello MAR. 2014. Partitioning of seed dispersal services between birds and bats in a fragment of the Brazilian
 273 Atlantic Forest. Zoologia 31(3) 245–255. Doi:10.1590/S1984-46702014000300006.
 274 Shah MNM, Johan KBM, Roslan A, Basri HZH, Pesiu E, Zahidin MA, Abdullah MT, Zalipah MN. 2021. Seed and pollen dispersal by small pteropidid
 275 bats in low land forests of Terengganu Peninsula Malaysia. Jurnal of wildlife and Parks, 36:75-93.
 276 Sheherazade, Ober HK, Tsang SM. 2019. Contributions of bats to the local economy through durian pollination in Sulawesi, Indonesia. Biotropica.
 277 51(6):913–922. [Doi.org/10.1111/btp.12712](https://doi.org/10.1111/btp.12712)
 278 Sheherazade, Tsang SM 2015. Quantifying the bat bushmeat trade in North Sulawesi, Indonesia, with suggestions for conservation action Glob Ecol
 279 Conserv 3 324-330. [Doi:10.1016/j.gecco.2015.01.003](https://doi.org/10.1016/j.gecco.2015.01.003).
 280 Stewart AB, Dudash MR. 2016. Flower-visiting bat species contribute unequally toward agricultural pollination ecosystem services in Southern
 281 Thailand. Biotropica. 49(2):239–248. [Doi.org/10.1111/btp.12401](https://doi.org/10.1111/btp.12401).
 282 Thavry H, Cappelle J, Bumrungsri S, Thona L, Furey N. 2017. The diet of the cave nectar bat (*Eonycteris spelaea* Dobson) suggests it pollinates
 283 economically and ecologically significant plants in Southern Cambodia. Zool Stud. 56(17). [Doi.org/10.6620%2FZS.2017.56-17](https://doi.org/10.6620%2FZS.2017.56-17)
 284 Suwannarong K, Schuler S. 2016. Bat consumption in Thailand. Infect Ecol Epidem 6:29941. [Doi.org/10.3402/iee.v6.29941](https://doi.org/10.3402/iee.v6.29941)
 285 Todd CM, Westcoot D, Rose K, Mathin JM, Welbergen JA. 2018. Slow growth and delayed maturation in a critically endangered insular Flying fox
 286 (*Pteropus natalis*). J. mammal 99(6):151-1521. [Doi.org/10.1093/jmammal/gvy110](https://doi.org/10.1093/jmammal/gvy110)
 287 Vardon M, Tidemann CR. 1998. Reproduction and maturity in the black flying fox *P. alecto* (Megachiroptera : Pteropodidae). Australian Journal of
 288 Zoology 46 : 329-344
 289 Weber N, Duengkae P, Fahr J et al. (2015) High-resolution GPS tracking of Lyle's flying fox between temples and orchards in Central Thailand. J
 290 Wildl Manage 79:957–968. [Doi.org/10.1002/jwmg.904](https://doi.org/10.1002/jwmg.904)
 291 Wiantoro S, Hitch AT, Engilis IE, Gunawan H, Engilis A. 2016. Bats (Chiroptera) recorded in the lowland of Southeast Sulawesi, Indonesia
 292 with notes on taxonomic status and significant range extensions. De Gruyter Mammalia 1-15. [Doi.org/10.1515/mammalia-2015-0153](https://doi.org/10.1515/mammalia-2015-0153)
 293 Win SS, Mya KM. 2015. The diet of the Indian Flying Fox *Pteropus giganteus* (Brünnich, 1782) (Chiroptera: Pteropodidae) in Myanmar—
 294 conflicts with local people? J Threat Taxa 7:7568–7572. [Doi.org/10.11609/JoTT.o4178.7568-72](https://doi.org/10.11609/JoTT.o4178.7568-72)

Comment [REV27]: Include the doi here