

EFFECT OF INCLUSION OF COCONUT MEAL IN THE DIETS ON NUTRIENT DIGESTIBILITY OF GROWING PIGS

by Florencia Sompie 2

Submission date: 22-Jun-2023 09:15AM (UTC+0700)

Submission ID: 2120628627

File name: Marie_dkk_2_EFFECT_OF_INCLUSION_OF_COCONUT_MEAL_IN_THE_DIETS.pdf (325.83K)

Word count: 2680

Character count: 13362

8
**EFFECT OF INCLUSION OF COCONUT MEAL IN THE DIETS
ON NUTRIENT DIGESTIBILITY OF GROWING PIGS**

8
Marie NAJOAN, Betty BAGAU, Fenny WOLAYAN, Florencia SOMPIE, Youdhi KOWEL

19
Faculty of Animal Husbandry, Sam Ratulangi University, North Sulawesi, Indonesia
Jl. Kampus Timur Kleak Manado. Sulawesi Utara. 95115

Corresponding author email: najoanmarie@yahoo.com

Abstract

Although its protein content is less than that of conventional ingredients commonly used as protein sources, copra meal represents the largest quantity of locally available feed protein in many tropical countries, such as Indonesia. It is, therefore, important that information about the nutritional value of copra meal is available. The present study was conducted to test the effect of inclusion of copra meal in growing pigs diets on nutrient digestibility. Twenty-four castrated Landrace Yorkshire pigs (3.5 – 4.0 months of age, ranged from 33 – 40 kg of body weight) were kept in individuals pens. Four treatments were arranged as follow: R1 = 95% basal diet + 5% copra meal; R2 = 90% basal diet + 10% copra meal; R3 = 85% basal diet + 15% copra meal; R4 = 80% basal diet + 20% copra meal, with six replications as blocks based on body weight. The present study was arranged in a 4 x 5 Completely Randomized Block Design with four treatments and five blocks. After a significant F test, Tukey's test (where necessary) was used to inspect differences among treatment means. Differences between treatment means were considered significant when $P < 0.05$. Variables measured were: dry feed and water consumption, nutrient digestibility (energy, crude protein, fat, crude fiber, calcium and phosphorus). The result showed that there was no significant effect ($P > 0.05$) of copra meal inclusion in diet on feed and water consumption, as well as nutrient digestibility (energy, crude protein, fat, crude fiber, calcium and phosphorus). It can be concluded that inclusion of copra meal up to 20% in growing pigs diets did not affect nutrient digestibility as well as performance of the pigs. Further study on higher level of copra meal inclusion in pig diets is needed.

Key words: copra, pigs, consumption, digestibility

INTRODUCTION

The demand for feed is increasing and co-products from the tropical food industries are increasingly used in diets fed to pigs. These co-products include copra meal and copra expellers (Stein et al., 2015). The coconut palm (*Cocos nucifera*) is widely distributed throughout Indonesia, especially in the Province of North Sulawesi. The production of coconut oil and copra meal in Indonesia is about 995.00 metric tons and copra meal are about 515.000 metric tons (BPS - Statistics Indonesia, 2017).

Coconut meal or copra meal is produced from the ripe fruit (nut) of the coconut palm (*Cocos nucifera*). The nut is split and the kernel is removed and dried below 6% moisture. This meal is called copra meal and still contains the oil. Coconut meal (copra meal) is a coproduct of the production of coconut oil. Copra meal is produced by expeller sometimes referred to as coconut meal or coconut oil meal. Copra meal is often fed to monogastric animals such as poultry and pigs (Sundu et al., 2006; Diarra et al., 2004;

Jaworsky et al., 2014; Rodjan et al., 2017; Park et al., 2012) and it could be deposited in adipose tissues (Rumokoy, 2012). Although the amino acid profile and digestibility in copra meal are less favourable than in soybean meal, it can provide significant protein and energy in swine diets and may be used to reduce feed costs (Stein et al., 2015).

The protein concentration of copra meal is between 14 and 22% and the protein has a low biological value and a very high Arg:Lys ratio (Stein et al., 2015). Somkune et al., (2013) reported that dried coconut meal contains about 26% crude protein and 26% crude fiber. Copra meal is high in fiber and the energy value is relatively low when fed to pigs. Its high fiber content is an important factor that limits its used in pigs diets (Kim et al., 2011) as it reduces digestibility of nutrients and efficiency of energy utilization (Gutierrez et al., 2014; Marin et al., 2017). High fiber co-products from the copra industries are by-products of the production of coconut oil. After entering and circulating in animal's body, this oil could be

part of intramuscular substances composition (Rumokoy and Toar, 2014). Although its protein content is less than that of conventional ingredients commonly used as protein sources, copra meal represents the largest quantity of locally available feed protein in many tropical areas, such as Indonesia. It is, therefore, important that information about the nutritional value of copra meal is available. The present study was conducted to test the effect of inclusion of copra meal in growing pigs diets on nutrient digestibility.

MATERIALS AND METHODS

Animals, diet, and management: twenty castrated Landrace X Yorkshire crossbred grower pigs aged 3.0 to 3.5 months old were assigned to four treatments as follow: T1 = 95% basal diet + 5% copra meal; T2 = 90% basal diet + 10% copra meal; T3 = 85% basal diet + 15% copra meal; and T4 = 80% basal diet + 20% copra meal. The present study was arranged in a 4 x 5 Completely Randomized Block Design with four treatments and five blocks. The initial body weight of animals was 28 to 37 kg. Pigs were divided into five blocks as follow: I = 28

to 29 kg; II = 30 to 31 kg; III = 32 to 33 kg; IV = 34 to 35 kg, and V = 36 to 37 kg. Pigs were placed in individual metabolic crate to facilitate the separate collection of urine and faeces. The diet, based on yellow corn, rice bran, soybean meal, bone meal, salt, and vitamin mix (premix), was formulated (Table 1) to meet or exceed all requirements for growing swine as defined on a percentage basis by the NRC (2012).

Table 1. Composition of basal (control) diet

Ingredients	%
Yellow Corn	50,0
Rice Bran	25,0
Soybean Meal	22,0
Bone Meal	1,0
Salt (NaCl)	1,0
Mineral(Pig Mix)	1,0
Total	100
<i>Nutrient Assayed</i>	
Dry matter (%)	18,31
Protein (%)	15,35
Crude fiber (%)	6,32
Ether extract (%)	5,87
Calcium (Ca) (%)	0,70
Phosphorus (P) (%)	0,68
Zinc (Zn)(mg/kg)	107
Digestible Energy (DE) (kcal/kg)	3436,0

Table 2. Composition of Experimental Diets

	Treatments			
	T1	T2	T3	T4
	%			
Control diet	95	90	85	80
Copra meal	5	10	15	20
Total	100	100	100	100
<i>Nutrient Assayed:</i>				
Dry matter (%)	84,31	84,31	83,22	84,94
Protein (%)	15,35	15,37	15,66	16,00
Crude fiber (%)	6,32	6,56	7,01	7,87
Ether extract (%)	5,87	5,88	5,92	5,97
Calcium (%)	0,70	0,71	0,72	0,72
Phosphorus (%)	0,68	0,69	0,70	0,72
DE (kcal/kg)	3436,20	3437,10	3445,20	3425,30

*) Provided per kg of diet: vitamin A, 4800 IU; vitamin D3, 470 IU; vitamin E, 22 IU; menadione, 2.2 g; riboflavin, 2.7 g; niacin, 22 g; d-pantothenic acid, 1.5 g; thiamine, 550 mg; d-biotin, 150 mg; vitamin B12, 20 mg

The diet was offered to the pigs as a mass. Fresh water was provided in the trough ad libitum until the time of feeding. Feed was provided and after 30 min, the uneaten portion was removed and weighed. The trough was refilled with water until the next feeding period. Feed and water intake were recorded during the total experimental period. The design of the trough in

the metabolism crate essentially eliminated wastage of feed or water. Trays placed below the crates revealed insignificant wastage. Total (24 h) faecal output was collected beginning at 08.00 h on day 70. Total samples were frozen and retained for later assay of dry matter, nitrogen, calcium, and phosphorus. Feed samples were collected during the experimental

period and retained daily for later analysis of dry matter, energy, protein, fat, crude fiber, calcium and phosphorus, according to methods defined by the A.O.A.C (2016). All data were analysed using the GLM Procedure (SAS Institute, 1989). A Completely Randomized Block Design (Steel and Torrie, 1997) was used to examine the treatment differences. After a significant F test, Tukey's test (where necessary) was used to detect differences among treatment means. Differences between treatment means were considered significant when $P < 0.05$.

RESULTS AND DISCUSSIONS

The results of the performance of sows and nutrient digestibility values of the experimental pigs are presented in Table 3. Feed and water intake, daily gain, nutrient digestibility (energy, protein, fat, crude fiber, calcium and phosphorus) of the pigs were all not affected ($P > 0.05$) by treatments. Increasing copra meal inclusion in the growing pig's diets had no significant effect on all parameters measured in the present study.

A report based on unpublished research by Jaworski et al., (2017) of University of Illinois at Urbana-Champaign, USA on research conducted from 2008 to 2017, indicated that growth performance declines with increasing inclusion of copra meal at day 20 of experimental period. Average daily gain, average daily feed intake, and gain:feed ratio all showed a linear decrease ($P = 0.05$) as the inclusion rate of copra meal increased. The inclusion of copra meal must be less than 25% in diets for growing-finishing pigs (Stein et al., 2015).

The differences in nutrient digestibility among experiments may be due to differences in nutrient composition, drying procedures, oil extraction procedures, and the degree and duration of heat processing that is used during extraction. Quality problems may be attributed to the high moisture content of copra during drying and storage (Stein et al., 2015). Copra meal may be included in diets fed to growing and finishing pigs by up to 30% without affecting growth performance (Stein et al., 2015), but negative effects of increasing levels of copra meal in the diet have been reported. Results with copra meal have been

improved if diets either were semi-purified diets or if they were formulated based on digestible amino acid rather than based on crude protein (Jaworski et al., 2017).

Copra meal is incorporated in pig diets as a source of plant protein. On the other hand, copra meal has some limitations due to its high fiber content and its low protein quality that can affect its nutrient digestibility when incorporated into pig diets at high level. As was proposed in the present study that the inclusion of copra meal up to 35% in the growing pig diets could affect digestibility of energy and protein as well as the performance of the pigs. Indeed, the inclusion of copra meal from 5, 10, 15, and 20% in the diets did not give a significant ($P > 0.05$) results in all parameters measured. It seems that as the level of copra meal inclusion increased in the diets has not given a major change Palatability of the diet did not change and so that feed consumption did not give a significant difference ($P > 0.05$). As a consequence, digestible energy (DE) in the present study among treatment did not give a significant difference as well due to the energy spent in fiber digestion was similar. Increment of copra meal among level was not too big so that that fiber content did not change much. This could also be the one that contributed to the insignificant result on energy spent in fiber digestion, although fiber content of copra meal is high enough. NRC (2012) stated that ingredients or diets high in fiber always followed by low energy content. In that matter, the increased in feed consumption will be followed by fiber consumption that increased energy used in digesting nutrients and reduced digestible energy per se of the feed as well as its feed efficiency. Copra meal contains between 10 and 16% crude fiber and approximately 47% total dietary fiber (Jaworsky et al., 2014). Concentrations of β -mannans, galactomannans, arabinoxylogalactan, and cellulose are relatively high and the water binding capacity of copra meal is high. Water binding capacity is an estimate of the amount of water that a fiber can absorb and hold after an external force has been applied to it via centrifugation. High water binding capacity will usually result in reduced feed intake of animals because of swelling in the intestinal tract. The relatively high concentrations of fermentable fiber in copra ingredient may result in increased needs for

4
dietary Threonine because dietary fiber increases the endogenous losses of amino acids, and therefore increases the loss of Threonine. Protein levels of copra meal and copra expellers typically range from 20 to 26% (Jaworsky et al., 2014). The concentration of gross energy in copra meal is greater than in corn, but because of the high concentration of fiber in copra meal and copra expellers, concentrations of digestible energy (DE) and metabolizable energy (ME) are less than in corn (NRC, 2012; Sulabo et al., 2013).

CONCLUSIONS

28
Inclusion of copra meal up to 20% in growing pigs diets did not affect nutrient digestibility as well as performance of the pigs. Further study on higher level of copra meal inclusion in pig diets is needed.

REFERENCES

- A.O.A.C. (2016). *Official Methods of Analysis of AOAC INTERNATIONAL*, 20th Edition.
- Diarra, D.S., Parera, D., Tabuaciri, P., Muhammad, U. (2005). Growth performance and carcass yield of broiler chickens fed commercial finisher and cassava copra meal based diets. *J. Appl. Anim. Res.*, 43, 352-356.
- Gutierrez, N.A., Seroo, N.V.L., Kerr, B.J., Zijlstra, R.T., Patience, J.F. (2004). Relationship among dietary fiber components and the digestibility of energy, dietary fiber, and amino acids and energy content of nine cocos products fed to growing pigs. *J. Anim. Sci.*, 92, 4505-4517.
- Jaworski, N.W., Shoulders, J., González-Vega, J.C., Stein, H.H. (2014). Effects of using copra meal, palm kernel expellers, or palm kernel meal in diets for weanling pigs. *Proc. Anim. Sci.*, 30, 243-51.
- Marin, M., Hodoşan, C., Dinişă, G., Nicolae, C. (2017). The influence of the chemical composition of maize, barley and peas hybrids on the digestibility of compound feed for pigs. *AgroLife Scientific Journal*, 6 (2), 127-132.
- National Research Council (2012). *Nutrient Requirements of Swine*: Eleventh Revised Edition. Washington, DC: The National Academies Press. <https://doi.org/10.17226/13298>.
- Park, C.S., Son, A.R., Kim, B.E. (2012). Prediction of gross energy and digestible energy in copra meal, palm kernel meal, and cassava root fed in pigs. *J. Anim. Sci.*, 90, 221-223.
- Rodjan, P., Buesery, K., Choopun, J., Boonchene, P. (2017). The effect of increased levels of dried coconut meal supplemented with an enzyme cocktail on diet utilization in growing pigs. *Songklanakarin J. Sci. Technol.*, 39(1), 101-108.
- Rumokoy, L.J.M. (2012). Precursor adipocyte development as media of lipid metabolism. *Jurnal Lassalian*, 9 (12), 25-31.
- Rumokoy, L.J.M., Toar, W.L. (2014). The content of lipids in intramuscular adipose as a quality determinant of cattle meat product. *E-journal Unisri*, 144-146.
- SAS Institute, Inc. (1989). *Users Guide*, SAS Institute Inc, Cary, NC.
- Somkune, N., Lanvy, C., Nontithai, S., Pongcharee, P., Ran Parklang, N. (2013). Used of solid coconut meal supplemented with exogenous enzyme on productive performance of broiler. *Khon Kaen Agric. J.*, 41, 430-433.
- Steel, R. G. D., Torrie, J. H. (1997). *Principles and Procedures of Statistics. A Biometrical Approach*. 3rd Ed. New York, USA: McGraw-Hill Book Co. Publishing House.
- Stein, H.H., Casas, G.A., Abellilla, J.J., Liu, Y., Sulabo, R.C. (2015). Nutritional value of high fiber co-products from the copra, palm kernel, and rice industries in diets fed to pigs. *J. Anim. Sci. and Biotech.*, 6, 56.
- Sulabo, R.C., Ju, W.S., Stein, H.H. (2013). Amino acid digestibility and concentration of digestible and metabolizable energy in copra meal, palm kernel expellers, and palm kernel meal fed to growing pigs. *J. Anim. Sci.*, 91, 1391-1409.
- Sundu, B., Kumar, A., Dingle, J. (2006). Response of broiler chicks fed increasing level of copra meal and enzymes. *Int. J. Poult. Sci.*, 5(1), 13-18.

EFFECT OF INCLUSION OF COCONUT MEAL IN THE DIETS ON NUTRIENT DIGESTIBILITY OF GROWING PIGS

ORIGINALITY REPORT

20%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	www.luwuk.today Internet Source	2%
2	Submitted to Universitas Brawijaya Student Paper	1%
3	pdffox.com Internet Source	1%
4	Andrea P Mallea, Maryane S F Oliveira, Diego A Lopez, Hans H Stein. "Nutritional value of a new source of cheese co-product fed to weanling pigs", Journal of Animal Science, 2023 Publication	1%
5	Submitted to University of the Philippines Los Banos Student Paper	1%
6	Li, Zhang, Liu, Yang, He, Cao, Yang, Zhong, Lin, Zhuo, Fang, Che, Feng, Xu, Li, Zhao, Jiang, Wu. "Effects of the Ratio of Insoluble Fiber to Soluble Fiber in Gestation Diets on Sow Performance and Offspring Intestinal Development", Animals, 2019 Publication	1%

7	www.mdpi.com Internet Source	1 %
8	2020.agricultureforlife.usamv.ro Internet Source	1 %
9	Sinta Maharani, Rusli Fidriyanto. "Substitution effects of chili leaves powder for copra meal in ruminant feed on In vitro rumen fermentation characteristics", AIP Publishing, 2023 Publication	1 %
10	J.Y. Sung, B.G. Kim. "Prediction models for apparent and standardized total tract digestible phosphorus in swine diets", Animal Feed Science and Technology, 2019 Publication	1 %
11	Submitted to Padjadjaran University Student Paper	1 %
12	Submitted to University of Malaya Student Paper	1 %
13	epubs.icar.org.in Internet Source	1 %
14	s-space.snu.ac.kr Internet Source	1 %
15	biolres.biomedcentral.com Internet Source	1 %
16	lib.dr.iastate.edu Internet Source	1 %

17

Roberto C. Guarte, Werner Mühlbauer, Manfred Kellert. "Drying characteristics of copra and quality of copra and coconut oil", *Postharvest Biology and Technology*, 1996

Publication

<1 %

18

updatepublishing.com

Internet Source

<1 %

19

www.cambridge.org

Internet Source

<1 %

20

Jong-Hoon Kim, Bon-Hwan Ku, Gwang-Pyo Ko, Man-Jong Kang, Kwang-Hee Son, Mi-Ae Bang, Ho-Yong Park. "Enzyme Feed Additive with Arazyme Improve Growth Performance, Meat Quality, and Gut Microbiome of Pigs", *Animals*, 2023

Publication

<1 %

21

Qingyun Li, Xiangshu Piao, Jundi Liu, Zhikai Zeng, Sai Zhang, Xinjian Lei. "Determination and prediction of the energy content and amino acid digestibility of peanut meals fed to growing pigs", *Archives of Animal Nutrition*, 2014

Publication

<1 %

22

U Hatta, O Sjofjan, N Rugaya, B Sundu. "Evaluation of crude cellulase from – fermented copra meal and its effect on feed digestibility and digestive organs development of broiler chickens ", *IOP*

<1 %

Conference Series: Earth and Environmental Science, 2020

Publication

23 doaj.org Internet Source <1 %

24 docsdrive.com Internet Source <1 %

25 www.european-poultry-science.com Internet Source <1 %

26 www.revistas.usp.br Internet Source <1 %

27 Jae-Cheol Jang, Dong Hyuk Kim, Jin Su Hong, Young Dal Jang, Yoo Yong Kim. "Effects of Copra Meal Inclusion Level in Growing-Finishing Pig Diets Containing β -Mannanase on Growth Performance, Apparent Total Tract Digestibility, Blood Urea Nitrogen Concentrations and Pork Quality", *Animals*, 2020
Publication <1 %

28 Jae Cheol Jang, Dong Hyuk Kim, Young Dal Jang, Yoo Yong Kim. "Amino acid digestibility in diets containing copra meal with β -mannanase fed to growing pigs", *Animal Bioscience*, 2021
Publication <1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On