

THE OMEGA-3 AND OMEGA-6 FATTY ACIDS POTENTIAL OF PUMPKIN, CANDLENUT AND NUTMEG SEEDS AS PHYTOADDITIVE FOR POULTRY. A REVIEW

by Florencia Sompie 11

Submission date: 22-Jun-2023 11:44AM (UTC+0700)

Submission ID: 2120704392

File name: ENUT_AND_NUTMEG_SEEDS_AS_PHYTOADDITIVE_FOR_POULTRY._A_REVIEW.pdf (477.63K)

Word count: 5329

Character count: 30769

61
THE OMEGA-3 AND OMEGA-6 FATTY ACIDS POTENTIAL OF PUMPKIN, CANDLENUT AND NUTMEG SEEDS AS PHYTOADDITIVE FOR POULTRY. A REVIEW

Jet Saartje MANDEY, Meity SOMPIE, Florencia Nery SOMPIE

34
Animal Husbandry Faculty, Sam Ratulangi University
Manado, North Sulawesi of Indonesia

46
Corresponding author email: jetsm_fapet@yahoo.co.id

Abstract

Fatty acids are important constituents of plants and known to possess antimicrobial activities. The biological activity and the possibility of the therapeutic of fatty acids of plant extracts as antimicrobial agents is reviewed. Pumpkin seed, candlenut, and nutmeg are rich in omega-3 and omega-6 fatty acids that possess antimicrobial activities, have potential antioxidant, antiinflammatory, antihyperlipidemic properties in poultry feeding. Pumpkin seed, candlenut, and nutmeg are a rich source of fixed and essential oil, triterpenes, and various types of phenolic compounds. The literature about the benefits of plants seed potency as an alternative phytoadditive for poultry was rare. This article provides an overview that on the potency and biological activity of the omega-3 and omega-6 from pumpkin seed, candlenut, and nutmeg as a basis for exploring it as a phytoadditive for poultry. The method used is the synthesis matrix. From the review of the article, it was concluded that pumpkin seed, candlenut, and nutmeg have the potential to be developed as an alternative feed for poultry, and have bioactive constituents that promote health.

49
Key words: antibacterial, omega-3 fatty acid, omega-6 fatty acid, phytoadditive, poultry.

INTRODUCTION

The common fatty acids in plants are saturated or simple unsaturated compounds of C16 or C18 chain length. Palmitic acid (C16) is the major saturated acid in leaf lipids and some seed oils, while stearic acid (C18) is a major saturated acid in seed fats in several plant families. Unsaturated acids based on C16 and C18 are widespread in leaf and seed oils. The tri-unsaturated linolenic acid, linoleic and oleic acids are common (Harborne and Baxter, 1993), and according to Harwood (1980), quantitatively, the major fatty acids in plant are palmitic, linoleic and, in particular, α -linolenic acids.

Many fatty acids are known have antibacterial and antifungal properties (Russel, 1991). The antibacterial properties of antimicrobial lipids have been known since long, when it was shown that fatty acids, a class of antimicrobial lipid, inhibited growth of the *Bacillus anthracis* pathogen that causes anthrax (Thormar, 2010). Antimicrobial lipids such as fatty acids and monoglycerides are promising antibacterial agents that destabilize bacterial cell

membranes, causing direct and indirect inhibitory effects (Yoon et al., 2018).

Fatty acids are released from lipids by the action of enzymes to become free fatty acids, which have potent biological activities (Desbois & Smith, 2010). Fatty acids (FAs) are potential therapeutic antimicrobial agents due to their potency, broad spectrum of activity and lack of classical resistance mechanism against the actions of these compounds (Desbois, 2012). Various long chain polyunsaturated FAs, which are found naturally at high levels in many marine organisms, have been shown to exert highly potent activity against Gram-positive bacteria, including eicosapentanoic acid (EPA; C20:5n-3) (Desbois, 2013) docosahexanoic acid (DHA; C22:6n-1) (Huang & Ebersole, 2010), γ -linoleic acid (GLA; C18:3n-6) and dihomo- γ -linolenic acid (DGLA; C20:3n-6). Similar to many other PUFAs, eicosapentaenoic acid (EPA; C20:5 n-3) exerts potent effects against Gram-positive species, including human pathogens *Bacillus cereus* and *S. aureus*.

Long-chain unsaturated fatty acids, such as α -linolenic acid, show antibacterial activity and are the key ingredients of antimicrobial food

additives and some antibacterial herbs (Zheng et al., 2005).

Long-chain unsaturated fatty acids are bactericidal to important pathogenic microorganisms, including Methicillin-resistant *Staphylococcus aureus* (Trington et al., 1992), *Helicobacter pylori* (Sun et al., 2003), and *Mycobacteria* (Seidel & Taylor, 2004). These antibacterial actions of fatty acids are usually attributed to long-chain unsaturated fatty acids including oleic acid, linoleic acid, and linolenic acid, while long-chain saturated fatty acids, including palmitic acid and stearic acid, are less active (Sun et al., 2003; Seidel & Taylor, 2004).

Pumpkin, scientific name *Cucurbita moschata* is part of the Cucurbitales order, Cucurbitaceae family and *Cucurbita* genus that has long been applied in Asia for medicinal goals (Call et al., 2006). Pumpkin seeds has many nutrients including polysaccharides, essential fatty acid, carotenoids, mineral, active proteins, and essential amino acids. The seeds have a high nutritional value (Fokou et al., 2004). That seeds had medicinal properties for their biological effects such as antimicrobial activities (Abd El-Aziz & Abd El-Kalek, 2011).

Candlenut, scientific name *Aleurites glauca*, belongs to euphorbiaceae family and grows widely in tropical and sub tropical regions. It is also known as Buah Keras in Malaysia, Kemiri in Indonesia, Indian Walnut in India, and Kukui in Hawaii. It is used in folk medicine to treat stomach in children, bad breath, skin sores, fever, headaches, tumors, diarrhoea, asthma and helps in rejuvenating the body after poisoning.

Candlenut seeds was classified as a type of stone fruit because they have physical characteristics of hard skin and shell shape, then the outer surface was roughly curved (Sinaga, 2016). Besides the consumption of core candlenut seeds is very large (Permana et al., 2017). Candlenut is a common spice that contains high levels of fatty acids.

Nutmeg, is dried kernel of broadly ovoid seed of *Myristica fragrans* Houtt (Family: Myristicaceae). It is widely used as spices in culinary preparations in alternative medicine as aphrodisiac (Tajuddin et al., 2003), memory enhancer, antidiarrhea (Grover et al.,

2002). This plant produces two spices: mace and nutmeg. Nutmeg is the seed kernel inside the fruit and mace is the red lacy covering on the kernel. Myristica species are natives of Moluccas, indigenous to India, Indonesia and Sri Lanka and now cultivated in many tropical countries (Pal et al., 2011). About 30-55% of the seed consists of oils and 45-60% consists of solid matter including cellulose materials.

Nutmeg seeds supplementation may improve blood lipids, ameliorate oxidative stress and this may be due to interactive or additive effects of the numerous bioactive constituents (Thomas & Krishnakumari, 2016). The medicinal use of nutmeg and its use as a spice suggest that it contains some constituents which are responsible for the reported biological activities (Al-Jurayy et al., 2012). Nutmeg extract ameliorates hyperglycemia and abnormal lipid metabolism in animal models (Arulmozhi et al., 2007).

This paper provides a comprehensive overview of various aspects of the species of pumpkin seeds, candlenut, and nutmeg in relation to its potential, which will be used as a benchmark for its use as phytoadditive.

MATERIALS AND METHODS

The technique and instrument used to present a comprehensive overview of the potential of pumpkin seeds, candlenut, and nutmeg is a synthesis matrix. The process is to integrate the results of the analysis of articles based on the similarities and differences of each article. Then make conclusions based on the identification and classification of the potential topic of pumpkin seeds, candlenut, and nutmeg (Ramdhani, et al., 2014). The synthesis matrix is a table/diagram that allows researchers to group and classify different arguments from several articles and combine different elements to get an impression / conclusion on the whole article in general (Murniati et al., 2018).

RESULTS AND DISCUSSIONS

The synthesis of research articles on the antimicrobial lipid potential of pumpkin seed, candlenut, and nutmeg shown in Tables 1, 2, and 3.

Table 1. Omega-3 and Omega-6 Potency of Pumpkin Seed

Descriptions/Issues of the Plant Seeds	References (Author, Year)
<ul style="list-style-type: none"> seed oil showed that predominant unsaturated were linoleic (42%) and oleic (38%), while the major saturated were palmitic (12.7%) and stearic (6%) 	Esuoso et al., 1998
<ul style="list-style-type: none"> oil showed the saturated fatty acids content was 27.73% and consisting of 16.41% palmitic acid and 11.14% stearic acid unsaturated fatty acids value was 73.03% and consisting of 18.14% oleic acid and 52.69% linoleic acid 	Alfawas, 2004
<ul style="list-style-type: none"> Up to 60.8% of the pumpkin seed oil is from the fatty acids, oleic (up to 46.9%), linolenic (up to 40.5%) and palmitic and stearic (up to 17.4%) 	Nakiae et al., 2006
<ul style="list-style-type: none"> Excellent source of protein, minerals, vitamins and unsaturated fatty acids 	Juranovic et al., 2003; Siegmund & Murkovic, 2004; Stevenson et al., 2007
<ul style="list-style-type: none"> highly nutritional and rich nutritional components such as unsaturated fatty acids especially palmitic acid, stearic acid, oleic acid and linoleic acid 	Raharjo et al., 2011
<ul style="list-style-type: none"> The predominant fatty acids were stearic, palmitic, oleic acid and linoleic acid 	Karanja et al., 2013
<ul style="list-style-type: none"> showed a high content of unsaturated fatty acids and the dominant fatty acids were palmitic, stearic, oleic, and linoleic acids the seeds were well endowed in crude oil, protein, carbohydrates and crude fibre. the oil contained unsaturated fatty acids and α-tocopherol 	Miura, 2013
<ul style="list-style-type: none"> Those essential fatty acids are belonging to the ω-6 and ω-3 family which exert amazing nutritional functions and play important role in many metabolic pathways 	Kulaitienė et al., 2018
<ul style="list-style-type: none"> Polyunsaturated linoleic fatty acid was the predominant fat component in pumpkin seed oil. In saturated fatty acids, palmitic and stearic acids predominated. The antioxidant activity increased proportionally with the phenolic content 	

Pumpkin seed, candlenut and nutmeg have been used as an alternative feed ingredient in poultry production.

Pumpkin seeds proximate analysis revealed a higher crude protein, moisture and mineral content (Nworgu et al., 2007).

Hajati et al. (2011) indicated that supplementation of diets with 5 g pumpkin seed oil/kg dry matter in corn soybean meal-wheat based diet can be profitable because it reduced broiler chickens mortality and it did not have any adverse effect on the performance of birds.

Table 2. Omega-3 and Omega-6 Potency of Candlenut

Descriptions/Issues of the Plant Seeds	References (Author, Year)
<ul style="list-style-type: none"> This plant possesses the antimicrobial activity against <i>S. typhi</i>, <i>Vibrio cholera</i>, and <i>E. coli</i>, lowering cholesterol and lipid absorption 	Pedrosa et al., 2002
<ul style="list-style-type: none"> anti-inflammatory and antipyretic also as an analgesic 	Niazi et al., 2010 Quintao et al., 2011
<ul style="list-style-type: none"> seed contain high proportions of polyunsaturated fatty acid (PUFA) such as ω-3, ω-6, and ω-9 	Martin et al., 2010; Rohaida et al., 2014
<ul style="list-style-type: none"> safe for internal uses, and its extract also has the potential to treat selected autoimmune inflammatory diseases by inhibiting the growth of the bacterial triggers 	Mpala et al., 2017
<ul style="list-style-type: none"> possess the antimicrobial activity against <i>S. typhi</i>, <i>Vibrio cholera</i>, and <i>E. coli</i>, lowering cholesterol and lipid absorption, anti-inflammatory, antipyretic, and an analgesic 	Pedrosa et al., 2002; Niazi et al., 2010
<ul style="list-style-type: none"> Based on spectroscopic analysis, these isolate of saponins may be predicted as triterpenoid saponins, diosgenin 	Amalia et al., 2020

Table 3. Omega-3 and Omega-6 Potency of Nutmeg

Descriptions/Issues of the Plant Seeds	References (Author, Year)
<ul style="list-style-type: none"> two types of oils extracted from the nutmeg seed, the essential oil and the fixed oil called the nutmeg butter 	Forrest & Heacock, 1972
<ul style="list-style-type: none"> As a plant seed, myristic acid is the main part of fat. Phenylalanine is the dominant amino acid in nutmeg 	Pathak & Ojha, 1957; Maya et al., 2006
<ul style="list-style-type: none"> the fatty acid composition of the triacylglycerol, the major lipid component, were myristic, palmitic, lauric, petroselinic, and stearic acids 	Niyas et al., 2003
<ul style="list-style-type: none"> Scientists reported that nutmeg have hypolipidemic and hypocholesterolemic effects, antimicrobial, antidepressant, aphrodisiac, memory-enhancing, antioxidant, and hepatoprotective properties 	Jaiswal et al., 2009
<ul style="list-style-type: none"> The major constituents fatty oil of Indian nutmeg were oleic acid, arachidic acid, palmitic acid The major constituents fatty oil of Sri Lankan nutmeg were myristic acid and palmitic acid 	Naher et al., 2013
<ul style="list-style-type: none"> The major biological compounds in the methanol extract were 9,12-Octadecadienoic acid methyl ester, cyclododecyne, and octadecanoic acid. the hexane extract constituents were margaric acid, oleic acid, and 9,12-octadecadienol. 	Anaduaka et al., 2020
<ul style="list-style-type: none"> Nutmeg has been reported to have antioxidant, anti-tumor, and antibacterial effects, and more 	Olaleye et al., 2006; Acuña et al., 2016; Le et al., 2017; Zhang et al., 2016; Gupta et al., 2013
<ul style="list-style-type: none"> Nutmeg extract posses antimicrobial antioxidant and anticancer activity. It support the fact that nutmeg extract can be used as a pure drug 	Chakraborty et al., 2015
<ul style="list-style-type: none"> Antioxidant activity of the mace essential oil examined, β-carotene in linoleic acid and percent inhibition in linoleic acid (67.9%) system. 	Din et al., 2021
<ul style="list-style-type: none"> Nutmeg seeds may improve blood lipids, ameliorate oxidative stress and this may be due to interactive or additive effects of the numerous bioactive constituents 	Thomas and Krishnakumari, 2016

Pumpkin seed oil feed trials on broiler birds have also proved to lower bird mortality, as well as reduced cholesterol and triglyceride concentrations in blood plasma (13). Mart'inez et al. (2010a, b) reported that 10% inclusion of pumpkin seed meal in broiler chicken diets served as a suitable substitute for soya bean meal as it enhanced the reduction in excessive abdominal fat, leading to increased production performance and improved organoleptic meat quality (42).

The inclusion of 0, 33, 66 and 100 (26) g of *Cucurbita moschata* in broiler diets, partially replacing soybean meal and vegetable oil, improved live performance and edible portions yield. In addition, abdominal fat and serum levels of harmful lipids were reduced, whereas (22) μm levels of beneficial lipids increased. There was no effect on meat sensory quality (Aguilar et al., 2011).

Tabari et al. (2016) investigated that use of diet supplemented with pumpkin seed oil improved body weight and (55) increased feed consumption in broiler chickens as a result of the positive effect of pumpkin seed oil (13) the intestine conditions leading to better digestion, absorption and utilization of nutrients and also due to the positive role of pumpkin seed oil on keeping a balanced microflora in the digestive tract.

That supplementary pumpkin seeds oils at a level 10 and 15 (g/kg) have a beneficial effect on productive trait, with no significantly effect of carcass characteristics, also pumpkin seeds oils in same levels reduced total plasma cholesterol concentrations and triglycerides in Japanese quail (Abbas et al., 2017). Pumpkin and flaxseed oils supplementation in feed mixtures of laying hens have a positive effect on the egg weight. Significantly higher average egg's weight during experiment was found after dietary oils supplementation. Tendency of the highest egg's weight was found after flaxseed oil supplementation (Herke et al., (6) 14).

Pumpkin seed extract is reported to be useful for immunomodulation, reproductive health, therapeutics over a wide range of disease conditions and stimulates metabolism of accumulated fats. (57) Pumpkin seeds are a valuable source of protein and fat. Their bioactivity offers prospects for natural control of pathogenic/parasitic organisms, stimulate nutri-

tion or enhance resistance to disease infections, and reduce abdominal fat and serum levels of harmful lipids, while increasing serum levels of beneficial lipids (Achilonu et al., 2018).

Broilers fed candlenut powder had significantly lower meat (51) cholesterol content compared to basal diets. The use of candlenut powder as feed additive at the level of 1% is safely recommended to give better blood profile and reduce meat cholesterol content of broilers (Putri et al., 2018). Supplementing either treated or untreated candlenut meal at 2% level was shown to enhance the fatty acid profiles in broiler chickens meat (Rohaida et al., 2014). Supplementing 2.5% of various components of candlenut kernel in the diet did not improve growth performance, carcass yield, (32) the chemical composition of broiler meat, and fatty acid composition of breast and thigh muscles of finishing broiler chickens (Rasid et al., 2019). The potential of the *A. mollucanas* nut as inhibitors of the growth of bacterial species associated with the onset of rheumatoid arthritis, ankylosing spondylitis and rheumatic heart disease (Mpala et al., 2017).

The *Myristica fragrans* seed meal supplementation at 0.25% enhanced the body weight gain, improved serum, and meat glutathione peroxidase and catalase, and reduced the broiler's meat cholesterol level and lipid oxidation (Adu et al., 2020). Supplementing either treated or untreated candlenut meal at 2% level was shown to enhance the fatty acid.

Supplementation 30% aqueous extract, 5% and 10% nutmeg increased weight and high profiles in broiler chickens meat (Rohaida et al., 2014). That 30% aqueous extract, 5% and 10% supplemented nutmeg (*Myristical fragrance*) increased weight and high density lipoprotein (HDL) concentration and decreased blood glucose, low density lipoprotein (LDL) (35), triglyceride and total cholesterol. Nutmeg diet exhibited significant anti-hyperglycemic in alloxan-induced diabetic rats (Oyindamola et al., 2017). The subchronic administration of 50, 100, and 200 mg/kg bw of nutmeg ethanolic extract did not cause the change of hematological parameters in rat (Bachri et al., 2017). *Myristica fragrans* ethanolic seed extract have hypolipidemic effect. *Myristica fragrans* ethanolic seed extract possess cardioprotective effect on experimentally

induced cardio toxic myocardial infarcted rats (Thomas & Krishnakumari, 2016)

The antioxidant and antiinflammatory activity possessed by nutmeg could be helpful in preventing or slowing the progress of various oxidative stress-related diseases and inflammatory diseases (Sethi & Dahiya, 2018). The subchronic administration of 50, 100, and 200 mg/kg bw of nutmeg ethanolic extract did not cause the change of hematological parameters in rat (Bachri et al., 2017). Seed of *M. fragrans* confirmed 29 anti-inflammatory properties and suggested that it may have deleterious effects on haemopoiesis at high doses (Bamidele et al., 2011).

As presented in this review, there is enormous potential for employing antimicrobial lipids to combat bacterial infections for animal health and human health. Over the past few decades, significant progress has been made towards understanding the relative potency and spectrum of antibacterial activity for different classes of antimicrobial lipids, in turn identifying particularly promising phytoadditive candidates through biological investigations.

CONCLUSIONS

47

The general conclusion of this literature study is that pumpkin seeds, candlenut, and nutmeg have bioactive constituents that promote health. By understanding how antimicrobial lipids function and the critical role of molecular self-assembly, need to begin to design new strategies to enhance 58 herapeutic performance. Recognizing the challenges of antibiotic-resistant bacteria and taking advantage of the low cost and abundant supply of antimicrobial lipids, there is excellent opportunity to further explore antimicrobial lipids as next-generation antibacterial agents for animal health and human health. All of these findings bring us to the new idea in developing and innovating nutraceuticals, pharmaceuticals, and products from pumpkin seeds, candlenut, and nutmeg as phytoadditive for poultry.

REFERENCES

Abbas, R.J., AlShaheen, S.A., & Majeed, T.I. (2017). Evaluation of the productive and physiological performance of japanese quail (*Coturnix coturnix japonica*) fed different levels of pumpkin (*Cucurbita*

moschata) seeds oil. *Interational Journal Veterinary Science*, 6 (1), 31-35.

Abd El-Aziz, A.B., & Abd El-Kalek, H.H. (2011). Antimicrobial proteins and oil seeds from pumpkin (*Cucurbita moschata*). *Nature and Science*, 9(3), 105-119.

Achilonu, M.C., Nwafor, I.C., Umesiobi, D.O., & Sedibe, M.M. (2018). Biochemical proximates of pumpkin (*Cucurbitaceae* spp.) and their beneficial effects on the general well-being of poultry species. *Journal of Animal Physiology and Animal Nutrition*, 102, 5-16.

Acuña, U.M., Carcache, P.J.B., Matthew, S. Blanco, E.J. C.D. (2016). New acyclic bis phenylpropanoid and neolignans, from *Myristica fragrans* Houtt., exhibiting PARP-1 and NF-κB inhibitory effects. *Food Chemistry*, 202. 269–275.

Adu, O.A., Gbore, F.A., Oloruntola, O.D., Falowo, A.B., & Olarotimi, O.J. (2020). The effects of *Myristica fragrans* seed meal and *Syzygium aromaticum* leaf meal dietary supplementation on growth performance and oxidative status of broiler chicken. *Bulletin of the National Research Centre*, 44.149.

Aguilar, Y.M., Yero, O.M., Navarro, M.I.V., Hurtado, C.A.B., López, J.A.C., Mejía, L.B.G. (2011). Effect of squash seed meal (*Cucurbita moschata*) on broiler performance, sensory meat quality, and blood lipid profile. *Brazilian Journal of Poultry Science*, 13 (4), 219-226.

Alfawaz, M.A. (2004). Chemical composition and oil characteristics of pumpkin (*Cucurbita maxima*) seed kernels. *Research Bulletin*, 129, 5-18.

Al-Jumaily, E.F., Al-Amiry, M.H.A., & Assad, J.I. (2012). Hepatotoxic activity of essential oil from nutmeg (*Myristica fragrans*) against tetrachloride-induced hepatic damage in mice. *IOSR Journal of Pharmacy and Biological Sciences*, 2 (6). 01-08.

Amalia, M., Alvionita, M., Subandi, Susanti, E., & Nugraheni, D. (2020). Isolation, identification, and inhibition of saponin isolates from pineapple (*Ananas comosus* L.) and candlenut (*Aleurites moluccanus* L.) against xanthine oxidase by in vitro assay. *IOP Conf. Series: Materials Science and Engineering*, 833, 012004.

Anaduaka, E.G., Uchendu, N.O., & Ezeanyika, L.U.S. (2020). Mineral, amino acid and fatty acid evaluations of *Myristica fragrans* seeds extracts. *Scientific African*, 10, e00567. 1-5

Arulmozhi, D.K., Kurian, R., Veeranjanyulu, A., & Bodhankar, S.L. (2007). Antidiabetic and antihyperlipidemic effects of *Myristica fragrans* in animal models. *Pharmaceutical Biology*, 45 (1), 64-68.

Bachri, M.S., Yuliani, S., & Sari, A.K. (2017). Effect of subchronic administration of nutmeg (*Myristica fragrans* Houtt) ethanolic extract to hematological parameters in rat. *IOP Conf. Series: Materials Science and Engineering*, 259, 012009.

Bamidele, O., Akinnuga, A.M., Alagbonsi, I.A., Ojo, O.A., Olorunfemi, J.O., & Akuyoma, M.A. (2011). Effects of ethanolic extract of *Myristica fragrans* Houtt. (nutmeg) on some heamatological indices in

- albino rats. *International Journal of Medicine and Medical Sciences*, 3(6), 215-218.
- Call, F., Huan, S., & Quanhong, L. (2006). A review on pharmacological activities and utilization technologies of pumpkin. *Plant Foods for Human Nutrition*, 61, 73-80.
- Chakraborty, P., Lavanya, P., & Abraham, J. (2015). Bioactivity of *Myristica fragrans* methanol extract. *World Journal of Pharmaceutical Research*, 4 (9), 1145-1157.
- Desbois, A.P., & Smith, V.J. (2010). Antibacterial free fatty acids: activities, mechanisms of action and biotechnological potential. *Applied Microbiology Biotechnology*, 85(6), 1629-1642.
- Desbois, A.P. (2012). Potential applications of antimicrobial fatty acids in medicine, agriculture and other industries. *Recent Patents on Anti-Infective Drug Discovery*, 7(2), 111-22.
- Desbois, A.P., & Lawlor, K.C. (2013). Antibacterial activity of long-chain polyunsaturated fatty acids against *Propionibacterium acnes* and *Staphylococcus aureus*. *Marine Drugs*, 11(11), 4544-4557.
- Din, M.U., Ali, A., Yasir, M., Jilani, M.I., Shoaib, S., Latif, M., Ahmad, A., Naz, S., Aslam, F., Iqbal, M., & Nazir, A. (2021). Chemical composition and in vitro evaluation of cytotoxicity, antioxidant and antimicrobial activities of essential oil extracted from *Myristica Fragrans* Houtt. *Polish Journal Environmental Studies*, 30(2), 1-6.
- Esuoso, K., Lutz, H., Kutubuddin, M., & Bayer, E. (1998). Chemical composition and potential of some underutilized tropical biomass. I: fluted pumpkin (*Telfairia occidentalis*). *Food Chemistry*, 61, 487-492.
- Farrington, M., Brenwald, N., Haines, D., & Walpole, E. (1992). Resistance to desiccation and skin fatty acids in outbreak strains of methicillin-resistant *Staphylococcus aureus*. *Journal Medicine Microbiology*, 36, 56-60.
- Fokou, E., Achu, M., & Tchouanguép, M. (2004). Preliminary nutritional evaluation of five species of egusi seeds in Cameroon. *African Journal Food Agriculture Nutrition Development*, 4(1), 1-11.
- Forrest, J.E., & Heacock, R.A. (1972). Nutmeg and mace, the psychotropic spices from *Myristica fragrans*. *Lloydia*, 35(4), 440-449.
- Glew, R.H., Glew, R.S., Chuang, L.T., Huang, Y.S., & Millson, M. (2006). Amino acid, mineral and fatty acid content of pumpkin seeds (*Cucurbita* spp) and *Cyperus esculentus* nuts in the Republic of Niger. *Plant Foods Human Nutrition*, 61, 51-56.
- Grover, J.K., Khandkar, S., Vats, V., Dhunoo, Y., & Das, D. (2002). Pharmacological studies on *Myristica fragrans*-antidiarrheal, hypnotic, analgesic and hemodynamic (blood pressure) parameters. *Methods Find Experiment Clinical Pharmacology*, 24, 675-680.
- Gupta, A.D., Bansal, V.K., Babu, V., & Maithil, N. (2013). Chemistry, antioxidant and antimicrobial potential of nutmeg (*Myristica fragrans* Houtt.). *Journal of Genetic Engineering & Biotechnology*, 11, 25-31.
- Hajati, H., Hasanabadi, H.A., & Waldroup, P.W. (2011). Effect of dietary supplementation with pumpkin oil (*Cucurbita pepo*) on performance and blood fat of broiler chickens during finisher period. *American Journal Animal Veterinary Science*, 6, 40-44.
- Harborne, J.B., & Baxter, H. (1993). *Phytochemical Dictionary: A Handbook of Bioactive Compounds from Plants*. London, UK: Taylor and Francis Ltd Publishing House, pp 34-38.
- Harwood, J.L. (1980) *Plant acyl lipids: structure, distribution and analysis*. In: Stumpf PK (ed) *The Biochemistry of Plants*. Vol 4. New York, USA: Academic Press Inc. Publishing House, pp 24-30.
- Herkel, R., Gálík, B., Bíro, D., Rolíneck, M., Šimko, M., Juráček, M., Majlát, M., & Arpášová, H. (2014). The effect of pumpkin and flaxseed oils on selected parameters of laying hens performance. *Acta fytotechn. Zootechnology*, 17 (3), 96-99.
- Jaiswal, P., Kumar, P., Singh, V.K., & Singh, D.K. (2009). Biological effects of *Myristica fragrans*. *Annual Review of Biomedical Sciences*, 11, 21-29.
- Juranovic, I., Breinhoelder, P., & Steffan, I. (2003). Determination of trace elements in pumpkin seed oils and pumpkin seeds by ICP-AES. *Journal Analytical Atomic Spectrometry*, 18, 54-58.
- Karanja, J.K., Mugendi, B.J., Khamis, F.M., & Muchugi, A.N. (2013). Nutritional composition of the pumpkin (*Cucurbita* spp.) seed cultivated from selected regions in Kenya. *Journal of Horticulture Letters*, 3 (1), 17-22.
- Kulaitienė, J., Černiauskiene, J., Jariene, E., Danilčenko, H., & Levickienė, D. (2018). Antioxidant activity and other quality parameters of cold pressing pumpkin seed oil. *Notulae Botanicae Horti Agrobotania*, 46 (1), 161-166.
- Martin, C., Moure, A., Martin, G., Carrillo, E., Dominquez, H., & Parajo, J.C. (2010). Fractional characterization of jatropha, neem, moringa, trisperma, castor and candlenut seeds as potential feedstock for biodiesel production in Cuba. *Biomass and Bioenergy*, 34, 553-538.
- Martínez, Y., Valdivi'e, M., Martínez, O., Estarroz, M., & Co'rdova, J. (2010a). Utilization of pumpkin (*Cucurbita moschata*) seed in broiler chicken diets. *Cuban Journal of Agricultural Science*, 44, 387-391.
- Martínez, Y., Valdivi'e, M., Estarroz, M., Solano, G., & Co'rdova, J. (2010b). Serum lipid profile of laying hens fed pumpkin (*Cucurbita maxima*) seed level. *Cuban Journal of Agricultural Science*, 44, 392-393.
- Maya, K.M., Zachariah, T.J., Krishnamurthy, K.S., Rema, J., & Krishnamoorthy, B. (2006). Fatty acids and leaf amino acids in *Myristica fragrans* Houtt. and related taxa. *Indian Journal Horticulture*, 63, 316-318.
- Miura, Y. (2013). The biological significance of ω -oxidation of fatty acids. *Proceeding Japan Academy, Series B Physical & Biological Sciences*, 89 (8), 370-382.
- Mpala, L.N., Chikowe, G.R., & Cock, I.E. (2017). *Aleurites moluccanus* (L.) Willd. extracts inhibit the growth of bacterial triggers of selected autoimmune inflammatory diseases. *Pharmacognosy Communications*, 7, 83-90.

- Mpala, L.N., Chikowe, G.R., & Cock, I.E. (2017). *Aleurites moluccanus* (L.) Willd. extracts inhibit the growth of bacterial triggers of selected autoimmune inflammatory diseases. *Pharmacognosy Communications*, 7 (2), 83-90.
- Murniarti, E., Naiggolan, B., Panjaitan, H., Pandiangan, L.E.A.M., Widyani, I.D.A., Dakhi, S. (2018). Writing matrix and assessing literature review: A methodological elements of a scientific project. *Journal of Asian Development*, 4 (2), 133-146.
- Naher, S., Rahman, M.M., Shahin, R., Hasan, A.S.M., Bhuiyan, M.N.H., & Ahsan, A. (2013). Comparative studies on physicochemical properties and analysis of fatty oil of the two varieties of the *Myristica fragrans* Houtt (nutmeg) seed. *International Journal of Pharmaceutical and Phytopharmacological Research*, 3 (2).
- Nakiaie, S.N., Rade, D., Kevin, D., Štrucelj, D., Mokrovič, Z., & Bartolija, M. (2006). Chemical characteristics of oils from naked and husk seeds of *Cucurbita pepo* L. *European Journal of Lipid Science Technology*, 108, 936-943.
- Niazi, J., Gupta, V., Chakarboroty, P., & Kumar, P. (2010). Anti-inflammatory and antipyretic activity of *Aleurites moluccana* leaves. *Asian Journal of Pharmaceutical and Clinical Research*, 3 (1), 35-37.
- Niyas, Z., Variyar, P.S., Gholap, A.S., & Sharma, A. (2003). Effect of gamma-irradiation on the lipid profile of nutmeg (*Myristica fragrans* Houtt.). *Journal of Agriculture Food Chemistry*, 51(22), 6502-6504.
- Nworgu, F.C. (2007). Economic importance and growth rate of broiler chickens served fluted pumpkin (*Telfaria occidentalis*) leaves extract. *African Journal of Biotechnology*, 6, 167-174.
- Olaleye, M.T., Akinnmoladun, C.A., & Akindahunsi, A.A. (2006). Antioxidant properties of *Myristica fragrans* (Houtt) and its effect on selected organs of albino rats. *African Journal of Biotechnology*, 5, 1274-1278.
- Oyindamola, E.A., Bolanle, A.O., & Abass, O.O. (2017). Effect of nutmeg (*Myristica fragrans*) on oxidative stress in alloxan-induced diabetic in Wistar albino rats. *World Journal of Pharmacy and Pharmaceutical Sciences*, 6 (4), 1901-1908.
- Pal, M., Srivastava, M., Soni, D.K., Kumar, A., & Tewari, S.K. (2011). Composition and anti-microbial activity of essential oil of *Myristica fragrans* from Andaman Nicobar Island. *International Journal of Pharmacology Life Science*, 2 (10), 1115-1117.
- Pathak, S.P., Ojha, V.N. (1957). The component glycerides of nutmeg butter (*Myristica fragrans*). *Journal of Science Food Agriculture*, 8, 537-540.
- Pedrosa, R., Meyre-Silva, C., Cechinel-Filho, V., Benassi, J.C., Oliveira, L.F.S., Zancanaro, V., Dal Magro, J., & Yunes, R.A. (2002). Hypolipidaemic activity of methanol extract of *Aleurites moluccana*. *Phytotherapy Research*, 16, 765-768.
- Pemana, K.D.A., Hartati, A., & Admadi, B. (2017). The effect of the concentration of sodium chloride solution (NaCl) as a immersion of the quality characteristics of Pati Ubi Talas (*Calocasia esculenta* L. Schott). Faculty of Agricultural Technology. Udayana University. *The Journal of Agroindustri Engineering and Management*, 5 (1), 60-70.
- Putri, F.T., Sudjarwo, E., & Sjoefjan, O. (2018). The Effect of dietary candlenut powder on blood profile and meat cholesterol content of broilers. *Agripet*, 18 (1), 63-66.
- Quintao, N.L., Meyre-Silva, C., Silva, G.F., Antonioli, C. S., Rocha, L.W., Lucinda-Silva, R.M., Malheiros, A., Souza, M.M., Filho, V.C., & Tania, M.B. (2011). *Aleurites moluccana* (L.) Willd. Leaves: Mechanical Antinociceptive Properties of a Standardized Dried Extract and Its Chemical Markers. *Evidence-Based Complementary and Alternative Medicine*, 2011.
- Raharjo, T.J., Nurliana, L., & Mastjeh, S. (2011). Phospholipids from pumpkin (*Cucurbita moschata* (Duch.) Poir) seed kernel oil and their fatty acid composition. *Indonesian Journal Chemistry*, 11 (1), 48-52.
- Ramdhani, A., Ramdhani, M.A., & Amin, A.S. (2014). Writing a literature review research paper: A step-by-step approach. *International Journal of Basics and Applied Science*, 3 (1), 47-56.
- Rasid, R.A., Baba, A.R., Yaakub, N.M., & Milan, A.R. (2019). Performance and carcass characteristics of broiler chickens fed various components of candlenut kernel. *Tropical Animal Science Journal*, 42 (3), 203-208.
- Rohaida, A.R., Alimon, A.R., & Sazili, A.Q. (2014). Fatty acid composition of breast and thigh muscles of broiler fed diets supplemented with candlenut kernel meal subjected to different heat treatments. *Malaysian Journal Animal Science*, 17, 47-60.
- Russel, A.D. (1991). Mechanisms of bacterial resistance to non-antibiotics: food additives and food pharmaceutical preservatives. *Journal Applied Bacteriology*, 71, 191-201.
- Seidel, V., & Taylor, P.W. (2004) In vitro activity of extracts and constituents of *Pelagonium* against rapidly growing mycobacteria. *International Journal of Antimicrobial Agents*, 23, 613-619.
- Sethi, J., & Dahiya, K. (2018). *Myristica Fragrans* (MF): potential role as an antioxidant and anti-inflammatory agent. *Journal of Natural & Ayurvedic Medicine*, 2 (2), 000120.
- Siegmund, B., & Murkovic, M. (2004). Changes in chemical composition of pumpkin seeds during the roasting process for production of pumpkin seed oil. 2. Volatile compounds. *Food Chemistry*, 84, 367-374.
- Sinaga, R. (2016). Physical and mechanical characteristics of candlenut (*Aleurites molucca* Wild.). *Journal Engineering Agriculture*, 4 (1), 97-106.
- Stevenson, D.G., Eller, F.J., Wang, L., Jane, J.L., Wang, T., & Inglett, G.E. (2007). Oil and tocopherol content and composition of pumpkin seed oil in 12 cultivars. *Journal of Agriculture Food Chemistry*, 55, 4005-4013.
- Sun, C.Q., O' Connor, C.J., & Robertson, A.M. (2003). Antibacterial actions of fatty acids and monoglycerides against *Helicobacter pylori*. *FEMS Immunology Medicine Microbiology*, 36, 9-17.
- Tabari, M.A., Ghazvinian, K.H., Irani, M., & Molaei, R. (2016). Effect of dietary supplementation of nettle

- root extract and pumpkin seed oil on production traits and intestinal microflora in broiler chickens. *Bulgary Journal Veterinary Medicine*, 19, 108-116.
- Tajuddin, S., Ahmad, A., Latif, I.A., & Qasmi, A. (2003). Aphrodisiac activity of 50% ethanolic extracts of *Myristica fragrans* Houtt. (nutmeg) and *Syzygium aromaticum* (L) Merr. & Perry. (clove) in male mice: a comparative study. *BMC Complementary Alternative Medicine*, 3, 6-10.
- Thomas, R.A., Krishnakumari, S. (2016). Lipid lowering effects of *Myristica fragrans*. *International Journal of Pharmacology and Pharmaceutical Sciences*, 3 (3), 10-13.
- Thormar, H. (2010). *Lipids and Essential Oils as Antimicrobial Agents*. New York, USA: John Wiley & Sons: Hoboken Publishing House.
- Yoon, B.K., Jackman, J.A., Valle-González, E.A., & Cho, N.J. (2018). Antibacterial free fatty acids and monoglycerides: biological activities, experimental testing, and therapeutic applications. *International Journal of Molecular Sciences*, 19 (1114), 1-40.
- Zhang, W.K., Tao, S.S., Li, T.T., Li, Y.S., Li, X.J., Tang, H.B., Cong, R.H., Ma, F.L., & Wan, C.J. (2016). Nutmeg oil alleviates chronic inflammatory pain through inhibition of COX-2 expression and substance P release in vivo. *Food Nutrition Research*, 60, 30849.
- Zheng, C.J., Yoo, J.S., Lee, T.G., Cho, H.Y., Kim, Y.H., & Kim, W.G. (2005). Fatty acid synthesis is a target for antibacterial activity of unsaturated fatty acids. *FEBS Letters*, 579, 5157-5162.

THE OMEGA-3 AND OMEGA-6 FATTY ACIDS POTENTIAL OF PUMPKIN, CANDLENUT AND NUTMEG SEEDS AS PHYTOADDITIVE FOR POULTRY. A REVIEW

ORIGINALITY REPORT

23%

SIMILARITY INDEX

16%

INTERNET SOURCES

17%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	www.mdpi.com Internet Source	1%
2	Moin ud Din, Abid Ali, Muhammad Yasir, Muhammad Idrees Jilani et al. "Chemical Composition and in vitro Evaluation of Cytotoxicity, Antioxidant and Antimicrobial Activities of Essential Oil Extracted from Myristica Fragrans Houtt", Polish Journal of Environmental Studies, 2020 Publication	1%
3	Submitted to Wigan and Leigh College, Greater Manchester Student Paper	1%
4	docplayer.net Internet Source	1%
5	Submitted to East Texas Baptist University Student Paper	1%
6	Submitted to Nguyen Tat Thanh University Student Paper	1%

7	Yashi Srivastava, Anil D. Semwal, Aishwarya Dhiman. "A comprehensive review on processing, therapeutic benefits, challenges, and economic scenario of unconventional oils", Journal of Food Processing and Preservation, 2021 Publication	1 %
8	dokumen.pub Internet Source	1 %
9	www.fins.uns.ac.rs Internet Source	1 %
10	Andrew P. Desbois. "Antibacterial free fatty acids: activities, mechanisms of action and biotechnological potential", Applied Microbiology and Biotechnology, 12/03/2009 Publication	<1 %
11	espace.library.uq.edu.au Internet Source	<1 %
12	hdl.handle.net Internet Source	<1 %
13	R Wafar, M Hannison, U Abdullahi, A Makinta. "Effect of Pumpkin (Cucurbita pepo L.) Seed Meal on the Performance and Carcass Characteristics of Broiler Chickens", Asian Journal of Advances in Agricultural Research, 2017 Publication	<1 %

14	oceanrep.geomar.de Internet Source	<1 %
15	www.biomedcentral.com Internet Source	<1 %
16	www.pakbs.org Internet Source	<1 %
17	Submitted to Natonal Institute of Technology Calicut Student Paper	<1 %
18	Sandra Neđeral Nakić. "Chemical characteristics of oils from naked and husk seeds ofCucurbita pepoL.", European Journal of Lipid Science and Technology, 11/2006 Publication	<1 %
19	citeseerx.ist.psu.edu Internet Source	<1 %
20	McCarthy, G.M.. "Dietary fish oil and rheumatic diseases", Seminars in Arthritis and Rheumatism, 199206 Publication	<1 %
21	T. K. Lim. "Myristica fragrans", Edible Medicinal And Non Medicinal Plants, 2012 Publication	<1 %
22	biblio.ugent.be Internet Source	<1 %
23	peer-reviewedjournals.com Internet Source	<1 %

24	"Innovations in Food Technology", Springer Science and Business Media LLC, 2020 Publication	<1 %
25	Mengjun Sun, Zichao Zhou, Jiachen Dong, Jichun Zhang, Yiru Xia, Rong Shu. "Antibacterial and antibiofilm activities of docosahexaenoic acid (DHA) and eicosapentaenoic acid (EPA) against periodontopathic bacteria", Microbial Pathogenesis, 2016 Publication	<1 %
26	edepot.wur.nl Internet Source	<1 %
27	livrepository.liverpool.ac.uk Internet Source	<1 %
28	sciendo.com Internet Source	<1 %
29	www.academicjournals.org Internet Source	<1 %
30	patents.google.com Internet Source	<1 %
31	pediatrics.aappublications.org Internet Source	<1 %
32	repository.ub.ac.id Internet Source	<1 %
33	vdocuments.site Internet Source	<1 %

34	www.entomoljournal.com Internet Source	<1 %
35	www.semanticscholar.org Internet Source	<1 %
36	Luigi Inguglia, Marco Chiaramonte, Vita Di Stefano, Domenico Schillaci et al. " fish waste oil: Fatty acids composition and antibacterial activity ", PeerJ, 2020 Publication	<1 %
37	Parfene, Georgiana, Vicentiu Horincar, Amit Kumar Tyagi, Anushree Malik, and Gabriela Bahrim. "Production of medium chain saturated fatty acids with enhanced antimicrobial activity from crude coconut fat by solid state cultivation of <i>Yarrowia lipolytica</i> ", Food Chemistry, 2013. Publication	<1 %
38	eprints.nottingham.ac.uk Internet Source	<1 %
39	ietd.inflibnet.ac.in Internet Source	<1 %
40	jast.modares.ac.ir Internet Source	<1 %
41	ph04.tci-thaijo.org Internet Source	<1 %
42	www.slpk.sk Internet Source	<1 %

- 43 Emeka Godwin Anaduaka, Nene Orizu Uchendu, Lawrence Uchenna Sunday Ezeanyika. "Mineral, amino acid and fatty acid evaluations of Myristica fragrans seeds extracts", Scientific African, 2020
Publication <1 %
-
- 44 Gilbert O. Fruhwirth. "Seeds and oil of the Styrian oil pumpkin: Components and biological activities", European Journal of Lipid Science and Technology, 11/2007
Publication <1 %
-
- 45 Maryam Jafari, Sayed Amir Hossein Goli, Mehdi Rahimmalek. "The chemical composition of the seeds of Iranian pumpkin cultivars and physicochemical characteristics of the oil extract", European Journal of Lipid Science and Technology, 2012
Publication <1 %
-
- 46 Mianty C Pakasi, J S Mandey, Y H.S Kowel, M N Regar. "PENGARUH PEMBERIAN TEPUNG LIMBAH LABU KUNING (Cucurbita moschata) DALAM RANSUM TERHADAP PERSENTASE KARKAS, POTONGAN KOMERSIL DAN LEMAK ABDOMINAL PADA AYAM PEDAGING", ZOOTEK, 2020
Publication <1 %
-
- 47 Mohamed Dowidar, Amany Ahmed, Hanaa Mohamed. "The Critical Nutraceutical Role <1 %

of Pumpkin Seeds in Human and Animal Health: An Updated Review", Zagazig Veterinary Journal, 2020

Publication

48

Sado-Kamdem, S.L.. "Effect of @a-linolenic, capric and lauric acid on the fatty acid biosynthesis in Staphylococcus aureus", International Journal of Food Microbiology, 20090228

Publication

<1 %

49

eprints.mui.ac.ir

Internet Source

<1 %

50

iecps2020.sciforum.net

Internet Source

<1 %

51

lrrd.org

Internet Source

<1 %

52

riubu.ubu.es

Internet Source

<1 %

53

www.afst.valahia.ro

Internet Source

<1 %

54

www.frontiersin.org

Internet Source

<1 %

55

www.ocl-journal.org

Internet Source

<1 %

56

Fotoon Sayegh, Ahmed Elazzazy, Stamatia Bellou, Anna Moustogianni, Ayman I. Elkady, Mohammed N. Baeshen, George Aggelis. "Production of polyunsaturated single cell

<1 %

oils possessing antimicrobial and anticancer properties", Annals of Microbiology, 2015

Publication

57

Nahed, Atta, Ghada, Ismaiel, Zein, H.. "Usage of oil and powder of bottle gourd and pumpkin seeds in production of high nutritive value biscuit", Egyptian Journal of Nutrition and Health, 2020

Publication

<1 %

58

Jackman, Joshua, Bo Yoon, Danlin Li, and Nam-Joon Cho. "Nanotechnology Formulations for Antibacterial Free Fatty Acids and Monoglycerides", Molecules, 2016.

Publication

<1 %

59

Juliana Maria de Mello Andrade, Rafaela Marin, Miriam Anders Apel, Maria do Carmo Bassols Raseira, Amélia Teresinha Henriques. "Comparison of the Fatty Acid Profiles of Edible Native Fruit Seeds from Southern Brazil", International Journal of Food Properties, 2012

Publication

<1 %

60

Maryam Haghghi, Leila Baghdadi Zare, Mina Ghiasi. "Biodiesel production from Spirulina algae oil over [Cu(H₂PDC)(H₂O)₂] complex using transesterification reaction: Experimental study and DFT approach", Chemical Engineering Journal, 2022

Publication

<1 %

61

Undurti N. Das. "Essential fatty acids: biochemistry, physiology and pathology", Biotechnology Journal, 2006

Publication

<1 %

62

Zeynep Aksoylu Özbek, Pelin Günç Ergönül. "Cold pressed pumpkin seed oil", Elsevier BV, 2020

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On