

ANALYSIS OF SPERMATOOZOA QUALITY USING PERCOLL DENSITY GRADIENT CENTRIFUGATION THROUGH THE ADMINISTRATION OF PHOSPHOLIPID + EGTA

by Eva L. Baideng 2

Submission date: 28-Jun-2023 12:20PM (UTC+0700)

Submission ID: 2123813216

File name: ATOZOA_QUALITY_USING_PERCOLL_DENSITY_GRADIENT_CENTRIFUGATION.pdf (473.78K)

Word count: 5594

Character count: 30184

ANALYSIS OF SPERMATOZOA QUALITY USING PERCOLL DENSITY GRADIENT CENTRIFUGATION THROUGH THE ADMINISTRATION OF PHOSPHOLIPID + EGTA

ROOIJE R.H. RUMENDE^{1,*}, EVA L. BAI DENG¹,
 FREDINE E.S. RARES², LAYA M. RARES³

*E-mail: rooije.rumende@gmail.com

Received: Oct. 27, 2021. Revised: Dec. 19, 2021. Accepted: Feb. 04, 2022. Published online: Feb. 15, 2022

ABSTRACT. The Percoll density gradient centrifugation (PDGC) method is frequently used in the sexing of spermatozoa. However, this method causes damage to the spermatozoa membranes, resulting in a decreased quality of spermatozoa. We analysed the impacts of phospholipid PC (phosphatidylcholine) and EGTA (ethylene glycol bis (β -aminoethyl ether) N,N,N',N'-tetraacetic acid) Ca²⁺ free buffer on the quality of bovine spermatozoa after the PDGC process, using semen from Friesian Holstein (FH) bulls aged 5 - 8 years. The following variables were observed: spermatozoa motility, spermatozoa viability, spermatozoa membrane integrity, spermatozoa that have not experienced capacitation, spermatozoa that have experienced capacitation and spermatozoa that have undergone acrosomal reaction. The results showed that the administration of phospholipid PC + EGTA Ca²⁺ free buffer to spermatozoa, followed by the PDGC process, could maintain or further improve the values of all variables. In the PDGC process, phospholipid PC 10% + EGTA Ca²⁺ free buffer at 1 mM was most suitable.

Keywords: PDGC; PC phospholipids; EGTA; separation spermatozoa.

INTRODUCTION

Separation of the X and Y chromosomes of spermatozoa can be carried out using the Percoll density gradient centrifugation (PDGC) method. This approach is easy in its implementation and application and a valid and inexpensive method. However, results from several recent studies indicate that the PDGC method may cause damage to spermatozoa membranes, which can further decrease the quality of spermatozoa (Susilawati *et al.*, 2017).

Damage to the spermatozoa membrane after the PDGC process is caused by three factors, namely loss of seminal plasma, the presence of chemical factors characterised by an increase in free radicals, as well as physical factors that occur due to

¹ Department of Biology, Faculty of Mathematics and Natural Sciences, Sam Ratulangi University, Manado, Indonesia

² Section of Microbiology, Faculty of Medicine, Sam Ratulangi University, Manado, Indonesia

³ Section of Ophthalmology, Faculty of Medicine, Sam Ratulangi University, Manado, Indonesia

friction/collision between spermatozoa. Of the three factors, the physical factor is the most important one. It is therefore assumed that once this is fixed, the other factors are also neutralised. To fix the physical factor, the integrity of the spermatozoa membrane, which is composed of lipids, proteins, carbohydrates and other substances, is crucial (Tucker and Jansen, 2002).

Although the PDGC method has been widely used and is proven to be efficient, there are factors that limit the success of separating X and Y spermatozoa, namely the occurrence of damage to the spermatozoa membrane due to mechanical forces during the centrifugation and washing process of spermatozoa, which decreases the quality of spermatozoa. Inappropriate speed and duration of centrifugation during spermatozoa separation will decrease the viability and motility of spermatozoa (Rasad *et al.*, 2019). In addition, centrifugation can cause the generation of Reactive Oxygen Species (ROS), namely an increase in free radicals due to membrane damage, affecting the physiological processes of spermatozoa (Zanella *et al.*, 2016). Rasad *et al.* (2019), who performed sexing using the PDGC method, found a decrease in motility by 39.87% for X spermatozoa and 34.73% for Y spermatozoa. This decrease occurred due to the influence of mechanical forces, the spermatozoa medium and the temperature during centrifugation. In addition, there was an increase in the percentage of abnormal spermatozoa due to collisions on the tube wall during the sexing process, resulting in damage to the morphology of spermatozoa.

To repair the damaged membrane structure, phospholipids have been

administered based on the results of previous studies (Bowyer and Davies, 1979). Research has also been carried out on human spermatozoa by Djauhari (2009) through the addition of phospholipids in tris amine methane diluent, which can maintain the quality of spermatozoa and increase the number of intact spermatozoa, inhibiting premature capacitation. In addition, Rumende *et al.* (2007) suggested that the addition of 10% PPC phospholipid (pure phosphatidyl choline 97.5%), in combination with the PDGC method, improved the quality of spermatozoa.

In bovine spermatozoa, damage to the structure of the spermatozoa membrane on the head, acrosomal hood or tail, caused by the PDGC process, affects the transport of calcium ions into and out of the spermatozoa membrane. An excessive increase in calcium ions in the cell causes the disruption of cell functions, resulting in a decreased quality of spermatozoa. Efforts to inhibit the entry of calcium ions into cells or to modulate the entry of calcium ions into cells, as well as attempts to conduct calcium chelation using various natural materials or more selective synthesis, have yielded promising results in the prevention and treatment of various diseases. For example, ethylene glycol bis (β -aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA) Ca^{2+} -free buffer is a scavenger of extracellular and intracellular Ca^{2+} ions that selectively binds Ca^{2+} with the highest affinity level (150 nM at pH 7.2) (Ellis-Davies and Kaplan, 1994; Lemonnier *et al.*, 2002). Another study has shown that 1 mM EGTA with free Ca^{2+} buffer in endothelial cell culture induced by various agonists, or administering 100 M H_2O_2 , may

cause Ca^{2+} depletion from endoplasmic reticulum depots and inhibit Ca^{2+} infusion from extracellular depots. EGTA is a scavenger of Ca^{2+} ions that selectively binds Ca^{2+} with high affinity (Hu and Corda, 1996; Sadova and Klishin, 2000; Parekh, 2002).

The purpose of this study was to reveal the effect of phospholipid PC (phosphatidyl choline) and EGTA (ethylene glycol bis (β -aminoethyl ether)-N,N,N',N'-tetraacetic acid) Ca^{2+} free buffer on the quality of bovine spermatozoa after PDGC. The existence of X and Y chromosomes in the sexing results was not observed because this study was conducted only on the sexing process spermatozoa and did not measure the separation of X and Y chromosomes. We looked at the impact of the sexing process on the quality of spermatozoa subjected to phospholipid + EGTA treatment. Based on previous studies, spermatozoa resulting from the sexing process generally show a decreased quality.

16 MATERIALS AND METHODS

This research was conducted at the Advanced Biology Laboratory, Department of Biology, Faculty of Math and Natural Sciences, Sam Ratulangi University, Manado, Indonesia. We used semen taken from 10 Friesian Holstein (FH) bulls aged 5 - 8 years and managed by the Sehati Farmers Group in Kakaskasen Dua Village, North Tomohon Sub-district, assisted by the Agriculture and Fisheries Service of Tomohon City. Semen was taken twice from each bull. The criteria for good-quality fresh semen were as follows: per-ejaculate volume ranging from 5 - 8 mL and concentrations between 700 and 2,000 million sperm per mL (Garner and Hafez, 2000; Berden and

Fuquay, 1984). Also, semen with a motility percentage above 70% is more viable than that with lower motility (Susilawati *et al.*, 2017). This has been supported by other studies: motility of at least 70% (Zenichiro *et al.*, 2002), motility > 70% and pH level of 5.9 - 7.3 (Hafez, 2000); viability > 80% (Ax *et al.*, 2000) and membrane integrity level of 72.90% (Check *et al.*, 1992).

This research is a laboratory experimental study using phospholipid PC 10% + EGTA Ca^{2+} free buffer (0.5 mM, 1 mM, 2 mM), administered prior to the PDGC process. After the PDGC, the following variables were observed: spermatozoa motility, spermatozoa viability, spermatozoa membrane integrity, spermatozoa not capacitated, spermatozoa capacitated, spermatozoa subjected to acrosomal reaction. Statistical analysis was done using one-way classification analysis of variance (ANOVA) to determine which level had the best significance for each variable, followed by the LSD test to determine the significance between untreated spermatozoa and control spermatozoa after PDGC. Duncan's multiple distance test was performed to determine the effect of each level on each variable. Data were analysed using the software package SPSS (Statistical Product and Service Solutions) version 22. Significance was set at 0.05 ($p \leq 0.05$).

Percoll density gradient centrifugation method

Making a Percoll density gradient

The density gradients used were 1.036, 1.038, 1.043, 1.047, 1.052, 1.055, 1.057, 1.060, 1.065 and 1.070, obtained by diluting Percoll with Tris aminomethane phospholipid (Djati, 2005: 80 mL Tris aminomethane, 20 mL phospholipid PC) to the density levels of 20, 25, 30, 35, 40, 45, 50, 55, 60 and 65%, respectively. Subsequently, of each sample, an aliquot of 0.5 mL was taken and placed into a tube, arranged sequentially from the highest to the lowest density level.

ANALYSIS OF SPERMATOZOA QUALITY USING PERCOLL DENSITY GRADIENT CENTRIFUGATION

Centrifugation

For this, 1 mL of the sample was placed into a tube containing a Percoll density gradient and centrifuged at 2,250 rpm (850 g) for 5 minutes. The obtained top layer was the seminal plasma, which was removed from the tube and second layer contained the Y spermatozoa. The third layer contained the X spermatozoa, which were taken out and placed into a tube containing 4% FBS in TCM 199, at a total volume of 3 mL, followed by centrifugation at 1,500 rpm (380 g) for 5 minutes. Subsequently, the supernatant was discarded, and a liquid containing the spermatozoa was obtained, of which 1 mL was taken. After treatment via PDGC for 30 minutes, the sample was stained for 15 minutes, followed by observing the quality of the spermatozoa.

Making coloring material

Eosin-Negrosin, as the staining agent in the viability test, was made by dissolving 1 g of eosin, 5 g of negrosin and 3 g of sodium citrate in 100 mL of distilled water.

Chlortetracycline (CTC) was made through the following steps: (1) 250 g of DABCO (Sigma, D-2522) was dissolved in 9 mL of glycerol and incubated in a water bath at 37°C for 3 - 4 hours. After this, 1 mL of PBS was added, and the mixture was stored in a freezer; (2) 0.246 g of Tris (Tris base; Sigma T-1503) and 0.7592 g of NaCl were dissolved in 100 mL of deionised water and stored in a refrigerator; (3) 6.057 g of Tris was dissolved in 59 mL of deionised water and stored in a refrigerator; (4) 12.5 g of paraformaldehyde was dissolved in 50 mL of deionised water, followed by adding 1 M of NaOH until the colour became lighter; (5) the solution obtained from the fourth stage was mixed with 1 M tris buffer solution obtained from the third stage at a ratio of 1:1 (the pH was adjusted to 7.4 with 0.2 M HCl) and stored in a refrigerator; (6) 0.0044 g of CTC powder dye (Sigma, C-7880) was mixed with 0.22 g of L-cysteine (hydrochloric monohydrate); the mixture was mixed with 5 mL of CTC buffer from

the second stage (pH was adjusted to 7.8 with 0.2 M HCl). Finally, three solutions were obtained, namely DABCO solution (stored in a freezer), CTC fixative solution (stored in a refrigerator) and CTC dye solution (stored in a refrigerator).

The hypoosmotic swelling test was formed by making a 150-m-osmos hypoosmotic solution that contained 7.35 g of sodium citrate 2H₂O and 13.52 g of fructose, dissolved in 1,000 mL of distilled water. The phospholipid PPC (pure phosphatidyl choline 97.5%) was obtained from Viva.

Spermatozoa motility observation

Briefly, 10 µL spermatozoa was dropped onto a concave glass object, covered with a glass cover measuring 22 × 22 mm and observed under a light microscope with 400x magnification using computer-assisted semen analysis (CASA). At least 200 motile spermatozoa were observed and divided into four groups: very good movement, good movement, non-progressive movement and not moving. The data obtained were used as a comparison in the observation of spermatozoa motility using CASA (Hafez, 2000).

Spermatozoa viability observations

The semen was dripped with 1 drop of eosin-negrosin solution, placed on a glass object, homogenised, smeared and dried. Observations were made under a light microscope with a magnification level of 400x. As shown in Fig. 1, only living spermatozoa absorbed the colour (Hafez, 2000).

Capacitation observation

The method used to observe capacitation was based on Djati (2005), who modified the Frase method as follows: (1) 100 µL of semen was placed in a 1.5-mL Eppendorf tube and mixed with 45 µL of CTC dye solution. Subsequently, the tube was covered with aluminium foil and vortexed for 9 minutes; (2) 8 µL of CTC fixative was added into the Eppendorf tube, and the mixture was vortexed for 1 minute;

(3) 10 μ L of DABCO was added into the Eppendorf tube and mixed carefully; (4) the Eppendorf tube was then covered with a cover glass and thick tissue paper, which were pressed carefully. The edges of the cover glass were covered with nail polish to protect the solution from drying. Observations were made with an epifluorescence microscope (Nikon) with 400x magnification. The characteristics are shown in Fig. 2. Spermatozoa that had not experienced capacitation showed a light colour on the entire head, whereas spermatozoa that had experienced capacitation showed a light colour only on the upper half of the head. Spermatozoa subjected to acrosomal reaction showed either a light colour or a circular ring appearance at the central part of the head.




Notes:
 (A) Living spermatozoa (do not absorb the dye);
 (B) Dead spermatozoa (absorbed the dye)

31
Figure 1 - Observation of spermatozoa viability with Eosin-Negrosin using a light microscope (OLYMPUS) with 1000X magnification

Observation of membrane integrity with hypoosmotic swelling test

For this, 1 mL of a 150-mL osmotic hypoosmotic solution was added to 0.1 mL of spermatozoa. The mixture was incubated at 37°C for 30 minutes. As shown in Fig. 3, when observed at 400x magnification, there was a characteristic membrane damage,

namely swelling of the spermatozoa or a circular tail end (Correa *et al.*, 1997).



Notes: (A) Uncapacitated spermatozoa; (B) Capacitated spermatozoa; (C) Spermatozoa that have undergone acrosomal reaction

Figure 2 - Observation of the physiological conditions of spermatozoa after CTC staining using epifluorescence microscope (NIKON) with 400x magnification



Notes: (A) The coiled spermatozoa are those with intact membrane integrity; (B) Spermatozoa without a coiled shape are those with damaged membrane integrity

Figure 3 - Observation of the spermatozoa membrane integrity using a light microscope (OLYMPUS)

17 RESULTS AND DISCUSSION

Fresh semen quality

The fresh semen samples that were taken from 10 Friesian Holstein (FH) bulls did not contain abnormal sperm, as shown in Table 1.

Based on the examination of the quality of fresh semen (Table 1), the semen was normal. The per-ejaculate volume ranged from 5 - 8 mL, with concentrations between 700 and 2,000 million sperm

ANALYSIS OF SPERMATOZOA QUALITY USING PERCOLL DENSITY GRADIENT CENTRIFUGATION

per mL (Garner and Hafez, 2000; Berden and Fuquay, 1984). According to the viability indicators, the semen may be used for further treatment.

Effect of Percoll density gradient centrifugation on cow spermatozoa quality


The results of the statistical analysis (Fig. 4) showed that the percentage of

spermatozoa quality for all tested variables, using the PDGC method, decreased significantly (LSD test $p \leq 0.001$), when compared with the control (without PDGC).

In other words, the PDGC method significantly decreased the quality of spermatozoa in all variables studied.

Table 1 - Results of quality inspection of fresh semen taken from 10 Friesian Holstein (FH) bulls before treatment with phospholipid 10%+EGTA Ca²⁺ free buffer (0.5 mM, 1 mM, 2 mM)

Parameter	Mean \pm SD
Volume (ml)	9.71 \pm 0.95
pH	6.30 \pm 0.08
Concentration (10 ⁶)	1518.70 \pm 303.25
Motility (%)	75.00 \pm 0.00
Viability (%)	93.15 \pm 2.78
Membrane integrity (%)	74.99 \pm 4.13
Uncapacitated spermatozoa (%)	54.09 \pm 4.61
Capacitated spermatozoa (%)	31.39 \pm 2.92
Acrosomal reaction spermatozoa	14.51 \pm 4.71




Notes: Motility (M), Viability (V), Membrane Integrity (MI), Not yet capacitated (NYC), Capacitated (C), Acrosomal reaction (AR)

Figure 4 - Percentage \pm SD of all variables before PDGC treatment and after PDGC treatment (LSD test results, $p \leq 0.01$)


Effect of PC phospholipids 10% + EGTA Ca²⁺ free buffer (0.5 mM, 1 mM, 2 mM, respectively) on spermatozoa after PDGC

Based on the results, the administration of 10% PC phospholipid concentration + EGTA Ca²⁺ free buffer (0.5, 1 and 2 mM) in spermatozoa through the PDGC method, the spermatozoa quality in each variable spermatozoa motility (Fig. 5), spermatozoa viability (Fig. 6), spermatozoa membrane integrity (Fig. 7), spermatozoa that have not experienced capacitation (Fig. 8), spermatozoa that have experienced capacitation (Fig. 9) and spermatozoa that have undergone acrosomal reaction (Fig. 10) increased significantly (ANOVA result $p \leq 0.001$), when compared with the control variables. In addition, the administration of PC phospholipid 10% concentration + EGTA Ca²⁺ free buffer at 1 mM resulted in a more significant effect when compared to the other levels (0.5 or 2 mM).




Notes: SP=Before treatment; A=Phospholipids 0% + EGTA Ca²⁺ free buffer (0 mM); B=Phospholipid 10%+ EGTA Ca²⁺ free buffer (0.5 mM); C=Phospholipid 10% + EGTA Ca²⁺ free buffer (1 mM); D=Phospholipids 10% + EGTA Ca²⁺ free buffer (2 mM); (a, b, c); Duncan's multiple distance test results 5%

Figure 5 - Average percentage ± SD of motility before and after treatment with phospholipids + EGTA Ca²⁺ free buffer



Notes: SP=Before treatment; A=Phospholipids 0% + EGTA Ca²⁺ free buffer (0 mM); B=Phospholipid 10%+ EGTA Ca²⁺ free buffer (0.5 mM); C=Phospholipid 10% + EGTA Ca²⁺ free buffer (1 mM); D=Phospholipids 10% + EGTA Ca²⁺ free buffer (2 mM); (a, b, c)=Duncan's multiple distance test results 5%


Figure 6 - Average percentage ± SD viability before and after treatment with phospholipids + EGTA Ca²⁺ free buffer



Notes: SP=Before treatment; A=Phospholipids 0% + EGTA Ca²⁺ free buffer (0 mM); B=Phospholipid 10%+ EGTA Ca²⁺ free buffer (0.5 mM); C=Phospholipid 10% + EGTA Ca²⁺ free buffer (1 mM); D=Phospholipids 10% + EGTA Ca²⁺ free buffer (2 mM); (a, b, c)=Duncan's multiple distance test results 5%


Figure 7 - Average percentage ± SD of membrane integrity before and after treatment with phospholipids + EGTA Ca²⁺ free buffer

ANALYSIS OF SPERMATOZOA QUALITY USING PERCOLL DENSITY GRADIENT CENTRIFUGATION



Notes: SP=5 Before treatment; A=Phospholipids 0% + EGTA Ca²⁺ free buffer (0 mM); B=Phospholipid 10%+ EGTA Ca²⁺ free buffer (0.5 mM); C=Phospholipid 10% + EGTA Ca²⁺ free buffer (1 mM); D=Phospholipids 10% + EGTA Ca²⁺ free buffer (2 mM); (a, b, c)=Duncan's multiple distance test results 5%

Figure 8 - Average percentage ± SD of uncapacitated spermatozoa before and after treatment with phospholipids + EGTA Ca²⁺ free buffer




Notes: SP=5 Before treatment; A=Phospholipids 0% + EGTA Ca²⁺ free buffer (0 mM); B=Phospholipid 10%+ EGTA Ca²⁺ free buffer (0.5 mM); C=Phospholipid 10% + EGTA Ca²⁺ free buffer (1 mM); D=Phospholipids 10% + EGTA Ca²⁺ free buffer (2 mM); (a, b, c)=Duncan's multiple distance test results 5%

Figure 9 - Average percentage ± SD of capacitated spermatozoa before and after treatment with phospholipids + EGTA Ca²⁺ free buffer

Based on the obtained results, the improvement of spermatozoa quality in all study variables after the administration of 10% PC phospholipid concentration + EGTA Ca²⁺ free buffer at either 0.5, 1 or 2 mM was caused by a significant

positive effect of PC phospholipids on spermatozoa quality; the balance of the calcium ion concentration between inside and outside the cell was well maintained through the administration of EGTA Ca²⁺ free buffer. The EGTA Ca²⁺ free buffer functions as a scavenger of extracellular and intracellular Ca²⁺ ions and selectively binds Ca²⁺, with the highest affinity level of 150 nM at pH 7.2.



Notes: SP=5 Before treatment; A=Phospholipids 0% + EGTA Ca²⁺ free buffer (0 mM); B=Phospholipid 10%+ EGTA Ca²⁺ free buffer (0.5 mM); C=Phospholipid 10% + EGTA Ca²⁺ free buffer (1 mM); D=Phospholipids 10% + EGTA Ca²⁺ free buffer (2 mM); (a, b, c)=Duncan's multiple distance test results 5%

Figure 10 - Average percentage ± SD of spermatozoa that have undergone acrosome reaction before and after treatment with phospholipids + EGTA Ca²⁺ free buffer

The results are in agreement with the findings of Ellis-Davies and Kaplan (1994) and Lemonnier *et al.* (2002). In another study, 1 mM EGTA with free Ca²⁺ buffer in an endothelial cell culture induced by various agonists or administration of 100 M H₂O₂ may cause Ca²⁺ depletion from endoplasmic reticulum depots and inhibits Ca²⁺ infusion from the extracellular matrix. In addition, EGTA was able to sharply reduce cytosolic Ca²⁺ with a 24 ng glucose exposure of 7 days. These results are in

accordance with the findings of Hu *et al.* (1988), Sadova and Klishin (2000) and Parekh (2002). This contributed greatly to the increase in cytosolic Ca^{2+} , and on day 7 of exposure, the increase in cytosolic Ca^{2+} was caused by influx from the extracellular matrix. Ellis-Davies and Kaplan (1994) obtained similar results.

Our findings support the hypothesis that the results obtained by other researchers may also be applied to spermatozoa cells. This is based on the understanding that calcium ions (Ca^{2+}) are present in the cell matrix, via molecule adhesion, inside cell membranes and inside cell organelles, either bound to protein receptors in the cytosol or free in the cytosol. The free calcium ion concentration inside the cell is only 10^{-7} M, whereas outside the cell, it is 10^{-3} ³⁵. This huge difference between the free calcium ion concentration in the cytoplasm and in the extracellular fluid causes an abundant flow of calcium ions into the cytoplasm when calcium ion channels both inside the cell and in the organelle membrane are opened. This assumption is supported by Carrafoli *et al.* (1990) and Dedman and Kaetzel (1998).

The concurrent role of PC phospholipids and EGTA Ca^{2+} free buffer in this study showed that the membrane remained stable and could function properly because PC phospholipids maintain and protect the spermatozoa membranes, including building structures and metabolic processes, whereas EGTA Ca^{2+} free buffer functions as a scavenger of extracellular and ¹tracellular Ca^{2+} ions. Synchronisation of the role of PC phospholipids and EGTA Ca^{2+} free buffer minimises the possibility of

extracellular matrix deformation and changes in the ion regulation system in spermatozoa membranes that are caused by the PDGC process. As a results, the membranes may function properly and ultimately, the quality of the spermatozoa is maintained or even improved. The mechanism underlying the increase in calcium ions in the cytoplasm is two-fold: 1) calcium ions enter when the canal is opened due to changes in membrane voltage (voltage-operated calcium channel, VOCC) and 2) calcium ions enter when the canal is opened due to binding of the ligand with the receptor (receptor-operated calcium channel, ROCC).

The opening of ⁴³ these two types of channels will cause the entry of calcium ions from outside the cell into the cell because of the large concentration gradient. The second pathway is by opening the ion channels in the cell organelle membrane. This channel is an ROCC; when a ligand binds to ²³ receptor on the membrane, it will cause the release of calcium ions from the organelle into the cell cytoplasm. Ion channels exist in two forms, namely in the closed and in the open state. In accordance with the findings of Breitbart and Naor (1999), calcium release-activated calcium channel (CRAC) is the main route of calcium entry in spermatozoa that do not have voltage-dependent calcium channels with a capacitive calcium entry (CCE) mechanism. This channel is related with the calcium deposition phase into the internal depot. The emptying ¹⁰ of internal depots appears to increase the calcium permeability in the plasma membrane, which causes the calcium ¹⁰ p₁₀ eau in spermatozoa cells to be highly dependent on the concentration

ANALYSIS OF SPERMATOZOA QUALITY USING PERCOLL DENSITY GRADIENT CENTRIFUGATION

of extracellular calcium. Cytosolic calcium release and IP_3 activation are closely related to the important role of acrosome calcium in regulating various cell functions, which is in line with Breitbart and Spungin (1997).

Furthermore, the incoming calcium will cause the intracellular calcium levels to increase or will fill the calcium storage. In addition, ligand binding to certain receptors will produce second-messenger IP_3 , which will bind to ROCC on the acrosome and cause the release of calcium into the cytoplasm. Increased levels of calcium in the cytosol will inhibit the ability of IP_3 to activate the channel. The increased calcium level in the cytoplasm will enter the mitochondria, where calcium is used for various oxidative phosphorylation enzymes for ATP synthesis.

Calcium will also bind to various protein receptors in cells, including protein kinase C. In the acrosomal membrane, there is also a transport system that requires energy to transport calcium into the acrosome. Calcium from the acrosome will bind to PLC - phosphatidylinositol 4,5-bisphosphate (PIP₂) - diacylglycerol (DAG), opening the plasma membrane so that calcium enters the cytoplasm. Calcium derived from the acrosome will affect CCE activity on the plasma membrane, which can occur through this pathway. Agonists that bind to certain receptors on the cell membrane will activate the PLC enzyme to cleave the PIP₂ precursor molecule into DAG and inositol (1,4,5) triphosphate (IP_3). The DAG, which also acts as a second messenger, will stimulate the activity of protein kinase-C (PKC), whereas IP_3 causes the release of calcium in the acrosome. The mechanism

by which IP_3 releases intracellular calcium involves its interaction with specific receptors on the acrosomal membrane and the opening of acrosomal calcium channels. These results are supported by the findings of Toole *et al.* (1996), Breitbart and Spungin (1997), Breitbart and Naor (1999), Dragileva *et al.* (1999) and Baker *et al.* (2000).

Based on the results, it can be concluded that the optimal phospholipid concentration of PC 10% + EGTA Ca^{2+} free buffer is 1 mM, which can be justified based on the following results: (1) the concentrations of phospholipid PC 10% + EGTA Ca^{2+} free buffer 1 and 2 mM have the same notation for variables: spermatozoa viability, spermatozoa membrane integrity, spermatozoa that have not experienced capacitation, spermatozoa that have experienced capacitation, and spermatozoa that have undergone acrosomal reaction; (2) phospholipid concentration of PC 10% + EGTA Ca^{2+} free buffer at a concentration of 1 mM for spermatozoa motility variable, produced the best notation and was significantly different from the concentration of phospholipid PC 10% + EGTA Ca^{2+} free buffer 2 mM.

CONCLUSIONS

Phospholipid PC 10% + Ca^{2+} free buffer at 0.5, 1 and 2 mM, combined with PDGC, significantly improved the quality of spermatozoa. The optimal concentration regarding all variables was PC 10% + Ca^{2+} free buffer at 1 mM.

It is necessary to develop an effective and accurate method to measure the movement of calcium ions in spermatozoa when treated with 10% PC phospholipid

concentration + EGTA Ca^{2+} free buffer + FURA2-AM.

A proportion of 10% PC phospholipid concentration + EGTA Ca^{2+} free buffer (1 mM) or 10% PC phospholipid concentration + EGTA Ca^{2+} free buffer (2 mM) can be used to obtain productive spermatozoa.

The best concentration of phospholipids found in this study may be recommended for breeding quality livestock with reproductive biology technology.

REFERENCES

- Ax, R.L., Dally, M., Didion, B.A., Lenz, R.W., Love, C.C., Varner, D.D., Havez B. & Bellin, M.E. (2000).** Semen evaluation. In: *Reproduction in Farm Animals*, 7th ed., E.S.E. Hafez and B. Hafez (Eds.), *Lea and Febiger*, Philadelphia, pp. 365-375.
- Baker, H.W.G., Liu, D.Y. & Martic, M.G. (2000).** The human acrosome reaction. *Asian J. Androl.*, 2(3): 172-178.
- Bowyer, D.E. & Davies, P.F. (1979).** Effect of concentration of perfusing free fatty acid on arterial lipid synthesis in perfused normal and atherosclerotic rabbit aortas. *Atherosclerosis*, 31(4):409-419, DOI: 10.1016/0021-9150(78)90136-3
- Berden, H.J. & Fuquay, J.W. (1984).** *Applied animal reproduction*. 2nd ed., *Res. Pub. Co., Inc. Prentice-Hall*, Reston, Virginia.
- Breitbart, H. & Spungin, B. (1997).** The biochemistry of the acrosome reaction. *Mol. Hum. Reprod.*, 3(3): 159-202, DOI: <https://doi.org/10.1093/molehr/3.3.195>
- Breitbart, H. & Naor, Z (1999).** Protein kinase in mammalian sperm capacitation reaction. *Rev. Reprod.*, Sep.4(3): 151-159, DOI: 10.1530/ror.0.0040151
- Carrafoli, E., Chiesi, M. & Gazzotti, P. (1990).** The membrane carriers related, and membrane potential in human endothelial cells. *Circulation*, 13: 1719-1725.
- Check, J.H., Katsoff, D., Rozak, J. & Lurie, D. (1992).** Effect of swim-up, Percoll and Sephadex sperm separation methods on the hypo-osmotic swelling test. *Human Reprod.*, 7.(1): 109-111, DOI: 10.1093/oxfordjournals.humrep.a137540
- Correa, J.R., Heersche, G. & Zavos, P.M. (1997).** Sperm membrane functional integrity and response of frozen-thawed bovine spermatozoa during hypoosmotic swelling test incubation at varying temperatures. *Theriogenology*, 47(3): 715-721, DOI: 10.1016/s0093-691x(97)00029-0
- Dedman, J.R. & Kaetzel, M.A. (2012).** Calcium as an intracellular second messenger. mediation by calcium-binding proteins - Chapter 7, *Cell Physiology Source Book* (4th ed.), 99-109, DOI: 10.1016/B978-0-12-387738-3.00007-X
- Djati, M.S. (2005).** Peran Buffer NaHCO_3 pada BO Medium Dalam Menginduksi Kapasitas, Reaksi Akrosom, Viabilitas Sperma, dan Penetrasi Sperma Terhadap Oosit pada Fertilisasi In vitro Sapi (Role of NaHCO_3 buffer in BO medium in inducing capacitance, acrosome reaction, sperm viability, and sperm penetration of oocytes *in vitro* fertilization of cattle). *Jurnal Ilmu-Ilmu Hayati (Journal of the Biological Sciences)*, Vol. 6(1) (in Indonesian)
- Djahhari, T. (2009).** Pengaruh Penambahan Fosfolipid Terhadap Kualitas Spermatozoa Manusia (Effects of addition of phospholipids on the quality of human spermatozoa). *Saintika Medika: Jurnal Ilmu Kesehatan dan Kedokteran Keluarga (Saintika Medika: Journal of Health and Family Medicine)*, Vol. 5(2) (in Indonesian)
- Dragileva, E., Rubinstein, S. & Breitbart, H. (1999).** Intracellular Ca^{2+} - Mg^{2+} - ATPase regulates calcium influx and acrosomal exocytosis in bull and ram spermatozoa. *Biol. Reprod.*, Vol. 61: 1226-1234.
- Ellis-Davies, G.C. & Kaplan; J.H. (1994).** Nitrophenyl-EGTA, a photolabile

ANALYSIS OF SPERMATOZOA QUALITY USING PERCOLL DENSITY GRADIENT CENTRIFUGATION

- chelator that selectively binds Ca²⁺ with high affinity and releases it rapidly upon photolysis. *Proc. Natl. Acad. Sci.*, 91(1): 187-191, DOI: 10.1073/pnas.91.1.187
- Garner, D.L. & Hafez, E.S.E. (2000).** Spermatozoa and seminal plasma. In: *Reproduction in Farm Animals* (7th ed.), Hafez, B., Hafez, E.S.E. (Eds.), Lea and Febiger, Philadelphia, pp. 96-109.
- Hafez, E.S.E. (2000).** X- and Y- chromosome-bearing spermatozoa. In: *Reproduction in Farm Animals* (7th ed.), Lea & Febiger, Philadelphia, pp. 440-446.
- Hu, Q., Corda, S., Zweier, J.L., Capogrossi, M.C. & Ziegelstein, R.C. (1988).** Hydrogen peroxide induces intracellular calcium oscillation in human aortic endothelial cells. *Circulation*, 97(3): 268-275, DOI: 10.1161/01.cir.97.3.268
- Lemonnier, L., Vitko, Y., Shuba, Y.M., Abeele, F.V., Prevarskaya, N. & Skryma, R. (2002).** Direct modulation of volume-regulation anion channels by Ca²⁺ chelating agents. *FEBS Letter*, 521: 152-156.
- Parekh, A.B. (2002).** The welcome prize lecture. Store-operated Ca²⁺ entry: dynamic interplay between endoplasmic reticulum, mitochondria and plasma membrane. *J Physiology.*, 547: 333-348, DOI: 10.1113/jphysiol.2002.034140
- Rasad, S.D., Setiawan, R., Solihati, N., Widyastuti, R. dan Nugraha, I. (2019).** Derajat Pemulihan dan Persentase Spermatozoa X dan Y Kambing Peranakan Etawah Setelah Separasi dengan Gradient Percoll (Degree of recoverand percentage of spermatozoa X and Y Etawah Crossbreeds after separation with Percoll gradients). *Jurnal Veteriner (Veterinary Journal)*. Vol. 20(1): 14-19 (in Indonesian).
- Rumende, R.R.H., Kalim, H., Widodo, A. dan Djati, M.S. (2007).** Peningkatan Kualitas Spermatozoa Sapi pada Proses Pemisahan Kromosom X dan Y dengan Sentrifugasi Gradien Densitas Percoll Melalui Pemberian Fosfolipid (Improving the quality of bovine spermatozoa in the separation process of X and Y chromosomes by centrifugation of Percoll density gradients by giving phospholipids). *Jurnal Kedokteran Brawijaya (Brawijaya Medical Journal)*. Vol. 23(2): 71-81 (in Indonesian).
- Sadova, M., Klishin, A., Huser, J. & Blatter, L.A. (2000).** Capacitative calcium entry is graded with degree of intracellular calcium store depletion in bovine vascular endothelial cells. *J. Physiol.*, 523(Pt 3): 549-559, DOI: 10.1111/j.1469-7793.2000.t01-3-00549.x
- Susilawati, T., Kusumawati, E.D., Isnaini, N., Yekti, A.P.A., Sudarwati, H. & Ridhowi, A. (2017).** Effect of sexing process using Percoll density gradient centrifugation and frozen on motility and damage to spermatozoa membrane of Filial Ongole. *Atlantis Press, Brawijaya University, Malang, Indonesia*.
- Toole, Ch., Roldan, M.B.O., Hampton, E.R.S. & Fraser, P. (1996).** A role for diacylglycerol in human sperm acrosomal exocytosis. *Mol. Hum. Reprod.*, 2(5): 317-326.
- Tucker, K.E. & Jansen, C.A.M. (2002).** Sperm separation techniques: comparison and evaluation of gradient product. In: *Proceedings 2nd International Workshop for Embryologists: Troubleshooting Activities in the ART Lab.*, R. Basuray and D Mortimer (Eds.).
- Zanella, E., Zanella, R., Poetini, M.R., Marques, M.G., Soares, J.C.M. & Bondan, C. (2016)** Oxidative status of boar semen during storage. *Am. J. Biochem. Biotechnol.*, 12(2): 95-101, DOI: 10.3844/ajbbsp.2016.95.101
- Zenichiro, K., Herliante dan Sarastina (2002).** Intruksi Praktis Teknologi Prosesing Semen Beku pada Sapi. (Technology of frozen semen processing for cattle). *Balai Besar Inseminasi Buatan Singosari, Malang, Indonesia*.

ANALYSIS OF SPERMATOZOA QUALITY USING PERCOLL DENSITY GRADIENT CENTRIFUGATION THROUGH THE ADMINISTRATION OF PHOSPHOLIPID + EGTA

ORIGINALITY REPORT

20%

SIMILARITY INDEX

11%

INTERNET SOURCES

17%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Rooije R. H. Rumende, Fredine E. S. Rares, Laya M. Rares. "Analysis of spermatozoa membrane integrity towards spermatozoa quality in sexing process with SGDP through the addition of phospholipid + EGTA", AIP Publishing, 2023

Publication

3%
- 2 Derubertis, F.R.. "Activation of the renal cortical and hepatic guanylate cyclase-guanosine 3',5'-monophosphate systems by nitrosoureas divalent cation requirements and relationship to thiol reactivity", BBA - General Subjects, 19771025

Publication

1%
- 3 fhpkat.kobv.de

Internet Source

1%
- 4 Constantin LUNGOCI, Carmen Doina JITAREANU, Carmen Simona GHITAU, Teodor ROBU. "INFLUENCE OF FOLIAR FERTILISERS ON BIOCHEMICAL AND PHYSIOLOGICAL PROPERTIES IN NEPETA

1%

RACEMOSA LAM.", Journal of Applied Life Sciences and Environment, 2022

Publication

5	Ronald J. Knox. "Activation of a calcium entry pathway by sodium pyrithione in the bag cell neurons ofAplysia", Journal of Neurobiology, 09/15/2004 Publication	1 %
6	Submitted to Universiti Sains Malaysia Student Paper	1 %
7	www.scribd.com Internet Source	1 %
8	ejournal.unsrat.ac.id Internet Source	1 %
9	hdl.handle.net Internet Source	1 %
10	Topics in the Neurosciences, 1986. Publication	1 %
11	T E Tallei, Y T Linelejan, S D Umboh, A A Adam, Muslem, R Idroes. " Endophytic Bacteria isolated from the leaf of Langusei (Tesym. & De Vr.) and their antibacterial activities ", IOP Conference Series: Materials Science and Engineering, 2020 Publication	1 %
12	vdoc.pub Internet Source	1 %

13 Laurentius Anthony Setiono, Rooije Roogers H. Rumende, Lalu Wahyudi. "Kajian Viabilitas Spermatozoa Sebelum dan Sesudah Sentrifugasi Gradien Densitas Percoll dengan Pemberian Fosfolipid", JURNAL BIOS LOGOS, 2022
Publication 1 %

14 www.researcherslinks.com
Internet Source 1 %

15 Imam Mustofa, Suherni Susilowati, Tri Wahyu Suprayogi, Yudit Oktanella, Djoko Agus Purwanto, Adeyinka Oye Akintunde. "Combination of nanoparticle green tea extract in tris-egg yolk extender and 39 °c thawing temperatures improve the sperm quality of post-thawed Kacang goat semen", Animal Reproduction, 2022
Publication <1 %

16 repository.untad.ac.id
Internet Source <1 %

17 pdffox.com
Internet Source <1 %

18 Yendraliza, Y Hendriyanto, D A Mucra, Zumarni, M Rodiallah. "The recovery rate of Simental spermatozoa frozen of post thawing by using tris dilution with different egg yolks", IOP Conference Series: Earth and Environmental Science, 2019
Publication <1 %

19

Louisa Adomaa OFORI, Eunice NIMO, John Tennyson AFELE, Olivia AGBENYEGA.

"INTEGRATING CASHEW (ANACARDIUM OCCIDENTALE) WITH FOOD CROPS: PERCEPTIONS OF FARMERS AT AMOMA, IN THE KINTAMPO SOUTH DISTRICT OF THE BONO EAST REGION OF GHANA", Journal of Applied Life Sciences and Environment, 2022

Publication

<1 %

20

Submitted to Universitas Sam Ratulangi

Student Paper

<1 %

21

www.mdpi.com

Internet Source

<1 %

22

Hoang Xuan Khoi, Kenji Shimizu, Yoshitaka Yoneda, Itaru Minagawa et al. "Monitoring the reactive oxygen species in spermatozoa during liquid storage of boar semen and its correlation with sperm motility, free thiol content and seasonality", Andrologia, 2021

Publication

<1 %

23

api.intechopen.com

Internet Source

<1 %

24

krishikosh.egranth.ac.in

Internet Source

<1 %

25

www.longdom.org

Internet Source

<1 %

26 Michaut, M.. "Synaptotagmin VI Participates in the Acrosome Reaction of Human Spermatozoa", *Developmental Biology*, 20010715
Publication <1 %

27 Submitted to University College London
Student Paper <1 %

28 Yangyang Kate Jiang, Eleanor A. Medley, Grant W Brown. "Two independent DNA repair pathways cause mutagenesis in template switching deficient *Saccharomyces cerevisiae*", Cold Spring Harbor Laboratory, 2023
Publication <1 %

29 repository.unhas.ac.id
Internet Source <1 %

30 repository.up.ac.za
Internet Source <1 %

31 A Rachmawati, Ismaya, B P Widyobroto, S Bintara, T Susilawati. "Effect of Different Bovine Serum Albumin (BSA) Levels on the Sperm Viability of Ongole Cross Bred Bull during 5°C Storage", *IOP Conference Series: Earth and Environmental Science*, 2020
Publication <1 %

32 Cebers, G.. "Modulation of AMPA/kainate receptors by cyclothiazide increases cytoplasmic free Ca^{2+} and $^{45}Ca^{2+}$ uptake in brain neurons", *European Journal*

of Pharmacology: Molecular Pharmacology,
19950718

Publication

33

D. Liu. "Calcium ionophore-induced acrosome reaction correlates with fertilization rates in vitro in patients with teratozoospermic semen", Human Reproduction, 04/01/1998

Publication

<1 %

34

Fei Liu, Jang-Yen Wu, Dorian Beasley, James A Orr. "TxA2-induced pulmonary artery contraction requires extracellular calcium", Respiration Physiology, 1997

Publication

<1 %

35

Hideo Mogami. "Calcium binding capacity of the cytosol and endoplasmic reticulum of mouse pancreatic acinar cells", The Journal of Physiology, 7/1999

Publication

<1 %

36

P Patriani, Hasnudi, E Mirwandhono, Nurzainah Ginting, U Hasanah. "Spermatozoa separation method with different percoll composition on motility, abnormalities, and mortality of local goat sperm", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

37

Stephen V. Straub, David R. Giovannucci, David I. Yule. "Calcium Wave Propagation in

<1 %

Pancreatic Acinar Cells", The Journal of General Physiology, 2000

Publication

38	khartoumspace.uofk.edu Internet Source	<1 %
39	mro.massey.ac.nz Internet Source	<1 %
40	raf.bioscientifica.com Internet Source	<1 %
41	www.docme.ru Internet Source	<1 %
42	www.jbc.org Internet Source	<1 %
43	www.ncbi.nlm.nih.gov Internet Source	<1 %
44	www.physiology.org Internet Source	<1 %
45	www.science.gov Internet Source	<1 %
46	www.usnsj.com Internet Source	<1 %
47	Grasa, P.. "Sperm survival and heterogeneity are correlated with fertility after intrauterine insemination in superovulated ewes", Theriogenology, 200502 Publication	<1 %

48 Sudhir K. Sopory, Meenakshi Munshi. "Protein Kinases and Phosphatases and Their Role in Cellular Signaling in Plants", *Critical Reviews in Plant Sciences*, 2010
Publication <1 %

49 "Poster Sessions", *Allergy*, 2013.
Publication <1 %

50 Joseph P.Y. Kao. "Caged Molecules: Principles and Practical Considerations", *Current Protocols in Neuroscience*, 11/2006
Publication <1 %

51 Vinod Kumar Yata. "Sperm Sexing and its Role in Livestock Production", Springer Science and Business Media LLC, 2022
Publication <1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On