

# Potential fungi isolated from mangrove ecosystem\_Biodiversitas Journal

*by* Rignolda Djamaluddin 23

---

**Submission date:** 24-Jul-2023 10:23AM (UTC+0700)

**Submission ID:** 2135796372

**File name:** fungi\_isolated\_from\_mangrove\_ecosystem\_Biodiversitas\_Journal.pdf (875.13K)

**Word count:** 5133

**Character count:** 28983

## Potential of fungi isolated from a mangrove ecosystem in Northern Sulawesi, Indonesia: Protease, cellulase and anti-microbial capabilities

AGUS TRIANTO<sup>1,2,\*</sup>, OCKY KARNA RADJASA<sup>1,3</sup>, SUBAGIYO<sup>1</sup>, HARTUTI PURNAWENI<sup>4</sup>,  
MUHAMMAD SYAIFUDIEN BAHRY<sup>3</sup>, RIGNOLDA DJAMALUDIN<sup>5</sup>, AIYEN TJOA<sup>6</sup>, IAN SINGLETON<sup>7</sup>,  
KAREN DIELE<sup>7</sup>, DARREN EVAN<sup>8</sup>

<sup>1</sup>Department Marine Science, Faculty of Fisheries and Marine Science, Universitas Diponegoro. Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275, Central Java, Indonesia. Tel. +62-24-7474698, \*email: agustrianto.undip@gmail.com

<sup>2</sup>Natural Product Laboratory, Centre for Research and Services, Universitas Diponegoro. Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275, Central Java, Indonesia

<sup>3</sup>Tropical Marine Biodiversity Laboratory, Faculty of Fisheries and Marine Science, Universitas Diponegoro. Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275, Central Java, Indonesia

<sup>4</sup>Department Public Administration, Faculty of Social and Politic Science, Universitas Diponegoro. Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275, Central Java, Indonesia

<sup>5</sup>Department of Marine Science, Faculty of Fisheries and Marine Science, Universitas Sam Ratulangi. Jl. Kampus Unsrat, Kleak-Bahu, Manado 95115, North Sulawesi, Indonesia

<sup>6</sup>Faculty of Agriculture, Universitas Tadulako. Jl. Soekarno Hatta Km. 9, Tondo, Mantikulore, Palu 94148, Central Sulawesi, Indonesia

<sup>7</sup>School of Applied Sciences, Edinburgh Napier University, Sighthill Campus, 9 Sighthill Court, Edinburgh, EH11 4BN, UK

<sup>8</sup>School of Natural and Environmental Sciences, Newcastle University, Agriculture Building, King's Road, Newcastle upon Tyne NE1 7RU, UK

Manuscript received: 17 January 2021. Revision accepted: 9 March 2021.

**Abstract.** Trianto A, Radjasa OK, Subagiyo, Purnaweni H, Bahry MS, Djamaludin R, Tjoa A, Singleton I, Diele K, Evan D. 2021. Potential of fungi isolated from a mangrove ecosystem in Northern Sulawesi, Indonesia: Protease, cellulase and anti-microbial capabilities. *Biodiversitas* 22: 1717- 1724. The high and relatively unexplored diversity of fungi present in the mangrove ecosystem represents a source of novel biotechnological importance. This study explored the potential of fungi isolated from the mangrove ecosystems to produce proteases and cellulases (commercially important enzymes) and their ability to inhibit pathogenic *Vibrio* species. Random samples of root, branch, leaf, sediments and litter were collected from 5 different mangrove sites in Manado, North Sulawesi, as a source of fungal isolates. The fungi were isolated on malt extract agar (MEA) and potato dextrose agar (PDA). The isolates were identified mainly based on the molecular methods (18S gene sequence) and examined for their ability to produce proteases, cellulases, and activity against several *Vibrio* species. Altogether 288 species of fungi were isolated from all samples. The fungi, isolated from leaves showed the highest diversity. A fungal isolate 19 Mba-C2-1 *Fusarium equiseti* from *Avicennia* sp. leaf showed the highest protease activity. While, the isolate 19 MT-05-3 *Hypocrea* sp. from sediment had the highest cellulase activity. From the root of *Rhizophora* sp., the isolate 19 MT-04-3 identified as *Trichoderma viride* had the strongest activity against a range of *Vibrio* species. This work indicates the high potential of fungi isolated from mangrove ecosystems as a source of commercially important enzymes and novel antimicrobial compounds.

**Keywords:** Aquaculture diseases, bioprospecting, Eurotiomycetes, fungal enzyme, internal transcribed spacer, Sordariomycetes

### INTRODUCTION

Mangrove is an essential component of our ecosystems and has huge but relatively unexplored biodiversity—particularly fungal biodiversity. These fungi could have huge biotechnology potential for production of enzyme and antibacterial compounds. Cellulase and protease enzymes promising biological prospects for discovering potential biocatalysts for use in hydrolysis of lignocellulosic materials and proteic residues. These enzymes can increase and ensure viable production of second-generation ethanol from different and alternative sources (Immaculatejeyasanta et al. 2011; Ramesh et al. 2014). *Vibrio* is a pathogenic-bacteria found in the environment and community with a high-risk infection (Igbino and Omoruyi 2016).

Manado is the largest coastal population in North Sulawesi, Indonesia, having 4.6% of 161 km<sup>2</sup> area under

forest and mangrove. It is an interesting area to harbor the richness of microorganisms. Mangrove forests are uniquely valuable coastal wetlands in the transition zone between land and sea, which moderates freshwater flows from inland while coping with tidal inundation. They sustain millions of people globally, contributing to their survival and welfare through protection against coastal erosion, provision of food and material for construction and firewood, and through filtering of water-borne pollutants, which improves the water quality (Brown and Djamiluddin 2017; Djamiluddin 2018; Hadika and Karumiasa 2020). Mangrove forests are also globally important carbon sinks with carbon densities exceeding 8 times those typical for terrestrial tropical forests (Hossain 2016). They are considered as high priority habitats in climate change mitigation and adaptation strategies (Nehren et al. 2017; Indarsih and Masruri 2019). Mangrove fungi are known to be rich sources of enzymes and secondary metabolites with

various applications such as cellulase, and antibacterial compounds (Sari et al. 2017; Maitig et al. 2018; Sibero et al. 2018). Many studies have shown the importance of using mangrove-derived fungi. The enzymes derived from mangrove-associated microorganisms have economic value for industrial and medical purposes. Previous study has shown that *Aspergillus niger*, *Halocyphina villosa*, and *Lignicola longirostris* are known to produce protease and cellulase (Immaculatejeyasanta et al. 2011).


Protease is an enzyme that performs proteolysis (protein catabolism by hydrolysis of peptide bonds). Proteases are used in industry, medicine, and daily life, e.g., in drug production, controlling blood clotting, as a substituent of detergents (Kamath et al. 2010). Proteases have been applied in environmental bioremediation of protein-polluted areas (through excess feeds) near fish or shrimp ponds to improve water quality (de Souza et al. 2015). Thus, proteases can also act as biocontrol of pathogens, such as *Vibrio*. Cellulase is an enzyme that can break the cellulose bonds into oligo, di, or mono-saccharides. It breaks cellulose through hydrolysis into simple saccharides called cellulose extrin (Kelecom 2002). Unlike other compounds, cellulose is an abundant natural biopolymer on earth. Microorganisms, such as fungi, produce cellulase to degrade cellulose by hydrolyzing the glycoside linkage of cellulose. Previous studies have reported that six fungi such as *Acremonium* sp., *Alternaria chlamydospora*, *Alternaria* sp., *Aspergillus* sp., *Fusarium* sp., and *Pestalotiopsis* sp., isolated from mangrove root of *Avicennia marina*, to produce cellulase (Maria and Sridhar 2002).

*Vibrio* is one of the notorious pathogens, and a leading cause of shrimp and fish aquaculture disease. Bacteria from the genus *Vibrio* have caused enormous losses in the shrimp and fish industry due to mass death and slowing of the growth rate of the fingerlings. The *Vibrio* also infects people consuming raw or undercooked seafood as a case in Japan where a man has reported to be infected by *V. vulnificus* through fish (Li et al. 2018). The objective of this study was to explore the mangrove-associated fungi as sources of protease and cellulase enzymes and/or anti-*Vibrio* compounds.

## MATERIAL AND METHODS

### Location of sampling sites

The natural mangroves are naturally formed mangrove ecosystems, while the restoration mangroves are the mangrove forests that humans have replanted. Samples were collected from 9 to 11 April 2019, from two natural and three restoration sites viz. Likupang Restoration (MSr), 1°40'33.82"N/125° 3'17.24"E; Likupang Natural (MSn), 1°40'41.76"N/125° 3'14.20"E; Tiwoho Natural (MT), 1°35'57.01"N/124°51'32.25"E; Bawoho Restoration (MBa), 1°34'51.69"N/124°49'3.26"E; Buyat Restoration (MB), 0°50'57.01"N/124°42'27.56"E (Figure 1). The collected samples (leaf, branch, root, and sediments) were put into sterile plastic bags to avoid contamination and brought to the laboratory in a cool-box (4 °C) for further treatment.



**Figure 1.** Map of the study areas in the Manado, North Sulawesi, Indonesia. Mangroves showing five sampling sites; Likupang Restoration, MSr (1); Likupang Natural, MSn (2); Tiwoho Natural, MT (3); Bawoho Restoration, MBa (4); and Buyat Restoration, MB (5)

### Screening for cellulolytic and proteolytic of fungi

A total of 288 fungal isolates were tested for their ability to produce cellulolytic and proteolytic enzymes. For Proteolytic screening, the isolates were inoculated on PDA media supplemented with 1 % skimmed milk powder and incubated for 3-5 days at 30 °C. A clear zone around the colony indicated the presence of protease activity (Bonugli-Santos et al. 2015). Screening of cellulolytic activity was done by inoculating the fungi on the media (containing 1% peptone, 0.05% yeast extract, 1% Carboxymethylulose (CMC), and 3% bacteriological agar) and then incubated for 3-5 days at 30 °C, and transferred into a refrigerator (4 °C) for overnight. Cellulase activity was detected by the presence of clear zone around the fungal colony after the addition of Congo Red. The clear zone diameter was classified into weak (< 2 mm), medium (3-4 mm), and strong (>5mm) based on the proteolytic and cellulolytic activities of fungi.

### Antibacterial assay

The *Vibrio* strains were chosen for antibacterial assay, which was conducted using the overlay method as used in the previous study (Trianto et al. 2019). The fungal isolates were inoculated on MEA media in triplicate. After the

growth of isolates which usually takes 1-7 days depending upon the growth rate, the *Vibrio* containing soft agar was poured onto the plates. The soft agar was composed of (0.3% (w/v), nutrient broth, 1% (w/v) NaCl and 0.7% (w/v) agar), containing one of the indicator strains with concentration of 0.5 McFarland. The following strains were used for antibacterial testing: *Vibrio harveyi*, *V. vulnificus*, and *V. parahaemolyticus*. The plates were incubated at the optimum temperature for bacterial growth ( $37 \pm 2$  °C) for 47 h. The anti-*Vibrio* activity was defined by the presence of clear zones around the bacterial isolates. The clear zone diameter formed was also used to classify the antibacterial activity of fungi into weak (< 2 mm), medium (3-4 mm), and strong (>5mm).

#### Molecular identification of the active fungi

The DNA of active isolates was extracted using the Zymo DNA kit. The universal primer, internal transcribed spacer (ITS) was used for the fingerprint region for fungal barcoding using polymerase chain reaction (PCR) thermal cycler. The PCR mix contained GoTaq® Green Master Mix 12.5µl, ITS 1 primer 0.25–2.5µl, ITS 4 primer 0.25–2.5µl, DNA template 1-5 µl, Nuclease-Free Water to total 30 µl. The thermal cycler setting used was denaturation at 95°C for 1 min; 34 cycles of denaturation at 95°C for 30 s, annealing at 56.1°C for 1 min, extension at 72°C for 1 min; last extension at 72°C for 7 min and cooling at 4°C until recovery of the samples. The PCR products were visualized by electrophoresis process, and sequencing was undertaken at Genetika Science, Jakarta Indonesia, and continued to 1st Base, Malaysia. The results were

compared with other sequences in the NCBI database using BLAST. The phylogenetic tree of sequence results was constructed by MEGA 7.0 (Kumar et al. 2016).

## RESULTS AND DISCUSSION

### Fungal isolates


A total of 288 fungal isolates were collected from 5 different locations (Figure 1) having vegetation of four genera of mangrove, viz. *Sonneratia* sp., *Rhizophora* sp., *Avicennia* sp., and *Lumnitzera* sp. Figure 2 showed the mangrove vegetation in North Sulawesi.

#### Isolation of mangrove-associated fungi


The total number of samples collected was 84 from 5 sites (Figure 1), which is shown in Figure 3. The highest number of samples was collected from Tiwoho Natural with 29 samples, followed by Likupang Restoration and Buyat Restoration locations, from which 17 and 16 samples were collected, respectively. Tiwoho Natural area was the most interesting location with respect to the diversity of mangrove samples obtained. This area is part of Bunaken National Park that is a protected mangrove area. The total number of fungi that were successfully isolated was 288 isolates. The highest amount of fungi was 96 isolates from Tiwoho Natural. The ratio from 5 locations (MSr, MSn, MT, MBa, MB) showed that the ratio of fungal association to the number of the sample was to be around 2.6 to 3.7. Interestingly, the highest ratio of 3.7 was observed in Buyat Restoration.



Figure 2. Samples of the study areas in Manado, North Sulawesi, Indonesia. A. *Sonneratia* sp., B. *Rhizophora* sp., C. *Lumnitzera* sp., D. Root and sediment of *Avicennia* sp., E. Litters, F. Mangrove forest



**Figure 3.** The number of samples and corresponding isolates collected from several locations in North-Sulawesi and their isolate-sample ratio



**Figure 4.** The fungal isolate (%) having cellulolytic and proteolytic activities

#### The cellulolytic and proteolytic activities

It was observed that the fungal isolates had cellulolytic and proteolytic activities, as shown in Figure 4. The percentage of the proteolytic enzyme was 2.1% to 21.1%. Moreover, the range of cellulolytic enzyme was identified

at 11.1% to 31.6%. Each location had distinct potential activity in terms of enzyme production. The location in Bahowo restored and Buyat Restoration had higher production of the enzyme with proteolytic and cellulolytic activities respectively, than other places. Tiwoho Natural ranked lowest in terms of enzyme production. Two other locations, Likupang Natural and Bahowo Restored had a similar result of proteolytic enzyme in range 11.1 to 21.1% and cellulolytic enzyme in range 17.6 to 22.2%.

The fungi isolated from leaves were found to be the most potent sources of the enzymes, where 33.3% of them produced protease and cellulase (Figure 5). Followed by roots, whose isolates produced 25% cellulases and 20% of proteolytic enzyme. Branches produced 5.6% of cellulases and 11.1% of proteolytic enzyme, moreover, sediment isolates produced 26.7% of cellulases and 6.7% of proteolytic enzyme, while, isolates from litters did not have any proteolytic activity.


#### Antibacterial assay

The result of antimicrobial activity of mangrove-fungi associated with *V. harveyi*, *V. vulnificus*, and *V. parahaemolyticus* was shown in Figure 6 and Figure 7. Based on the collection sites, most of them had antimicrobial activity with varying potential. The highest anti-*Vibrio* activity (47.6%) was represented by fungal isolate from Bahowo Restoration which had antimicrobial activity against *V. harveyi*. While, fungal isolates from Likupang Natural had the highest antimicrobial activity (20%) against *V. vulnificus*. A total 45.1% of fungal isolate from Tiwoho Natural had highest antimicrobial activity against *V. parahaemolyticus*. Then, based on the fungal association of the part of mangrove (Figure 7), it was known that most of them had strong antimicrobial activity against *V. harveyi*, *V. vulnificus*, and *V. parahaemolyticus*. There is no antimicrobial activity against *V. vulnificus*, and *V. parahaemolyticus* as observed on the fungal isolates from Litters. The strongest activity was investigated on fungi from sediment against *V. harveyi* at 45.5%.


**Table 1.** Isolate source, BLAST identified potential species and biological activity of fungi associated mangrove as proteolytic, cellulolytic and anti-*Vibrio*

Isolate code	Source of mangrove part	Biological activity			Identified species	ACC number
		Proteolytic	Cellulolytic	Anti- <i>Vibrio</i>		
19 Mba-C2-4	Leaf of <i>Avicennia</i> sp.	+			<i>Pestalotiopsis theae</i>	AY924274.1
19 Mba-C2-1	Leaf of <i>Avicennia</i> sp.	+			<i>Fusarium equiseti</i>	KT459349.1
19 Mba-C1-1	Branch of <i>Avicennia</i> sp.			+	<i>Penicillium citrinum</i>	KT844552.1
19 MSr-B3-4	Root of <i>Sonneratia</i> sp.	+	+		<i>Fusarium equiseti</i>	MF471699.1
19 MSr-B3-5	Root of <i>Sonneratia</i> sp.	+			<i>Pestalotiopsis microspora</i>	KT459349.1
19 MB-B7-4	Litters <i>Rhizophora</i> sp.		+		<i>Nigrospora sphaerica</i>	KC505176.1
19 MSr-B2-3	Leaf of <i>Sonneratia</i> sp.		+		<i>Hypocrea jecorina</i>	MN310399.1
19 MT-10-2	Leaf of <i>Rhizophora</i> sp.		+		<i>Aspergillus aculeatus</i>	MH892845.1
19 MT-04-3	Root of <i>Rhizophora</i> sp.			+	<i>Trichoderma viride</i>	MK841023.1
19 MSr-C4-3	Sediment		+		<i>Diaporthe stewartii</i>	KU204517.1
19 MT-05-3	Sediment		+		<i>Hypocrea</i> sp.	MG711900.1


Noted: Likupang Restoration (MSr); Likupang Natural (MSn); Tiwoho Natural (MT); Bahowo Restoration (MBa); Buyat Restoration (MB)



**Figure 5.** The fungal isolate (%) from various samples having cellulolytic and proteolytic activities



**Figure 6.** Site wise fungal isolate (%) active against *Vibrio harveyi*, *V. vulnificus*, and *V. parahaemolyticus* (%)



**Figure 7.** Source-wise fungal isolate (%) active against *Vibrio harveyi*, *V. vulnificus*, and *V. parahaemolyticus*

**Molecular identification of fungal isolates**

Phylogenetic trees were build using maximum likelihood method with bootstrap replicates of 1000 in MEGA 7.0.26 bioinformatics software. The tree represents phylogenetic diversity between restoration area and natural area. The accession number of each species was written in bold italic style right after the species name. The scale bar at the bottom of the figure represents the distance of evolutionary sequence.

**Biological activity of isolated fungi**

The proteolytic and cellulolytic activity as well as anti-*Vibrio* activity of the fungal isolates, their BLAST identified potential species, source of mangrove parts are shown in Table 1. *Pestalotiopsis theae* and *Fusarium equiseti* isolated from leaf of *Avicennia* sp., *Fusarium equiseti*, *Pestalotiopsis microspora* PKT2, and *Fusarium equiseti* isolated from root of *Sonneratia* sp. were the five fungi which had proteolytic activities. The genus *Pestalotiopsis* causes leaf spots, petiole/rachis blights, and sometimes a bud rot of palms. In other words, unlike the other leaf spot and petiole blight pathogens, which attack either leaf blade or the leaf petiole, *Pestalotiopsis* attacks all parts of the leaf from base to tip (Elliott 2018). *Fusarium equiseti* is commonly found in tropical and subtropical areas, and is considered to be a weak pathogen on cereals and is occasionally to be associated with fusarium head blight infected kernels. Isolation of *Fusarium* species in greater number and frequency may be due to the high nutrient level in the mangrove ecosystem (Selvi and Sivakumar 2013).

**Discussion**

Mangrove ecosystems are a relatively unexplored source of fungal diversity and these fungi represent a potential important commercial reservoir of novel enzymes and compounds with various activities e.g. new antibiotics, larvicides, etc. (Thatoi et al. 2013; Pringgenies et al. 2018; Sibero et al. 2018). Bonugli-Santos et al. 2015 reported that mangrove associated fungi are endowed with rich sources of enzymes with biotechnological application, such as proteolytic and/or oxidative enzymes, with alginate lyase, amylase, cellulase, chitinase, glucosidase, inulinase, keratinase, ligninase, lipase, nuclease, phytase, protease, and xylanase.

This work focused on isolating fungi from natural and restored mangrove systems in Northern Sulawesi and it was thought that the different sites could harbor different species due to variation in mangrove species, tidal systems, and restoration methods used. A total of 288 species with the highest diversity found in Tiwoho Natural. Fungi in classes Sordariomycetes and Eurotiomycetes were among the most prevalent in mangrove environments which is in agreement with previous studies (Lee et al. 2019). Previous study has reported the diversity of fungi associated with different parts such as fruits, leaves, pneumatophores, and sediments. They reported that fruit and leaf is the highest amount. On average, our study showed that fungi isolate produced protease and cellulose enzymes. The fungal association from leaf showed the highest production of proteolytic and cellulolytic enzymes (Figure 5). Mangrove fungal cellulolytic activity is reported to be affected by environmental conditions (pH, temperature, substrates), fungal community, and culture conditions (Hossain 2016). Fungi that have proteolytic activity have the ability to produce protease enzymes that are secreted into their environment. The proteolytic enzyme works to hydrolyze protein compounds into oligopeptides, short-chain peptides and amino acids. This extracellular protease enzyme is very important for bacterial life because it provides nitrogen

compounds that can be transported into cells. The types of fungi that can secrete protease have great potential to be used as a source of aquaculture probiotics, especially in shrimp farming (Setyati et al. 2016).


Our study showed Likupang has complex aquatic dynamics. Likupang-restored mangrove area had abandoned shrimp pond with water rich in protein and cellulose from the waste of shrimp and unconsumed feed. The abundance of pollutant sources from the ponds provides a source of nutrients needed by microbes, therefore Likupang-restored had a high number of isolates and isolate-sample ratio (Saiya and Katoppo 2015; Ruete et al. 2016). Moreover to the contrary, it was surprising that in Likupang-Natural has the lowest ratio of isolates to samples. Sridhar and Seetharam (2001) reported loss of over 300 species of aquatic hyphomycete species due to negative effect of polluted water on fungal diversity (18). Laboratory experiments, water pollutants that contain low concentrations of Cd, Cu, and Zn have been shown to inhibit growth and reproduction of aquatic hyphomycetes and fungi respond by synthesizing specific stress peptides (Krauss et al. 2011).

Mangrove leaves were the best source of fungal isolates producing protease and cellulase enzymes (Figure 5). The study of Chi et al. (2019) revealed that fungi associated with leaves of mangrove had a highly diverse fungal community, where a total of 110 taxa were recovered from isolation and metabarcoding methods; among them, Ascomycota was dominant, which includes *Corynespora cassiicola* (6.90%), *F. oxysporum* (6.40%) and *Guignardia* sp. (6.40%). Only specifically fungal isolates sourced from

sediments reported anti-*Vibrio* activity. The fungi that live in seawater sediment are reported to grow well than in freshwater. Sediment nutrient content is affected by physiology and environment among the mangrove vegetation. Previous studies have reported that fungi from different locations, bottom sediment and depth, identified Phycmycetes, Ascomycetes, and Deuteromycetes (Sikumar 2013).

Fungi use mechanisms like extracellular precipitation, valence transformation and active uptake (e.g. bio-sorption to cell wall and pigments, intracellular compartmentation, complexation and crystallization, and sequestration) and therefore could be used to degrade, accumulate or remove metal pollutants. Thus, screening of metal tolerant fungi has the potential of providing strains with improved metal accumulation. However, the search for microorganisms capable of metal bio-sorption and sequestration has mainly focused on contaminated sites (Ojuederie and Babalola 2017; Igiri et al. 2018).

An interesting fungi *Hypocrea jecorina* have been found in litters of *Rhizophora* sp. with cellulolytic enzymatic activity. *Hypocrea jecorina* (anamorphic *Trichoderma reesei*) is a saprophyte noted for its ability to abundantly secrete native hydrolytic enzymes. These enzymes are used in various industrial applications, such as pulp and paper production, and the food and feed industries, and the textile industry (Steiger et al. 2011). *Penicillium citrinum* and *Trichoderma viride* have been successfully isolated from branch of *Avicennia* sp. and root of *Rhizophora* sp., respectively.



**Figure 8.** Phylogenetic trees constructed with sequence of active isolates along with BLAST derived sequences

We showed anti-*Vibrio* activity of *Penicillium citrinum* and *Trichoderma viride*. Previous study has shown four new compounds, penicitrinone E, penicitrinol J, penicitrinol K, and citrinolactone D, that were isolated together with six known compounds from the marine-derived *Penicillium* sp. ML226 (Wang et al. 2013). While, penicitrinone E, penicitrinol J and penicitrinol K showed modest selective cytotoxicity against HepG-2 cell line, citrinolactone D showed weak cytotoxicity against HepG-2 and HeLa cell lines. penicitrinol J and penicitrinol K also showed mild antimicrobial activity against *Staphylococcus aureus* (Wang et al. 2013).

### ACKNOWLEDGEMENTS

This work was supported by grants from the DRPM-Ministry of Research Technology and Higher Education, Indonesia through Basic Research Scheme with contract number: No. 201-01/UN7.P4.3/PP/2019. and No. 257-15/UN7.P4.3/PP/2019.

### REFERENCES

Bonugli-Santos RC, dos Santos Vasconcelos MR, Passarini MRZ, Vieira GAL, Lopes VCP, Mainardi PH, dos Santos JA, Duarte L de A, Otero IVR, Yoshida AM da S, Feitosa VA, Pessoa Jr A, Sette LD. 2015. Marine-derived fungi: Diversity of enzymes and biotechnological applications. *Front Microbiol* 6: 269. DOI: 10.3389/fmicb.2015.00269.

Brown B, Djalaluddin R. 2017. A site history and field guide for ecological mangrove rehabilitation in Tiwoho Village, Bunaken National Marine Park, North Sulawesi, Indonesia. *Center Int For Res* 1-40. DOI: 10.17528/cifor/006397.

Chi WC, Chen W, He CC, Guo SY, Cha HJ, Tsang LM, et al. 2019. A highly diverse fungal community associated with leaves of the mangrove plant *Acanthus ilicifolius* var. *xiamenensis* revealed by isolation and metabarcoding analyses. *PeerJ* 7: e7293. DOI: 10.7717/peerj.7293.

Djalaluddin R. 2018. The mangrove flora and their physical habitat characteristics in Bunaken National Park, North Sulawesi, Indonesia. *Biodiversitas* 19 (4): 1303-1312. DOI: 10.13057/biodiv/d190417.

Elliott ML. 2018. *Pestalotiopsis* (Pestalotia) Diseases of Palm. IFAS Extension, University of Florida, Miami, FL.

Hadika A, Karuniasa M. 2020. Mangrove's vegetation structure and composition (a study: Manado City, North Sulawesi Province). In: Saiya HG et al. (eds) *Proc Int Conf Environ Sci Sustain Develop*, 22-23 October 2019, Jakarta, Indonesia. DOI: 10.4108/eai.22-10-2019.2291477.

Hossain MM. 2016. On identifying the SARIMA model to forecast the humidity of some selected stations in Bangladesh. *Int J Environ Sci* 6: 1080-1089. DOI: 10.6088/ijes.6102.

Igbino EO, Omoruyi G, Idemudia OG. 2016. Anti-vibrio potentials of acetone and aqueous leaf extracts of *Ocimum gratissimum* (Linn). *Trop J Pharm Res* 15 (4): 743-750. DOI: 10.4314/tjpr.v15i4.11.

Igiri BE, Okoduwa SIR, Idoko GO, Akabuogu EP, Adeyi AO, Ejiogu IK. 2018. Toxicity and bioremediation of heavy metals contaminated ecosystem from tannery wastewater: A review. *J Toxicol* 2018: 25680386. DOI: 10.1155/2018/2568038.

Immaculatejeyasanta K, Madhanraj P, Patterson J, Panneerselvam A. 2011. Case study on the extracellular enzyme of marine fungi associated with mangrove driftwood of Muthupet Mangrove, Tamil Nadu, India. *J Pharmacy Res* 4: 1385-1387.

Indarsih R, Masruri MS. 2019. Mangrove conservation as an abrasion strategy risk reduction based on ecosystem in the coastal area of the Rembang Regency. *IOP Conf Ser: Earth Environ Sci* 271: 012021.

Kamath P, Subrahmanyam VM, Rao JV, Raj PV. 2010. Optimization of cultural conditions for protease production by a fungal species. *Indian J Pharm Sci* 72 (2): 161-166. DOI: 10.4103/0250-474X.65017.

Kelecom A. 2002. Secondary metabolites from marine microorganisms. *An Acad Bras Ciênc* 74: 151-170. DOI: 10.1590/s0001-37652002000100012.

Krauss F, Bärlocher F, Krauss GJ. 2011. Effects of pollution on aquatic hyphomycetes. In: Tsui CKM, Hyde KD (eds) *Freshwater Mycology*, Fungal Diversity Press, Hong Kong.

Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 33 (7): 1870-1874. DOI: 10.1093/molbev/msw054.

Lee NLY, Huang D, Quek ZBR, Lee JN, Wainwright BJ. 2019. Mangrove-associated fungal communities are differentiated by geographic location and host structure. *Front Microbiol* 10: 2456. DOI: 10.3389/fmicb.2019.02456.

Li L, Wang L, Zhang C, Chen P, Luo X. 2018. A case of *Vibrio vulnificus* related wound infection diagnosed by next-generation sequencing. *IDCases* 15: e00497. DOI: 10.1016/j.idcr.2019.e00497.

Sari SLA, Setyaningsih R, Wibowo NFA. 2017. Isolation and screening of cellulolytic fungi from *Salacca zolacca* leaf litter. *Biodiversitas* 18 (3): 1282-1288. DOI: 10.13057/biodiv/d180355.

Maitig AMA, Alhoot MAM, Tiwari K. 2018. Isolation and screening of extracellular protease enzyme from fungal isolates of soil. *J Pure Appl Microbiol* 12 (4): 2059-2067. DOI: 10.22207/JPAM.12.4.42.

Maria GL, Sridhar KR. 2002. Richness and diversity of filamentous fungi on woody litter of mangroves along the west coast of India. *Curr Sci* 83 (12): 1573-1580.

Nehren U, Ho Dac Thai H, Trung ND, Raedig C, Alfonso S. 2017. Sand dunes and mangroves for disaster risk reduction and climate change adaptation in the coastal zone of Quang Nam Province, Vietnam. In: Nauditt A, Ribbe L (eds) *Land Use and Climate Change Interactions in Central Vietnam*. Water Resources Development and Management. Springer, Singapore. DOI: 10.1007/978-981-10-2624-9\_13.

Ojuederie OB, Babalola OO. 2017. Microbial and plant-assisted bioremediation of heavy metal polluted environments: A review. *Int J Environ Res Public Health* 14 (12): 1504. DOI: 10.3390/ijerph14121504.

Pringgienies D, Widiyadmi R, Yudiati E, Bahry MS, Djunaedi A. 2018. Potentials of mangrove fruit extract *Xylocarpus granatum* for *Aedes aegypti* mosquito repellent. *J Trop Mar Sci* 1: 1-6. DOI: 10.33019/jour.trop.mar.sci.v1i1.657. [Indonesian]

Ramesh K, Natarajan M, Sridhar H, Uma Vanitha M, Umamaheswari S. 2014. Anti-vibrio activity of mangrove and mangrove associates on shrimp pathogen, *Vibrio harveyi* VSH5. *Global Veterinaria* 12 (2): 270-276. DOI: 10.5829/idosi.gv.2014.12.02.82211

Ruete A, Snäll T, Jönsson M. 2016. Dynamic anthropogenic edge effects on the distribution and diversity of fungi in fragmented old-growth forests. *Ecol Appl* 26 (5): 1475-1485. DOI: 10.1890/1512-71.

Saiya HG, Katoppo DR. 2015. Waste management of shrimp farms as starting point to develop integrated farming systems (case study: Kuwaru Coast, Bantul, Yogyakarta, Indonesia). *J Degrade Min Land Manage* 3: 423-432. DOI: 10.15243/jdmlm.2015.031.423.

Selvi KV, Sivakumar T. 2013. Isolation, identification and characterization of *Fusarium* species from mangrove habitat of Pichavaram, Tamil Nadu, India. *Int J Curr Microbiol Appl Sci* 2: 33-49.

Setyati WA, Habibi AS, Subagiyo, Ridlo A, Nirwani S, Pramesti R. 2016. Skrining dan seleksi bakteri simbiosis spongs penghasil enzim ekstraseluler sebagai agen bioremediasi bahan organik dan biokontrol vibriosis pada budidaya udang. *Jurnal Kelautan Tropis* 19: 11-20. DOI: 10.14710/jkt.v19i1.595. [Indonesian]

Sibero MT, Herdikiawan D, Radjasa OK, Sabdono A, Trianto A, Triningsih DW. 2018. Antibacterial activity of sponge-associated fungi against *Vibriosis* agents in shrimp and its toxicity to *Litopenaeus vannamei*. *AAEL Bioflux* 11: 10-18.

Sivakumar T. 2013. A review on biodiversity of marine and mangrove fungi. *Int J Curr Res Aca Rev* 1 (1): 26-44.

de Souza PM, de Assis Bittencourt ML, Caprara CC, de Freitas M, de Almeida RPC, Silveira D, Fonseca YM, Filho EXF, Jr Pessoa A, Magalhães PO. 2015. A biotechnology perspective of fungal proteases. *Braz J Microbiol* 46 (2): 337-346. DOI: 10.1590/S1517-83824620140359.



- Sridhar KR, Seetharam RN. 2001. Aquatic hyphomycetes and leaf litter processing in unpolluted and polluted habitats. In: Misra JK, Hom BW (eds) *Trichomyces and other fungal groups*. Science Publishers, NH, USA..
- Steiger MG, Vitikainen M, Uskonen P, Brunner K, Adam G, Pakula T, Penttilä M, Saloheimo M, Mach RL, Mach-Aigner AR. 2011. Transformation system for *Hypocrea jecorina* (*Trichoderma reesei*) that favors homologous integration and employs reusable bidirectionally selectable markers. *Appl Environ Microbiol* 77 (1): 114-121. DOI: 10.1128/AEM.02100-10.
- Thatoi H, Behera BC, Mishra RR. 2013. Ecological role and biotechnological potential of mangrove fungi : A review. *Mycology* 4 (1): 54-71. DOI: 10.1080/21501203.2013.785448.
- Trianto A, Sabdono A, Radjasa OK, Pramesti R, Putrajaya NTS, Bahry MS, Triningsih DW, Sulistiowati, Afriyanto R. 2019. Optimization production of antifungal substance from a sponge-associated *Trichoderma harzianum* cultivated in the tofu dregs and rice bran. *Res J Biotechnol* 14: 68-73.
- Wang ML, Lu CH, Xu QY, Song SY, Hu ZY, Zheng ZH. 2013. Four new citrinin derivatives from a marine-derived *Penicillium* sp. fungal strain. *Molecules* 18 (5): 5723-5735. DOI: 10.3390/molecules18055723.

# Potential fungi isolated from mangrove ecosystem\_Biodiversitas Journal

---

## ORIGINALITY REPORT

---

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

15%

PUBLICATIONS

3%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1 N Z Zahari, A Calder, P M Tuah, F N Cleophas, F A F Adnan, K Budin, S A M Ali, S A Rahim. "Potential of Wild Fungi (*Amanita princeps* & *Tylopilus felleus*) in Mycoremediation of Selected Heavy Metals in Soil", IOP Conference Series: Earth and Environmental Science, 2021  
Publication 1%
  - 2 Rhesi Kristiana, Diah Ayuningrum, Macellyne Yohanna, Dio Dirgantara et al. "Characterization and identification of antibacterial compound from *Pseudoalteromonas piscicida* associated with *Chromodoris lochi*", AIP Publishing, 2019  
Publication 1%
  - 3 [repo.unsrat.ac.id](http://repo.unsrat.ac.id)  
Internet Source 1%
  - 4 [www.tandfonline.com](http://www.tandfonline.com)  
Internet Source 1%
-

5	B.C. Behera, B.K. Sethi, R.R. Mishra, S.K. Dutta, H.N. Thatoi. "Microbial cellulases – Diversity & biotechnology with reference to mangrove environment: A review", Journal of Genetic Engineering and Biotechnology, 2017 Publication	1 %
6	amb-express.springeropen.com Internet Source	1 %
7	peerj.com Internet Source	1 %
8	repositorio.utfpr.edu.br:8080 Internet Source	1 %
9	Rignolda Djameluddin, Marco Fusi, Brama Djabar, Darren Evans et al. "Point of (no) return? Vegetation structure and diversity of restored mangroves in Sulawesi, Indonesia, 14–16 years on", Restoration Ecology, 2023 Publication	1 %
10	Submitted to Universiti Malaysia Terengganu UMT Student Paper	<1 %
11	Yassir Zouhri, Jean-Michel M. Berthé, Fanny Bonnet, Bernard Stasik et al. "Ferrocene/ferrocenium redox shuttling: Peroxide decomposition catalysis applied to vinyl monomers polymerization under suspension conditions", Polymer, 2022	<1 %

12 [dx.doi.org](https://dx.doi.org) Internet Source <1 %

---

13 R. L. King, S. M. Matheson, E. M. Baxter, S. A. Edwards. "Sow behaviour and piglet weight gain after late cross-fostering in farrowing crates and pens", *animal*, 2020  
Publication <1 %

---

14 [dokumen.pub](https://dokumen.pub) Internet Source <1 %

---

15 [link.springer.com](https://link.springer.com) Internet Source <1 %

---

16 "Etiology and Integrated Management of Economically Important Fungal Diseases of Ornamental Palms", Springer Science and Business Media LLC, 2020  
Publication <1 %

---

17 Submitted to Griffith University Student Paper <1 %

---

18 Patricia Velez, María C. González, Edmundo Rosique-Gil, Joaquín Cifuentes et al. "Community structure and diversity of marine ascomycetes from coastal beaches of the southern Gulf of Mexico", *Fungal Ecology*, 2013  
Publication <1 %

---

19	<a href="http://journals.tubitak.gov.tr">journals.tubitak.gov.tr</a> Internet Source	<1 %
20	<a href="http://www.ecoevoblog.com">www.ecoevoblog.com</a> Internet Source	<1 %
21	Sara Elmahdi, Salina Parveen, Sylvia Ossai, Ligia V. DaSilva, Michael Jahncke, John Bowers, John Jacobs. "Vibrio parahaemolyticus and Vibrio vulnificus Recovered from Oysters during an Oyster Relay Study", Applied and Environmental Microbiology, 2018 Publication	<1 %
22	<a href="http://wikivividly.com">wikivividly.com</a> Internet Source	<1 %
23	Submitted to Higher Education Commission Pakistan Student Paper	<1 %
24	S Jumiayati, M Arsyad, Rajindra, D A T Pulubuhu, A Hadid. "Cocoa based agroforestry: An economic perspective in resource scarcity conflict era", IOP Conference Series: Earth and Environmental Science, 2018 Publication	<1 %
25	<a href="http://assets.researchsquare.com">assets.researchsquare.com</a> Internet Source	<1 %
26	<a href="http://ijseas.com">ijseas.com</a> Internet Source	<1 %

<1 %

27

[rucforsk.ruc.dk](http://rucforsk.ruc.dk)

Internet Source

<1 %

28

"Progress in Mycology", Springer Science and Business Media LLC, 2021

Publication

<1 %

29

Songtao Li, Hongling Zhao, Zhifeng Yin, Shuhua Deng, Yang Gao, Yunfeng Li, Tao Dai. "Synthesis and Antitumor Activity Evaluation of New Phenanthrene-Based Tylophorine Derivatives", Letters in Organic Chemistry, 2019

Publication

<1 %

30

[hdl.handle.net](http://hdl.handle.net)

Internet Source

<1 %

31

Yves M. Galante, Luca Merlini, Tiziana Silveti, Paola Campia, Bianca Rossi, Fiorenza Viani, Milena Brasca. "Enzyme oxidation of plant galactomannans yielding biomaterials with novel properties and applications, including as delivery systems", Applied Microbiology and Biotechnology, 2018

Publication

<1 %

32

[abisys.in](http://abisys.in)

Internet Source

<1 %

33

[eprints.utm.edu.my](http://eprints.utm.edu.my)

Internet Source

&lt;1 %

34

[microbiologyjournal.org](http://microbiologyjournal.org)

Internet Source

&lt;1 %

35

[scholarworks.waldenu.edu](http://scholarworks.waldenu.edu)

Internet Source

&lt;1 %

36

[repository.lppm.unila.ac.id](http://repository.lppm.unila.ac.id)

Internet Source

&lt;1 %

37

"Recent Advancement in White Biotechnology Through Fungi", Springer Science and Business Media LLC, 2019

Publication

&lt;1 %

38

Gyeong Tae Eom, Seung Hwan Lee, Bong Keun Song, Keun-Wo Chung, Young-Wun Kim, Jae Kwang Song. "High-level extracellular production and characterization of *Candida antarctica* lipase B in *Pichia pastoris*", *Journal of Bioscience and Bioengineering*, 2013

Publication

&lt;1 %

39

M. Muwawa Edith, M. Makonde Huxley, C. Obieze Chinedu, M. Jefwa Joyce, H. P. Kahindi James, P. Khasa Damase. "Isolation, characterization and biotechnological potential of tropical culturable rhizospheric fungi from four mangrove species in Kenya",

&lt;1 %

# African Journal of Microbiology Research, 2020

Publication

---

40 [pure.uva.nl](http://pure.uva.nl) <1 %  
Internet Source

---

41 Nalin N. Wijayawardene, Alan J. L. Phillips,  
Diana Santos Pereira, Dong-Qin Dai et al.  
"Forecasting the number of species of  
asexually reproducing fungi (Ascomycota and  
Basidiomycota)", Fungal Diversity, 2022  
Publication

---

42 [journal.frontiersin.org](http://journal.frontiersin.org) <1 %  
Internet Source

---

43 [journal.ipb.ac.id](http://journal.ipb.ac.id) <1 %  
Internet Source

---

44 [vdoc.pub](http://vdoc.pub) <1 %  
Internet Source

---

45 "Approaches in Bioremediation", Springer  
Science and Business Media LLC, 2018  
Publication

---

46 "Industrially Important Fungi for Sustainable  
Development", Springer Science and Business  
Media LLC, 2021  
Publication

---

47 Monica D. Nakei, Gerald Misinzo, Hamisi  
Tindwa, Ernest Semu. "Degradation of <1 %



polyethylene plastic bags and bottles using microorganisms isolated from soils of Morogoro, Tanzania", *Frontiers in Microbiology*, 2022

Publication

---

48

Zhang, Li, Yezhou Liu, Xin Tian, and Zhenhua Tian. "Antimicrobial Capacity and Antioxidant Activity of Enzymatic Hydrolysates of Protein from Rushan Bay Oyster (*C. rassostrea gigas*): Bioactivity of Enzymatic Hydrolysates for Oyster", *Journal of Food Processing and Preservation*, 2014.

Publication

---

<1 %

49

Wei Xu, Shuangshuang Guo, Ka-Lai Pang, Zhu-Hua Luo. "Fungi associated with chimney and sulfide samples from a South Mid-Atlantic Ridge hydrothermal site: Distribution, diversity and abundance", *Deep Sea Research Part I: Oceanographic Research Papers*, 2017

Publication

---

<1 %

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On