BONE DEPROTEINIZATION SKIPJACK (Katsuwonus pelamis L) WITH NaOH IN CONCENTRATION AND TIME DIFFERENCE HYDROLISIS

by Betty Bagau1

Submission date: 21-Aug-2023 08:32AM (UTC+0700)

Submission ID: 2148582037

File name: DEPROTEINIZATION_SKIPJACK_Katsuwonus_pelamis_L_WITH_NaOH_IN.pdf (351.84K)

Word count: 2734 Character count: 15066

UNIVERSITATEA DE ȘTIINȚE AGRICOLE ȘI MEDICINĂ VETERINARĂ "ION IONESCU DE LA BRAD" IAȘI

LUCRĂRI ȘTIINȚIFICE

VOL. 54 MEDICINĂ VETERINARĂ

NR. 1

12

EDITURA "ION IONESCU DE LA BRAD" IAȘI 2011

COLEGIUL DE REDACȚIE

Redactor responsabil:

Prof. univ. dr. Gheorghe SOLCAN

Redactor adjunct:

Prof. univ. dr. Octavian Zaharie OPREAN

Membri:

Prof. univ. dr. Vasile VULPE Conf.univ.dr. Constantin SPĂTARU Conf. univ.dr. Eleonora GUGUIANU Prof. univ. dr. Dan DRUGOCIU

COMISIA DE REFERENȚI ȘTIINȚIFICI

Prof. univ. dr. Octavian Zaharie OPREAN, FMV Iași

Prof. univ. dr. Abdelfatah NOUR – Universitatea Pudue, SUA

of. univ. dr. H.C. Francois CRESPEAU – ENV Alfort, France

Prof. univ. dr. Gheorghe SOLCAN, FMV Iași

grof. univ. dr. Liviu MIRON, FMV Iași

Prof. univ. dr. Gheorghe SAVUTA, FMV Iași

Prof. univ. dr. Gabriel PREDOI - FMV București

rof. univ. dr. Ioan Ștefan GROZA – FMV Cluj Napoca

Prof. univ. dr. Gheorghe DĂRĂBUŞ -FMV Timișoara

Prof. univ. dr. Corneliu COTEA, FMV Iași

Prof. univ. dr. Mihai CARP-CĂRARE, FMV Iași

Conf. univ. dr. Şerban MOROŞAN – INSERM Paris

Prof. univ. dr. H.C. Liviu RUNCEANU, FMV Iași

Prof. univ. dr. Tudor PERIANU, FMV Iași

Prof. univ. dr. Ioan COMAN, FMV Iași

Prof. univ. dr. Elena VELESCU, FMV Iași

Prof.univ.dr. Tita Rostita Balia, Univ. Padjadjaran Bandung

Conf.univ.dr. Hendronoto Lengkey, Univ. Padjadjaran Bandung

Volumul a fost editat cu sprijinul financiar al Ministerului Educației, Cercetării, Tineretului și Sportului

ISSN: 1454-7406

CUPRINS

NR.1

| TUBULAR CHOLANGIOCARCINOMA AT DOG MORPHOLOGICAL ASPECTS OPREAN O.Z., PAȘCA S., LAZĂR M., ROTARU Anca, ANCUȚA Andreea, RAILEANU Gabriela | 7 |
|---|----|
| CYTOKINES AND LIPID PEROXIDATION RESPONSES TO CHRONIC TOXOPLASMA GONDII INFECTION IN PREGNANT WOMEN BEFORE AND AFTER SPIRAMYCIN TREATMENT Afaf, D, Abd elmageid and Abeer, A. Abd el Aleim. | 11 |
| THE CYTOLOGICAL DIAGNOSIS BY FINE NEEDLE ASPIRATION BIOPSY OF DIFFERENT TYPES OF CUTANEOUS AND SUBCUTANEOUS TUMORS ON COMPANY ANIMALS Ancuța Andreea- Roxana, Dragu Corina-Claudia, Oprean O.Z | 19 |
| HISTOLOGICAL AND SEM STUDIES ON THE OSTRICH'S ESOPHAGUS (STRUTHIO CAMELUS L) Attia,H.F; Mazher,K; Al-Kafafy ,M Rashed. R and Abdel-Aziz,A. | 25 |
| BONE DEPROTEINIZATION SKIPJACK (Katsuwonus pelamis L) WITH NaOH IN CONCENTRATION AND TIME DIFFERENCE HYDROLISIS Betty BAGAU and Hendronoto Arnoldus W. LENGKEY | 33 |
| MORPHOLOGICAL, HISTOLOGICAL AND CYTOGENETICS ASPECTS IN MALE PSEUDOHERMAPHRODITES DOMESTIC PIGS (SUS SCROFA DOMESTICA) CIORNEI Cristina, COTEA C.V., SOLCAN Carmen, BEEK Josine, CORNILLIE P. | 37 |
| MORPHOPATHOLOGICAL PECULIARITIES OF NERVES AND ARTERIES INVOLVEMENT IN NECROBACILLARY PODODERMATITIS OF SHEEP Grigore DUMITRAS | 45 |
| MOUSE MODEL FOR HUMAN PLACENTAL STEM CELLS ENGRAFTMENT Groza I.Ş, Pall Emoke, Daria Groza, Cenariu M., Ciupe Simona, Morar I. | 48 |
| SERUM BIOCHEMICAL PROFILE AND KIDNEY'S HISTOLOGICAL CHANGES IN DRUG- INDUCED RENAL FAILURE IN EXPERIMENTAL RATS Nabaa Kamal Al shafei, Afaf Desouky Abd Elmagid , Hussein Abd El-Maksoud, Yakout A. El-senosi, and Abdelfattah Nor | 53 |
| | |

| IMMUNOSUPRESSANT ACTION OF AFLATOXIN B ON PRIMARY LYMPHOID ORGANS IN DUCKLINGS | 69 |
|---|-----|
| Carmen Solcan, Cotea C., Panagachi Gogu Mihaela, Gh. Solcan | |
| BOVINE ADIPONECTIN: EXPRESSION AND REGULATION IN DIFFERENTIATED BOVINE ADIPOCYTES | 79 |
| Mohamed Mohamed Soliman,, Tamer Ahmed Ismaeil,, Hossam Fouad Attia, Omayma Almed Ragab | |
| EFFECT OF LACTOBACILLUS FERMENTED WHEAT BRAN ON RAT LIPID METABOLISM FED CHOLESTEROL ENRICHED DIET LUCIA CARMEN TRINCA, MIRELA ADINA CAPRARIU | 91 |
| LUCIA CARIVIEN TRINCA, IVIIRELA ADINA CAPRARIU | |
| HISTOLOGICAL AND HISTOCHEMICAL STUDY ON THE OVARY OF THE NILE TILAPIA (OREOCHROMIS NILOTICUS) IN REFERENCE TO THE AGE | 96 |
| EL-Zoghby, I.M.A.; Mohammed, R. B. Y; Shaheen, A.A.A and Abdelfattah, N. | |
| HISTOLOGICAL AND IMMUNOHISTOCHEMICAL STUDIES ON THE CLITORIS OF THE DROMEDARY SHE-CAMEL (CAMELUS DROMEDARIUS) | 107 |
| EL-Zoghby, I. M. A. and Emam, M. A. | |
| NEUROLOGICAL DEFICITS AND CHANGES IN THE BRAINSTEM AUDITORY EVOKED POTENTIALS IN THE CENTRAL VESTIBULAR SYSTEM DISORDERS | 119 |
| Armașu Mihaela, Musteață M., Zbângu Adina, Solcan Gh. | |
| INVESTIGATIONS OF BIO-MOS EFFECT ON INDEX NUMBERS OF PRODUCTION AND HEMATOLOGICAL INDEXES IN PIGLETS AND SOWS BALANESCU S. | 125 |
| CLINICAL, PARACLINICAL, ECHOGRAPHICAL AND ELECTROCARDIOGRAPHICAL ASPECTS IN ISCHEMIC CARDIOMYOPATHY OF DOG | 129 |
| V.BOGHIAN, Diana Mocanu, S.I.Beşchea-Chiriac, Luminita Diana Hriţcu, Gh. Solcan | |
| MILK QUALITY AS AFFECTED BY MILKING MACHINE | 134 |
| Cristina BULBAŞA (PANAITE), Ch. HANZEN, L. THERON, D. DRUGOCIU | |
| OBSERVATIONS REGARDING THE CLINICAL ASPECTS AND COMPLICATIONS OF | 137 |
| ANTERIOR UVEITIS IN DOGS AND CATS Ioana BURCOVEANU, I. BURTAN, Roxana TOPALĂ, L.C. BURTAN, M. FÂNTÂNARIU, | |
| CIOBANU | |
| EPIDEMIOLOGY OF DICROCELIOSIS AND CHANGES IN SOME CHEMICAL INDICES OF MUSCULAR TISSUE AND LIVER OF CATTLE IN THE REPUBLIC OF MOLDOVA | 143 |

| PERFORMANCE, FECUNDITY AND SEMEN CHARACTERISTICS IN NILE TILAPIA (OREOCHROMIS NILOTICUS) Eman A. Abd El-Gawad; Mohamed M.M. Kandiel; Amany A. Abbass; and Adel A. Shaheen DETERMINING OF TOXOPLASMOSIS SEROPREVALENCE, BY ELISA, IN LAMBS IN TIMIS COUNTY Ionela Hotea, Marius Ilie, Mirela Imre, Denisa Sorescu, Gheorghe Dărăbuș FOLLICULAR DYNAMICS AND UTERINE STATUS AFTER SYNCHRONIZATION OF OVULATION N EARLY POST-PARTURIENT BUFFALO COWS Sosa, G.A.; Kandiel, M.M.; Gad, B.A. and El-Azab, A.I. THE DIAGNOSTIC AND PROGNOSTIC VALUE OF PERITONEAL FLUID ANALYSIS IN COLIC IN ARABIAN HORSE Wael M. EL-Deeb and Taha A. Fouda NR.2 INVESTIGATION ON HEPATITIS E INFECTION IN PIG FARMS Adriana ANIȚĂ, Luanda LUDU, Dragoș ANIȚĂ, Gheorghe SAVUȚA IDENTIFICATION OF NEW TICK-BORNE ENCEPHALITIS FOCI — A PILOT SEROPREVALENCE STUDY IN THE EAST OF ROMANIA Oana COJOCARU, V. PĂDURARU, B. OLĂERIU, Gh. SAVUȚA THE CONTRIBUTION OF ECOLOGICAL EDUCATION TO BIODIVERSITY CONSERVATION AND ENVIRONMENT PROTECTION VICTOR Donea FERMENTATIONS OF VARIOUS WHEY TYPES WITH USING Kluyveromyces lactis IN THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | | |
|--|--|-----|
| IMPACT OF SOME ORGANOPHOSPHORUS INSECTICIDES ON GROWTH PERFORMANCE, FECUNDITY AND SEMEN CHARACTERISTICS IN NILE TILAPIA (OREOCHEO MIS NILOTICUS) Eman A. Abd El-Gawad; Mohamed M.M. Kandiel; Amany A. Abbass; and Adel A. Shaheen DETERMINING OF TOXOPLASMOSIS SEROPREVALENCE, BY ELISA, IN LAMBS IN TIMIS COUNTY Ionela Hotea, Marius Ilie, Mirela Imre, Denisa Sorescu, Gheorghe Dărăbuș FOLLICULAR DYNAMICS AND UTERINE STATUS AFTER SYNCHRONIZATION OF OVULATION EARLY POST-PARTURIENT BUFFALO COWS Sosa, G.A.; Kandiel, M.M.; Gad, B.A. and El-Azab, A.I. THE DIAGNOSTIC AND PROGNOSTIC VALUE OF PERITONEAL FLUID ANALYSIS IN COLIC IN ARABIAN HORSE Wael M. EL-Deeb and Taha A. Fouda NR.2 INVESTIGATION ON HEPATITIS E INFECTION IN PIG FARMS Adriana ANIȚĂ, Luanda LUDU, Dragoș ANIȚĂ, Gheorghe SAVUȚA IDENTIFICATION OF NEW TICK-BORNE ENCEPHALITIS FOCI — A PILOT SEROPREVALENCE STUDY IN THE EAST OF ROMANIA Oana COJOCARU, V. PĂDURARU, B. OLĂERIU, Gh. SAVUȚA THE CONTRIBUTION OF ECOLOGICAL EDUCATION TO BIODIVERSITY CONSERVATION AND ENVIRONMENT PROTECTION VICTOR Donea FERMENTATIONS OF VARIOUS WHEY TYPES WITH USING Kluyveromyces lactis IN THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | D. Erhan , Ş. Rusu, P. Pavaliuc, O. Chihai, Maria Zamornea, V. Buza, T. Anghel, G. | |
| TIMIS COUNTY Ionela Hotea, Marius Ilie, Mirela Imre, Denisa Sorescu, Gheorghe Dărăbuș COLLICULAR DYNAMICS AND UTERINE STATUS AFTER SYNCHRONIZATION OF OVULATION IN EARLY POST-PARTURIENT BUFFALO COWS Sosa, G.A.; Kandiel, M.M.; Gad, B.A. and El-Azab, A.I. THE DIAGNOSTIC AND PROGNOSTIC VALUE OF PERITONEAL FLUID ANALYSIS IN COLIC IN ARABIAN HORSE Wael M. EL-Deeb and Taha A. Fouda NR.2 INVESTIGATION ON HEPATITIS E INFECTION IN PIG FARMS Adriana ANIȚĂ, Luanda LUDU, Dragoș ANIȚĂ, Gheorghe SAVUȚA IDENTIFICATION OF NEW TICK-BORNE ENCEPHALITIS FOCI – A PILOT SEROPREVALENCE STUDY IN THE EAST OF ROMANIA Oana COJOCARU, V. PĂDURARU, B. OLĂERIU, Gh. SAVUȚA THE CONTRIBUTION OF ECOLOGICAL EDUCATION TO BIODIVERSITY CONSERVATION AND ENVIRONMENT PROTECTION Victor Donea FERMENTATIONS OF VARIOUS WHEY TYPES WITH USING Kluyveromyces lactis IN THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | IMPACT OF SOME ORGANOPHOSPHORUS INSECTICIDES ON GROWTH PERFORMANCE, FECUNDITY AND SEMEN CHARACTERISTICS IN NILE TILAPIA (OREOCHROMIS NILOTICUS) Eman A. Abd El-Gawad; Mohamed M.M. Kandiel; Amany A. Abbass; and Adel A. | 150 |
| FOLLICULAR DYNAMICS AND UTERINE STATUS AFTER SYNCHRONIZATION OF OVULATION BEARLY POST-PARTURIENT BUFFALO COWS Sosa, G.A.; Kandiel, M.M.; Gad, B.A. and El-Azab, A.I. THE DIAGNOSTIC AND PROGNOSTIC VALUE OF PERITONEAL FLUID ANALYSIS IN COLIC IN ARABIAN HORSE Wael M. EL-Deeb and Taha A. Fouda NR.2 INVESTIGATION ON HEPATITIS E INFECTION IN PIG FARMS Adriana ANIȚĂ, Luanda LUDU, Dragoş ANIȚĂ, Gheorghe SAVUȚA IDENTIFICATION OF NEW TICK-BORNE ENCEPHALITIS FOCI – A PILOT SEROPREVALENCE STUDY IN THE EAST OF ROMANIA Oana COJOCARU, V. PĂDURARU, B. OLĂERIU, Gh. SAVUȚA THE CONTRIBUTION OF ECOLOGICAL EDUCATION TO BIODIVERSITY CONSERVATION AND ENVIRONMENT PROTECTION Victor Donea FERMENTATIONS OF VARIOUS WHEY TYPES WITH USING Kluyveromyces lactis IN THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | TIMIS COUNTY | 161 |
| COLIC IN ARABIAN HORSE Wael M. EL-Deeb and Taha A. Fouda NR.2 INVESTIGATION ON HEPATITIS E INFECTION IN PIG FARMS Adriana ANIȚĂ, Luanda LUDU, Dragoș ANIȚĂ, Gheorghe SAVUȚA IDENTIFICATION OF NEW TICK-BORNE ENCEPHALITIS FOCI – A PILOT SEROPREVALENCE STUDY IN THE EAST OF ROMANIA Oana COJOCARU, V. PĂDURARU, B. OLĂERIU, Gh. SAVUȚA THE CONTRIBUTION OF ECOLOGICAL EDUCATION TO BIODIVERSITY CONSERVATION AND ENVIRONMENT PROTECTION Victor Donea FERMENTATIONS OF VARIOUS WHEY TYPES WITH USING Kluyveromyces lactis IN THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | FOLLICULAR DYNAMICS AND UTERINE STATUS AFTER SYNCHRONIZATION OF OVULATION IN EARLY POST-PARTURIENT BUFFALO COWS | 165 |
| INVESTIGATION ON HEPATITIS E INFECTION IN PIG FARMS Adriana ANIȚĂ, Luanda LUDU, Dragoș ANIȚĂ, Gheorghe SAVUȚA IDENTIFICATION OF NEW TICK-BORNE ENCEPHALITIS FOCI – A PILOT SEROPREVALENCE STUDY IN THE EAST OF ROMANIA Oana COJOCARU, V. PĂDURARU, B. OLĂERIU, Gh. SAVUȚA THE CONTRIBUTION OF ECOLOGICAL EDUCATION TO BIODIVERSITY CONSERVATION AND ENVIRONMENT PROTECTION Victor Donea FERMENTATIONS OF VARIOUS WHEY TYPES WITH USING Kluyveromyces lactis IN THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | COLIC IN ARABIAN HORSE | 182 |
| Adriana ANIȚĂ, Luanda LUDU, Dragoș ANIȚĂ, Gheorghe SAVUȚA IDENTIFICATION OF NEW TICK-BORNE ENCEPHALITIS FOCI – A PILOT SEROPREVALENCE STUDY IN THE EAST OF ROMANIA Oana COJOCARU, V. PĂDURARU, B. OLĂERIU, Gh. SAVUȚA THE CONTRIBUTION OF ECOLOGICAL EDUCATION TO BIODIVERSITY CONSERVATION AND ENVIRONMENT PROTECTION Victor Donea FERMENTATIONS OF VARIOUS WHEY TYPES WITH USING Kluyveromyces lactis IN THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | NR.2 | |
| SEROPREVALENCE STUDY IN THE EAST OF ROMANIA Oana COJOCARU, V. PĂDURARU, B. OLĂERIU, Gh. SAVUȚA THE CONTRIBUTION OF ECOLOGICAL EDUCATION TO BIODIVERSITY CONSERVATION AND ENVIRONMENT PROTECTION Victor Donea FERMENTATIONS OF VARIOUS WHEY TYPES WITH USING Kluyveromyces lactis IN THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | | 199 |
| CONSERVATION AND ENVIRONMENT PROTECTION Victor Donea FERMENTATIONS OF VARIOUS WHEY TYPES WITH USING Kluyveromyces lactis IN THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | SEROPREVALENCE STUDY IN THE EAST OF ROMANIA | 203 |
| THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER | CONSERVATION AND ENVIRONMENT PROTECTION | 209 |
| Sunardi, Iwan Setiawan | THE PRODUCTION OF BIOETHANOL AND ORGANIC LIQUID FERTILIZER Gemilang Lara Utama, Tita Roostita Balia, Kurnani Tubagus Benito Achmad, | 214 |

| INVESTIGATION OF FECXI MUTATION IN THREE ROMANIAN SHEEP BREEDS S.E. GEORGESCU, G. HRINCA, Marieta COSTACHE | 220 |
|---|-----|
| INFLUENCE OF GROWTH CONDITIONS ON B.U.T. BIG TURKEY HYBRID AT SLAUGHTER Roxana Lazăr, Alina Narcisa Postolache, Boișteanu P. C. | 224 |
| | |
| EPIDEMIOLOGICAL INVESTIGATION REGARDING PRRSV INFECTION ON PIGLETS FROM NE MOLDAVIAN AREA | 231 |
| MIHAI (MILEA) LEONTINA-SMĂRĂNDIȚA , Elena VELESCU , TOMASZ STADEJEK, TAHAR AIT-ALI | |
| n | |
| EFFECTS OF THE REMEDY BIOR ON THE HEALTH AND PRODUCTIVITY IN YOUNG RABBITS | 236 |
| V.MACARI, Natalia <mark>IACUB</mark> , Ana <mark>MACARI, D</mark> . MAŢENCU | |
| THE EVALUATION OF THE IMMUNE RESPONSE LEVEL INDUCED BY A NEW ANTI – | 241 |
| CONTAGIOUS AGALACTIA VACCINE FORMULATION IN GOATS AND SHEEP | |
| Laurenț OGNEAN,., C. CHIURCIU , F. ZĂVOIU , Cristina CERNEA, P. SEVCIUC, | |
| I.NICOLAE , P. ŞTIUBE , C.TUDORAN , A.NICA , Meda MOLDOVAN | |
| PRION PROTEIN DETECTION IN SMALL RUMINANTS FROM SOUTHEAST ROMANIAN FARMS | 251 |
| MARIA RODICA OTELEA(), STELIAN BARAITAREANU(), MIHAELA ZAULET(), FANEL | |
| OTELEA(), DOINA DANES(,) | |
| THE INFLUENCE OF THE REMEDY BIOR ON THE HEALTH AND PRODUCTIVITY IN | 257 |
| BROILER CHICKS | |
| V. MACARI, V. PUTIN, Natalia PAVLICENCO, Gh. CAZACU | |
| DETECTION OF CIPROFLOXACIN-RESISTANT E.COLI ISOLATES FROM POULTRY | 262 |
| FAECES USING THE EUCAST AGAR DILUTION METHOD | |
| Ramona MORARU, Anne-Marie Pourcher, P. Dabert, V. Nastasa, Mariana Grecu , M. Mares | |
| STUDY ON SOME CHARACTERISTICS OF KOSTA MALE GOAT | 271 |
| Jan Alex SIWI, Enang Wasja ADIKARTA, Primiani EDIANINGSIH | |
| THE MONITORING OF WATER TEMPERATURE, PH, AND DISSOLVED OXYGEN | 274 |
| VARIATION, IN IZVORU-MUNTELUI BICAZ MAN-MADE LAKE, BETWEEN 009-00. | |
| Ramona SORIC, Liviu MIRON, Dumitru ACATRINEI | |

| IMMUNOMODULATING PROPERTIES OF CALENDULA OFFICINALIS AND ECHINACEA ANGUSTIFOLIA EXTRACTS IN ANTIGEN PRIMED HENS Marina SPINU, Carmen SANDRU, Mihaela NICULAE, D. CADAR, Timea KISS, Armela | 278 |
|--|-----|
| BORDEANU, Florina-Alexandra KRUPACI, Silvana POPESCU, R. ŞTEFAN | |
| THE ACTION OF STREPTOMYCETES BIOMASSES AT IMMUNOLOGICAL EFFICACY | 282 |
| AND SOME BIOCHEMICAL BLOOD INDEXES OF VACCINATED CHICKENS | |
| Nicolae STARCIUC, OLGA POSTOLACHI, Ion SCUTARU, Tudor SPATARU, NATALIA | |
| OSADCI, Ruslan ANTOCI | |
| ASSESSMENT OF VIRAL ARTERITIS IMPACT BY SEROLOGICAL EXAM IN HORSES | 287 |
| FROM IASSY COUNTY | |
| Oana Tanase, Constantin Pavli, Remus Plesca, Luanda Ludu, Gheorghe Savuta, | |
| Elena Velescu | |
| EVALUATION OF THE ANTIFUNGAL EFFICIENCY OF ORGANIC ACIDS IN OILSEED | 291 |
| CONSERVATION IN PRODUCTION CIRCUMSTANCES | |
| Alina TEODOR, Ioan COMAN, Daniel TEFELEŞ | |
| PLANT-ANIMAL MODELS FOR HEAVY METALS ACCUMULATION IN AGRICULTURAL | 298 |
| SYSTEMS | |
| Liliana TUDOREANU, Mario CODREANU, Gheorghe V. GORAN, Victor CRIVINEANU | |

BONE DEPROTEINIZATION SKIPJACK (Katsuwonus pelamis L) WITH NaOH IN CONCENTRATION AND TIME DIFFERENCE HYDROLISIS

Betty BAGAU¹ and Hendronoto Arnoldus W. LENGKEY²

¹Faculty of Animal Husbandry University of Sam Ratulangi Manado, Indonesia ²Universitas Padjadjaran, Bandung, Indonesia E-mail betty_bagau@yahoo.com, hawlengkey@yahoo.com

INTRODUCTION

In tuna fish processing business is handled in large and small scale industrial waste generated both solid waste and liquid waste that has not been fully utilized, and the wastes can cause environmental pollution. Tuna has around 61.79% as edible portion, the part that can be eaten, while the remaining 38.21% are wast 2 ncluding bone parts (Prih et al. 1982). According to Lengkey, et al (2011), even there are indicated that skipjack tuna gill meal in ration has no significantly effect on broiler carcass, but it can replace the function of fish meal in the ration.

Tuna fish bone (Katsuwonus pelamis L) as well as animal bones in general, containing intercellular substance (about 70%) and the 30% remains, as inorganic salts of organic matrix. 95% of the organic component is formed from collagen, the basic substanct of the remainder consists of proteoglycan and non-collagen molecules that the pear to be involved in the regulation of bone mineralization. Collagen in the bone is approximately half of the total body collagen, and its structure is similar to the collagen in other tissues binder. Almost all of it, is type I fiber. Three-dimensional space on the structure called hole zones, are home to deposits of minerals.

The contribution of the basic substance of proteoglycan in bone has much smaller proportion than in the cartilage, mainly composed of chondroitin sulphate and hyaluronic acid. The basic substances that controlling water content in the bone, and possibly involved in regulating collagen fiber formation. Non-collagen organic material consists of osteocalcin (osla proteins) involved in the binding of calcium during the mineralization process, osteonectin that serves as a bridge between collagen and mineral components, sialoprotein (rich in salicylic acid) and some other proteins. Inorganic matrix is minerals which consists mainly of calcium and phosphate in the form of hydroxyapatite crystals. The crystals are arranged along the collagen fibers. Bone hardness depends on the concentration of inorganic material in the matrix, while the strength depends on the organic materials, particularly collagen fibers.

Collagen is part of the protein fibers or fibrous proteins that have multiple polypeptide chains linked by a variety of crosslinking to form *a triple helix*. Collagen is the protein part of the stromal type. These proteins can not be extracted with water, diluted acid, alkali or salt solution at a concentration 0.01 to 0.1. Collagen can expand because of its molecular tissue structure weakened because of the treatment given when the pH below 4 or increased to 10.

In the manufacture of bone gelatin or the use of bone as a source of minerals then the fat and non-collagen protein should be reduced to the minimum, deproteinization is a process that aims to eliminate or dissolve the protein as much as possible from the substrate, usually done by using a chemical solution that is alkaline (Suryani et al., 2005).

MATERIALS AND METHOD

Materials


Materials are solid waste processing tuna fish bone in the form of components, and the chemicals used were NaOH solution. The tools are: scales, knives, waste container.

Research Procedure

Deproteinization carried out with sodium hydroxide solution, at various concentrations and longer processing time. Before deproteinization with NaOH washing, reduction of bone size and cooking in boiling water to facilitate spending on the remnants of meat (non-collagen protein), and the layers that contain fat deposits. Removing of fat from the bone tissue (degreasing), conducted at a temperature between the melting point of fat and bone albumin coagulation (between 32-80°C) to produce an optimum fat solubility (Junianto, et al, 2006).

Research Methods

This study uses Factorial Completely Randomized Design Patterns 3 X 4 replications 3 times where the factors are : A = the concentration of alkali (given the symbol K) and B = processing time (given the symbol W). Factor A: the concentration of NaOH ($K_1 = 2\%$, $K_2 = 3\%$; $K_3 = 4\%$) Factor B: Processing Time ($W_1 = 12$ hours; $W_2 = 24$ hours; $W_3 = 36$ hours; $W_4 = 48$ hours).



RESULTS AND DISCUSSION

Research on the use of NaOH as a source of lye to changes in bone protein content of tuna (Katsuwonus Pelamis L) shown in Figure 1.




Figure 1. Effect of treatment on protein content (%)Skipjack Bone (Katsuwonus pelamis L).

In Table 1, shows that the combination treatment of K_4W_4 (NaOH concentration of 4% with hydrolysis time 48 h) gives the most highly hydrolyzed protein as indicated by the lowest protein content of 7.71%, while the protein hydrolyzed the most. The lowest is a combination treatment of K_1W_1 (concentration of 2% NaOH with hydrolysis time of 12 hours) that still contains the highest protein of 12.70%.

Table 1.Duncan Test Results (Effect of NaOH Concentration on Protein Levels of Bone Skipjack (Katsuwonus pelamis L).

| Treatment | The average protein content | Significance (0.05) |
|----------------|-----------------------------|---------------------|
| | % | |
| K ₁ | 12:24 | a |
| K ₂ | 10.97 | b |
| K 3 | 8.64 | С |

Information.: Different letters indicated significance in columns (P < 0.05)

K $_1$ = Concentration of NaOH 2%, K $_2$ = concentration of NaOH 3%; K $_3$ = Concentration of NaOH 4%.

By analysis of vasance, showed that the concentration of NaOH and the duration of hydrolysis time, has significant (P <0.01) effect on protein content of tuna fish bone, means that protein content changes because the concentration and the duration of hydrolysis by NaOH. Furthermore, to know the difference between treatments is using Duncan's test, are presented in Table 2.

Based on Duncan's test data in Table 2, K_3 treatment (concentration of NaOH 4%) gives lowest bone protein content (8.64%). This means that the higher concentration of NaOH will cause more hydrolizable Skipjack (*Katsuwonus pelamis* L) bone protein. The results are consistent with the opinion of Stevens and Verhe (2004) that the protein

components in the form of bone collagen will be hydrolyzed if the pH is increased up to pH 10.

Table 2. Duncan Test Results of Hydrolysis Time with NaOH against Skipjack (*Katsuwonus pelamis* L) Bone Protein Levels.

| | % | |
|----------------|-------|----|
| W ₁ | 11.76 | a |
| W ₂ | 10.94 | a |
| W ₃ | 10:05 | ab |
| W ₄ | 9.70 | b |

Informatan: Different letters indicated significancy in columns (P < 0.05)

 $W_1 = 12$ Hours Hydrolysis, $W_2 = 24$ Hours Hydrolysis, $W_3 = 36$ Hours Hydrolysis;

W₄ = 48 Hours Hydrolysis

In Table 2, according to Duncan test, W_4 (48 hours) has the lowest levels of bone protein (9.70%) this means that the length of hydrolysis time, would result more hydrolizable Skipjack (*Katsuwonus pelamis* L) bone protein. This results are similar with Bagau (2010), that using of 4% NaOH solution concentration for 24 hours at room temperature, dried and milled, processed tuna bone still has 10.29% protein content.

CONCLUSION

Results and discussion indicated, the influence of concentration and time of deproteinization NaOH hydrolysis, as measured by the levels of hydrolizable bone protein, can be concluded that the concentration of NaOH 4% with 48 hours hydrolysis time, give the lowest of bone proteins tuna (*Katsuwonus pelamis* L).

REFERENCES

- 1. Bagau, B., 2010. Pengolahan Limbah Padat Ikan Cakalang (*Katsuwonus pelamis* L) pagan HCl dan NaOH. UNPAD. Bandung.
- Junianto, Kiki H, Ine M, 2006. Produksi Gelatin dari Tulang Ikan dan Pemanfaatannya Sebagai Bahan Pembuatan Cangkang Kapsul. Laboran Penelitian Hibah Bersaing. Depatermen Pendidikan Tinggi.
- 3. Lengkey, Hendronoto A.W., Tuti Widjastuti, Maya Ludong, 2011. Various levels effect of Skipjack tuna gill meal (Katsowunus pelamos L) in ration on broiler car and abdominal fat. Lucrari Stiintifice Seria Zootehnie Vol. 55 (16), Editura Ion Ionescu de La Brad, Iasi, Romania. p 237-239.
- 4. Prih, S., A. Poernomo, D. Suryaningrum dan S, Ilyas, 1982. Evaluasi Kandungan Potein Ikan Indonesia. Laporan Penelitian Teknologi Perikanan DEPTAN. Jakarta.
- 5. Stevens, C.V and Verhe, R.G. (2004) Renewable Bioresources, Scope and Modification for Non-Food Applications, John Wiley & Sons Ltd, Chichester
- Suryanti, S. Hadi, R. Paranginangin, 2006. Ekstraksi Gelatin dari Tulang Ikan Kakap Merah (*Lutjanus sp*) Secara Asam. Jurnal Pascapanen dan Bioteknologi Kelautan dan Perikanan Vol. 1 No. 1.

BONE DEPROTEINIZATION SKIPJACK (Katsuwonus pelamis L) WITH NaOH IN CONCENTRATION AND TIME DIFFERENCE HYDROLISIS

| | RULISIS | | | |
|---------|--------------------------------|--|--|---------------------|
| ORIGINA | ALITY REPORT | | | |
| SIMILA | 4% ARITY INDEX | 10% INTERNET SOURCES | 6% PUBLICATIONS | % STUDENT PAPERS |
| PRIMAR | Y SOURCES | | | |
| 1 | bsclupa Internet Sour | n.asm.md | | 4% |
| 2 | id.123de Internet Sour | | | 1 % |
| 3 | etd.repo | ository.ugm.ac. | id | 1 % |
| 4 | jurnal.u Internet Sour | ntan.ac.id | | 1 % |
| 5 | develop concent ovulatin | A Atanasov. "Ement and estra crations in non- ng post-partum es", Reproductions, 2018 | ndiol-17β ovulating and Bulgarian Mu | ¶ % rrah |
| 6 | energyf Internet Sour | romwasteandw rce | ood.weebly.co | 1 % |
| 7 | Babeș-E | Bolyai University | / | 1 % |

| 8 | W Atmaka, B Yudhistira, M I S Putro. " Characteristic study of chitosan addition in Tilapia () bone based gelatin film ", IOP Conference Series: Earth and Environmental Science, 2018 Publication | 1 % |
|----|--|-----|
| 9 | sinta3.ristekdikti.go.id Internet Source | 1 % |
| 10 | Lobna Badawi, Gaber Ibrahim, Ahmed Ali, Attia Omar. "The Effect of Using Moringa (Moringa oleifera lam.) Leaves as Feed Additives on Growth Performance, Feed Utilization and Immune Response of Red Tilapia (Oreochromis sp.) Fingerlings", Sinai Journal of Applied Sciences, 2021 | 1 % |
| 11 | Guo, Chun-Yuan. "BIOCHEMICAL MARKERS OF BONE TURNOVER: ASSAY METHODS AND CLINICAL APPLICATION", Current Topics in Osteoporosis, 2005. | 1 % |
| 12 | Şt. Tudose-Sandu-Ville, V.V. Cotea, Cintia Colibaba, B. Nechita, M. Niculaua, Maria Codreanu. "Phenolic Compounds in Merlot Wines Obtained Through Different Technologies in Iaşi Vineyard, Romania", Cercetari agronomice in Moldova, 2012 Publication | 1 % |

| T | 13 |
|----|----|
| W | |
| ار | |

Taghreed Ibrahim. "Diseases of Nile Tilapia with Special Emphasis on Water Pollution", Journal of Environmental Science and Technology, 2019

<1%

Publication

14

Musa Yapen, Bertie Elias Kaseger, Nurmeilita Taher. "ANALISA KADAR AIR DAN UJI ORGANOLEPTIK PADA IKAN MUJAIR (Oreochcromis mossambicus) DI PASAR BRESEHATI", MEDIA TEKNOLOGI HASIL PERIKANAN, 2016

<1%

Publication



repository.sustech.edu

Internet Source

<1%

Exclude quotes Off
Exclude bibliography Off

Exclude matches

Off